Cristobal C Richart

List of Publications by Year in Descending Order

Source: https://exaly.com/author-pdf/7589363/cristobal-c-richart-publications-by-year.pdf

Version: 2024-04-10

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

62 4,200 94 33 h-index g-index citations papers 4,652 5.8 100 4.35 avg, IF L-index ext. citations ext. papers

#	Paper	IF	Citations
94	Lipocalin, Resistin and Gut Microbiota-Derived Propionate Could Be Used to Predict Metabolic Bariatric Surgery Selected Outcomes. <i>Processes</i> , 2022 , 10, 143	2.9	
93	The Potential Protective Role of RUNX1 in Nonalcoholic Fatty Liver Disease. <i>International Journal of Molecular Sciences</i> , 2021 , 22,	6.3	1
92	Relationship between IL-8 Circulating Levels and TLR2 Hepatic Expression in Women with Morbid Obesity and Nonalcoholic Steatohepatitis. <i>International Journal of Molecular Sciences</i> , 2020 , 21,	6.3	9
91	Hepatocyte Notch Signaling Deregulation Related to Lipid Metabolism in Women with Obesity and Nonalcoholic Fatty Liver. <i>Obesity</i> , 2020 , 28, 1487-1493	8	2
90	Circulating microbiota-derived metabolites: a "liquid biopsy?. <i>International Journal of Obesity</i> , 2020 , 44, 875-885	5.5	30
89	Low Circulating Levels of Neurotensin in Women with Nonalcoholic Fatty Liver Disease Associated with Severe Obesity. <i>Obesity</i> , 2018 , 26, 274-278	8	12
88	Targeted metabolomic approach in men with carotid plaque. PLoS ONE, 2018, 13, e0200547	3.7	8
87	Hepcidin in morbidly obese women with non-alcoholic fatty liver disease. PLoS ONE, 2017, 12, e018706.	53.7	13
86	Adipo/cytokines in atherosclerotic secretomes: increased visfatin levels in unstable carotid plaque. <i>BMC Cardiovascular Disorders</i> , 2016 , 16, 149	2.3	17
85	Redundant roles of the phosphatidate phosphatase family in triacylglycerol synthesis in human adipocytes. <i>Diabetologia</i> , 2016 , 59, 1985-94	10.3	22
84	Proteomic Profile of Unstable Atheroma Plaque: Increased Neutrophil Defensin 1, Clusterin, and Apolipoprotein E Levels in Carotid Secretome. <i>Journal of Proteome Research</i> , 2016 , 15, 933-44	5.6	19
83	Increased Circulating Levels of Alpha-Ketoglutarate in Morbidly Obese Women with Non-Alcoholic Fatty Liver Disease. <i>PLoS ONE</i> , 2016 , 11, e0154601	3.7	7
82	miR33a/miR33b* and miR122 as Possible Contributors to Hepatic Lipid Metabolism in Obese Women with Nonalcoholic Fatty Liver Disease. <i>International Journal of Molecular Sciences</i> , 2016 , 17,	6.3	30
81	PNPLA3 Expression Is Related to Liver Steatosis in Morbidly Obese Women with Non-Alcoholic Fatty Liver Disease. <i>International Journal of Molecular Sciences</i> , 2016 , 17,	6.3	17
80	Interleukin-17A Gene Expression in Morbidly Obese Women. <i>International Journal of Molecular Sciences</i> , 2015 , 16, 17469-81	6.3	18
79	Downregulation of de Novo Fatty Acid Synthesis in Subcutaneous Adipose Tissue of Moderately Obese Women. <i>International Journal of Molecular Sciences</i> , 2015 , 16, 29911-22	6.3	11
78	Evaluation of the pharmacogenetics of immune recovery in treated HIV-infected patients. <i>Expert Opinion on Drug Metabolism and Toxicology</i> , 2014 , 10, 81-101	5.5	6

(2011-2014)

77	Endocannabinoid receptors gene expression in morbidly obese women with nonalcoholic fatty liver disease. <i>BioMed Research International</i> , 2014 , 2014, 502542	3	21
76	Altered fatty acid metabolism-related gene expression in liver from morbidly obese women with non-alcoholic fatty liver disease. <i>International Journal of Molecular Sciences</i> , 2014 , 15, 22173-87	6.3	37
75	Clinical and adipocytokine changes after bariatric surgery in morbidly obese women. <i>Obesity</i> , 2014 , 22, 188-94	8	20
74	Downregulation of lipogenesis and fatty acid oxidation in the subcutaneous adipose tissue of morbidly obese women. <i>Obesity</i> , 2014 , 22, 2032-8	8	25
73	Plasma visfatin levels and gene expression in morbidly obese women with associated fatty liver disease. <i>Clinical Biochemistry</i> , 2013 , 46, 202-8	3.5	49
72	Long-term changes in leptin, chemerin and ghrelin levels following different bariatric surgery procedures: Roux-en-Y gastric bypass and sleeve gastrectomy. <i>Obesity Surgery</i> , 2013 , 23, 1790-8	3.7	81
71	Retinol binding protein-4 circulating levels were higher in nonalcoholic fatty liver disease vs. histologically normal liver from morbidly obese women. <i>Obesity</i> , 2013 , 21, 170-7	8	43
70	Adipocytokine levels in women with anorexia nervosa. Relationship with weight restoration and disease duration. <i>International Journal of Eating Disorders</i> , 2013 , 46, 855-61	6.3	29
69	Omega-3 docosahexaenoic acid and procyanidins inhibit cyclo-oxygenase activity and attenuate NF- B activation through a p105/p50 regulatory mechanism in macrophage inflammation. <i>Biochemical Journal</i> , 2012 , 441, 653-63	3.8	48
68	Increased levels and adipose tissue expression of visfatin in morbidly obese women: the relationship with pro-inflammatory cytokines. <i>Clinical Endocrinology</i> , 2012 , 77, 691-8	3.4	41
67	Leptin and adiponectin, but not IL18, are related with insulin resistance in treated HIV-1-infected patients with lipodystrophy. <i>Cytokine</i> , 2012 , 58, 253-60	4	22
66	Adipogenic/lipid, inflammatory, and mitochondrial parameters in subcutaneous adipose tissue of untreated HIV-1-infected long-term nonprogressors: significant alterations despite low viral burden. <i>Journal of Acquired Immune Deficiency Syndromes (1999)</i> , 2012 , 61, 131-7	3.1	18
65	Mirror extreme BMI phenotypes associated with gene dosage at the chromosome 16p11.2 locus. <i>Nature</i> , 2011 , 478, 97-102	50.4	322
64	Lipodystrophy and insulin resistance in combination antiretroviral treated HIV-1-infected patients: implication of resistin. <i>Journal of Acquired Immune Deficiency Syndromes (1999)</i> , 2011 , 57, 16-23	3.1	19
63	Upregulation of lipocalin 2 in adipose tissues of severely obese women: positive relationship with proinflammatory cytokines. <i>Obesity</i> , 2011 , 19, 2295-300	8	64
62	New adipokines vaspin and omentin. Circulating levels and gene expression in adipose tissue from morbidly obese women. <i>BMC Medical Genetics</i> , 2011 , 12, 60	2.1	113
61	No relationship between TNF-Igenetic variants and combination antiretroviral therapy-related lipodystrophy syndrome in HIV type 1-infected patients: a case-control study and a meta-analysis. <i>AIDS Research and Human Retroviruses</i> , 2011 , 27, 143-52	1.6	9
60	FABP 4 is associated with inflammatory markers and metabolic syndrome in morbidly obese women. <i>European Journal of Endocrinology</i> , 2011 , 164, 539-47	6.5	88

59	Anti-inflammatory profile of FTO gene expression in adipose tissues from morbidly obese women. <i>Cellular Physiology and Biochemistry</i> , 2010 , 26, 1041-50	3.9	25
58	Association of retinol-binding protein-4 (RBP4) with lipid parameters in obese women. <i>Obesity Surgery</i> , 2010 , 20, 1258-64	3.7	35
57	Polymorphisms in the interleukin-10 gene promoter and the risk of alcoholism and alcoholic liver disease in Caucasian Spaniard men. <i>Alcohol</i> , 2010 , 44, 211-6	2.7	8
56	Effect of TNF-alpha genetic variants and CCR5 Delta 32 on the vulnerability to HIV-1 infection and disease progression in Caucasian Spaniards. <i>BMC Medical Genetics</i> , 2010 , 11, 63	2.1	18
55	Parallel downregulation of retinol-binding protein-4 and adiponectin expression in subcutaneous adipose tissue of non-morbidly obese subjects. <i>European Journal of Endocrinology</i> , 2009 , 161, 87-94	6.5	12
54	Obesity and Critical Care. Clinical Pulmonary Medicine, 2009, 16, 202-211	0.3	3
53	PPARgamma Pro12Ala polymorphism in HIV-1-infected patients with HAART-related lipodystrophy. <i>Current HIV Research</i> , 2009 , 7, 533-40	1.3	12
52	A study on the TNF-alpha system in Caucasian Spanish patients with alcoholic liver disease. <i>Drug and Alcohol Dependence</i> , 2008 , 92, 91-9	4.9	22
51	The IL-6 system in HIV-1-infection and in HAART-related fat redistribution syndromes. <i>Aids</i> , 2008 , 22, 893-6	3.5	18
50	Human subcutaneous adipose tissue LPIN1 expression in obesity, type 2 diabetes mellitus, and human immunodeficiency virusassociated lipodystrophy syndrome. <i>Metabolism: Clinical and Experimental</i> , 2007 , 56, 1518-26	12.7	20
49	Monocyte chemoattractant protein-1 in obesity and type 2 diabetes. Insulin sensitivity study. <i>Obesity</i> , 2007 , 15, 664-72	8	40
48	Circulating retinol-binding protein-4, insulin sensitivity, insulin secretion, and insulin disposition index in obese and nonobese subjects. <i>Diabetes Care</i> , 2007 , 30, 1802-6	14.6	118
47	LMNA messenger RNA expression in highly active antiretroviral therapy-treated HIV-positive patients. <i>Journal of Acquired Immune Deficiency Syndromes</i> (1999), 2007 , 46, 384-9	3.1	8
46	Lamivudine in the treatment of hepatitis B virus-related polyarteritis nodosa. <i>Journal of Clinical Rheumatology</i> , 2007 , 13, 298-9	1.1	5
45	Effect of genetic variants of CCR2 and CCL2 on the natural history of HIV-1 infection: CCL2-2518GG is overrepresented in a cohort of Spanish HIV-1-infected subjects. <i>Journal of Acquired Immune Deficiency Syndromes (1999)</i> , 2007 , 44, 132-8	3.1	24
44	Clinical Characteristics and Outcome of Infective Endocarditis in Individuals of the General Population Managed at a Teaching Hospital Without Cardiac Surgery Facilities. Study of 120 Cases. <i>Revista Espanola De Cardiologia (English Ed)</i> , 2006 , 59, 1131-1139	0.7	О
43	Polymorphisms of alcohol-metabolizing enzymes and the risk for alcoholism and alcoholic liver disease in Caucasian Spanish women. <i>Drug and Alcohol Dependence</i> , 2006 , 84, 195-200	4.9	29
42	Polymorphism of RANTES chemokine gene promoter is not associated with long-term nonprogressive HIV-1 infection of more than 16 years. <i>Journal of Acquired Immune Deficiency Syndromes</i> (1999), 2006 , 41, 17-22	3.1	21

(2002-2006)

41	Absence of CXCR4 C-terminal polymorphisms in HIV-1-infected and uninfected Spaniards. <i>Journal of Acquired Immune Deficiency Syndromes (1999)</i> , 2006 , 42, 382-3	3.1	2
40	Polymorphisms in the interleukin-6 receptor gene are associated with body mass index and with characteristics of the metabolic syndrome. <i>Clinical Endocrinology</i> , 2006 , 65, 88-91	3.4	41
39	Spanish HIV-1-infected long-term nonprogressors of more than 15 years have an increased frequency of the CX3CR1 249I variant allele. <i>Journal of Acquired Immune Deficiency Syndromes</i> (1999), 2005 , 40, 527-31	3.1	35
38	Lack of association of SDF-1 35A variant allele with long-term nonprogressive HIV-1 infection is extended beyond 16 years. <i>Journal of Acquired Immune Deficiency Syndromes (1999)</i> , 2005 , 40, 276-9	3.1	16
37	Mannose-binding lectin gene polymorphisms are associated with gestational diabetes mellitus. Journal of Clinical Endocrinology and Metabolism, 2004 , 89, 5081-7	5.6	44
36	Proximal tubular kidney damage and tenofovir: a role for mitochondrial toxicity?. Aids, 2004, 18, 1741-2	3.5	26
35	Circulating soluble CD14 monocyte receptor is associated with increased alanine aminotransferase. <i>Clinical Chemistry</i> , 2004 , 50, 1456-8	5.5	9
34	Resistin, adiponectin, ghrelin, leptin, and proinflammatory cytokines: relationships in obesity. <i>Obesity</i> , 2004 , 12, 962-71		362
33	Genetic polymorphisms of ADH2, ADH3, CYP4502E1 Dra-I and Pst-I, and ALDH2 in Spanish men: lack of association with alcoholism and alcoholic liver disease. <i>Journal of Hepatology</i> , 2004 , 41, 744-50	13.4	66
32	Substitution of nevirapine, efavirenz, or abacavir for protease inhibitors in patients with human immunodeficiency virus infection. <i>New England Journal of Medicine</i> , 2003 , 349, 1036-46	59.2	268
31	CD14 monocyte receptor, involved in the inflammatory cascade, and insulin sensitivity. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2003 , 88, 1780-4	5.6	79
30	Severe neutropenia and thrombocytopenia associated with infliximab. <i>Annals of Internal Medicine</i> , 2003 , 139, W-W63	8	50
29	The interleukin-6 (-174) G/C promoter polymorphism is associated with type-2 diabetes mellitus in Native Americans and Caucasians. <i>Human Genetics</i> , 2003 , 112, 409-13	6.3	145
28	A polymorphism in the promoter of the tumor necrosis factor-alpha gene (-308) is associated with coronary heart disease in type 2 diabetic patients. <i>Atherosclerosis</i> , 2003 , 167, 257-64	3.1	93
27	G protein beta3 gene variant, vascular function, and insulin sensitivity in type 2 diabetes. Hypertension, 2003 , 41, 124-9	8.5	26
26	Shedding of TNF-alpha receptors, blood pressure, and insulin sensitivity in type 2 diabetes mellitus. <i>American Journal of Physiology - Endocrinology and Metabolism</i> , 2002 , 282, E952-9	6	57
25	Paraoxonase Gln-Arg(192) and Leu-Met(55) gene polymorphisms and enzyme activity in a population with a low rate of coronary heart disease. <i>Clinical Biochemistry</i> , 2002 , 35, 197-203	3.5	39
24	Progressive multifocal leukoencephalopathy in chronic lymphocytic leukemia after treatment with fludarabine. <i>Leukemia and Lymphoma</i> , 2002 , 43, 433-6	1.9	42

23	Platelet count and interleukin 6 gene polymorphism in healthy subjects. <i>BMC Medical Genetics</i> , 2001 , 2, 6	2.1	18
22	Circulating interleukin 6 levels, blood pressure, and insulin sensitivity in apparently healthy men and women. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2001 , 86, 1154-9	5.6	418
21	Genetic polymorphism of alcohol dehydrogenase in europeans: the ADH2*2 allele decreases the risk for alcoholism and is associated with ADH3*1. <i>Hepatology</i> , 2000 , 31, 984-9	11.2	201
20	Left ventricular hypertrophy in hypertensive patients with autosomal dominant polycystic kidney disease: influence of blood pressure and humoral and neurohormonal factors. <i>American Journal of Nephrology</i> , 2000 , 20, 193-200	4.6	31
19	Echocardiographic evaluation in patients with autosomal dominant polycystic kidney disease and end-stage renal disease. <i>American Journal of Kidney Diseases</i> , 1999 , 34, 264-72	7.4	4
18	Asthma, mite sensitization, and sleeping in bunks. <i>Annals of Allergy, Asthma and Immunology</i> , 1999 , 82, 531-3	3.2	6
17	Plasma levels of the soluble fraction of tumor necrosis factor receptors 1 and 2 are independent determinants of plasma cholesterol and LDL-cholesterol concentrations in healthy subjects. <i>Atherosclerosis</i> , 1999 , 146, 321-7	3.1	40
16	Genetic heterogeneity within ulcerative colitis determined by an interleukin-1 receptor antagonist gene polymorphism and antineutrophil cytoplasmic antibodies. <i>European Journal of Gastroenterology and Hepatology</i> , 1999 , 11, 413-20	2.2	21
15	Ambulatory blood pressure and left ventricular mass in normotensive patients with autosomal dominant polycystic kidney disease. <i>Journal of the American Society of Nephrology: JASN</i> , 1999 , 10, 1020)- [2·7	52
14	Influence of chronic alcohol abuse and liver disease on hepatic aldehyde dehydrogenase activity. <i>Alcohol</i> , 1998 , 15, 3-8	2.7	29
13	Lipoprotein(a) and the significance of the association between platelet glycoprotein IIIa polymorphisms and the risk of premature myocardial infarction. <i>Atherosclerosis</i> , 1998 , 140, 155-9	3.1	33
12	Recovery of minimal change nephrotic syndrome and acute renal failure in a patient withRenal cell carcinoma. <i>American Journal of Nephrology</i> , 1998 , 18, 433-5	4.6	14
11	Fatal Listeria meningitis, endocarditis and pericarditis in a patient with haemochromatosis. Scandinavian Journal of Infectious Diseases, 1997 , 29, 308-9		25
10	Alcohol dehydrogenase of human and rat blood vessels. Role in ethanol metabolism. <i>FEBS Letters</i> , 1997 , 405, 26-30	3.8	47
9	Minimal-change glomerulopathy and carcinoma. Report of two cases and review of the literature. <i>American Journal of Nephrology</i> , 1993 , 13, 69-72	4.6	19
8	Granulomatous hepatitis and fever of unknown origin: influence of AIDS epidemics. <i>Journal of Clinical Gastroenterology</i> , 1993 , 17, 89	3	3
7	Decrease of skin and bronchial sensitization following short-intensive scheduled immunotherapy in mite-allergic asthma. <i>Chest</i> , 1993 , 103, 183-7	5.3	18
6	Atypical liver alcohol dehydrogenase in the Spanish population: its relation with the development of alcoholic liver disease. <i>Alcoholism: Clinical and Experimental Research</i> , 1993 , 17, 782-5	3.7	12

LIST OF PUBLICATIONS

5	Effect of ascorbic acid on gastrointestinal aluminium absorption. <i>Lancet, The</i> , 1991 , 338, 1467	40	22
4	IgA nephropathy and polycystic kidney disease. American Journal of Nephrology, 1988, 8, 477-8	4.6	67
3	Systemic vasculitis resembling periarteritis nodosa in the lupus-like syndrome induced by hydralazine. <i>American Journal of Nephrology</i> , 1987 , 7, 71-3	4.6	17
2	More on unusual forms of meningococcal disease: septic polyarthritis. <i>American Journal of Medicine</i> , 1986 , 81, 561-2	2.4	2
1	Retinol Binding Protein-4 Circulating Levels Were Higher in Nonalcoholic Fatty Liver Disease Vs. Histologically Normal Liver From Morbidly Obese Women. <i>Obesity</i> ,	8	4