

Monika E Hegi

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/7590806/monika-e-hegi-publications-by-year.pdf>

Version: 2024-04-28

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

141
papers

28,129
citations

68
h-index

167
g-index

168
ext. papers

33,326
ext. citations

8.8
avg, IF

6.58
L-index

#	Paper	IF	Citations
141	DNA methylation-based age acceleration observed in IDH wild-type glioblastoma is associated with better outcome-including in elderly patients.. <i>Acta Neuropathologica Communications</i> , 2022 , 10, 39	7.3	1
140	Compensatory CSF2-driven macrophage activation promotes adaptive resistance to CSF1R inhibition in breast-to-brain metastasis.. <i>Nature Cancer</i> , 2021 , 2, 1086-1101	15.4	6
139	Hyperpolarized C-glucose magnetic resonance highlights reduced aerobic glycolysis in vivo in infiltrative glioblastoma. <i>Scientific Reports</i> , 2021 , 11, 5771	4.9	4
138	BET inhibitors repress expression of interferon-stimulated genes and synergize with HDAC inhibitors in glioblastoma. <i>Neuro-Oncology</i> , 2021 , 23, 1680-1692	1	5
137	EANO guidelines on the diagnosis and treatment of diffuse gliomas of adulthood. <i>Nature Reviews Clinical Oncology</i> , 2021 , 18, 170-186	19.4	204
136	The state of neuro-oncology during the COVID-19 pandemic: a worldwide assessment. <i>Neuro-Oncology Advances</i> , 2021 , 3, vdab035	0.9	0
135	Metabolic and transcriptomic profiles of glioblastoma invasion revealed by comparisons between patients and corresponding orthotopic xenografts in mice. <i>Acta Neuropathologica Communications</i> , 2021 , 9, 133	7.3	1
134	An integrated pipeline for comprehensive analysis of immune cells in human brain tumor clinical samples. <i>Nature Protocols</i> , 2021 , 16, 4692-4721	18.8	2
133	Diagnostic Surveillance of High-Grade Gliomas: Towards Automated Change Detection Using Radiology Report Classification. <i>Communications in Computer and Information Science</i> , 2021 , 423-436	0.3	
132	Interrogation of the Microenvironmental Landscape in Brain Tumors Reveals Disease-Specific Alterations of Immune Cells. <i>Cell</i> , 2020 , 181, 1643-1660.e17	56.2	200
131	Glioblastoma in adults: a Society for Neuro-Oncology (SNO) and European Society of Neuro-Oncology (EANO) consensus review on current management and future directions. <i>Neuro-Oncology</i> , 2020 , 22, 1073-1113	1	178
130	CBIO-13. EPIGENETIC DEREGLATION OF NUCLEAR TRANSLOCATION - A NOVEL MECHANISM FOR TREATMENT RESISTANCE IN GLIOBLASTOMA. <i>Neuro-Oncology</i> , 2020 , 22, ii18-ii18	1	
129	COVID-31. THE STATE OF NEURO-ONCOLOGY DURING THE COVID-19 PANDEMIC: A WORLDWIDE ASSESSMENT. <i>Neuro-Oncology</i> , 2020 , 22, ii27-ii27	1	
128	The TICKing clock of EGFR therapy resistance in glioblastoma: Target Independence or target Compensation. <i>Drug Resistance Updates</i> , 2019 , 43, 29-37	23.2	19
127	Mechanisms of Resistance to EGFR Inhibition Reveal Metabolic Vulnerabilities in Human GBM. <i>Molecular Cancer Therapeutics</i> , 2019 , 18, 1565-1576	6.1	6
126	Improving survival in molecularly selected glioblastoma. <i>Lancet, The</i> , 2019 , 393, 615-617	40	20
125	MGMT promoter methylation status testing to guide therapy for glioblastoma: refining the approach based on emerging evidence and current challenges. <i>Neuro-Oncology</i> , 2019 , 21, 167-178	1	92

124	Promoter Methylation Cutoff with Safety Margin for Selecting Glioblastoma Patients into Trials Omitting Temozolomide: A Pooled Analysis of Four Clinical Trials. <i>Clinical Cancer Research</i> , 2019 , 25, 1809-1816	12.0	59
123	Novel, improved grading system(s) for IDH-mutant astrocytic gliomas. <i>Acta Neuropathologica</i> , 2018 , 136, 153-166	14.3	162
122	The DNA methylome of DDR genes and benefit from RT or TMZ in IDH mutant low-grade glioma treated in EORTC 22033. <i>Acta Neuropathologica</i> , 2018 , 135, 601-615	14.3	54
121	In vivo characterization of brain metabolism by H MRS, C MRS and FDG PET reveals significant glucose oxidation of invasively growing glioma cells. <i>International Journal of Cancer</i> , 2018 , 143, 127-138	7.5	12
120	Toward methylation-based classification of central nervous system tumors. <i>Neuro-Oncology</i> , 2018 , 20, 579-581	1	1
119	Glioma epigenetics: From subclassification to novel treatment options. <i>Seminars in Cancer Biology</i> , 2018 , 51, 50-58	12.7	197
118	European Association for Neuro-Oncology (EANO) guideline on the diagnosis and treatment of adult astrocytic and oligodendroglial gliomas. <i>Lancet Oncology, The</i> , 2017 , 18, e315-e329	21.7	599
117	Correlation of immune phenotype with IDH mutation in diffuse glioma. <i>Neuro-Oncology</i> , 2017 , 19, 1460-1468		144
116	Is more better? The impact of extended adjuvant temozolomide in newly diagnosed glioblastoma: a secondary analysis of EORTC and NRG Oncology/RTOG. <i>Neuro-Oncology</i> , 2017 , 19, 1119-1126	1	82
115	Evidence-based management of adult patients with diffuse glioma - AuthorsReply. <i>Lancet Oncology, The</i> , 2017 , 18, e430-e431	21.7	4
114	Survey on current practice within the European Low-Grade Glioma Network: where do we stand and what is the next step?. <i>Neuro-Oncology Practice</i> , 2017 , 4, 241-247	2.2	11
113	Effect of Tumor-Treating Fields Plus Maintenance Temozolomide vs Maintenance Temozolomide Alone on Survival in Patients With Glioblastoma: A Randomized Clinical Trial. <i>JAMA - Journal of the American Medical Association</i> , 2017 , 318, 2306-2316	27.4	875
112	Ubiquitin Specific Peptidase 15 (USP15) suppresses glioblastoma cell growth via stabilization of HECTD1 E3 ligase attenuating WNT pathway activity. <i>Oncotarget</i> , 2017 , 8, 110490-110502	3.3	18
111	WIF1 re-expression in glioblastoma inhibits migration through attenuation of non-canonical WNT signaling by downregulating the lncRNA MALAT1. <i>Oncogene</i> , 2016 , 35, 12-21	9.2	101
110	Early detection of human glioma sphere xenografts in mouse brain using diffusion MRI at 14.1 T. <i>NMR in Biomedicine</i> , 2016 , 29, 1577-1589	4.4	7
109	Reply to F. Felix et al and M.F. Fay et al. <i>Journal of Clinical Oncology</i> , 2016 , 34, 3107-8	2.2	1
108	LTKB-01: PROSPECTIVE, MULTI-CENTER PHASE III TRIAL OF TUMOR TREATING FIELDS TOGETHER WITH TEMOZOLOMIDE COMPARED TO TEMOZOLOMIDE ALONE IN PATIENTS WITH NEWLY DIAGNOSED GLIOBLASTOMA. <i>Neuro-Oncology</i> , 2016 , 18, i1-i1	1	14
107	Does Valproic Acid or Levetiracetam Improve Survival in Glioblastoma? A Pooled Analysis of Prospective Clinical Trials in Newly Diagnosed Glioblastoma. <i>Journal of Clinical Oncology</i> , 2016 , 34, 731-9	2.2	108

106	Sensitivity Analysis of the MGMT-STP27 Model and Impact of Genetic and Epigenetic Context to Predict the MGMT Methylation Status in Gliomas and Other Tumors. <i>Journal of Molecular Diagnostics</i> , 2016 , 18, 350-361	5.1	54
105	Cilengitide in newly diagnosed glioblastoma: biomarker expression and outcome. <i>Oncotarget</i> , 2016 , 7, 15018-32	3.3	49
104	Current management of low-grade gliomas. <i>Current Opinion in Neurology</i> , 2016 , 29, 782-788	7.1	19
103	Phase II Study of Radiotherapy and Temozolimus versus Radiochemotherapy with Temozolomide in Patients with Newly Diagnosed Glioblastoma without MGMT Promoter Hypermethylation (EORTC 26082). <i>Clinical Cancer Research</i> , 2016 , 22, 4797-4806	12.9	77
102	Temozolomide chemotherapy versus radiotherapy in high-risk low-grade glioma (EORTC 22033-26033): a randomised, open-label, phase 3 intergroup study. <i>Lancet Oncology, The</i> , 2016 , 17, 1521-1532	21.7	294
101	Chromosome 7 gain and DNA hypermethylation at the HOXA10 locus are associated with expression of a stem cell related HOX-signature in glioblastoma. <i>Genome Biology</i> , 2015 , 16, 16	18.3	65
100	Two cilengitide regimens in combination with standard treatment for patients with newly diagnosed glioblastoma and unmethylated MGMT gene promoter: results of the open-label, controlled, randomized phase II CORE study. <i>Neuro-Oncology</i> , 2015 , 17, 708-17	1	148
99	Programmed death ligand 1 expression and tumor-infiltrating lymphocytes in glioblastoma. <i>Neuro-Oncology</i> , 2015 , 17, 1064-75	1	356
98	Genome-wide DNA methylation detection by MethylCap-seq and Infinium HumanMethylation450 BeadChips: an independent large-scale comparison. <i>Scientific Reports</i> , 2015 , 5, 15375	4.9	15
97	CSIG-05THE UBIQUITIN SPECIFIC PEPTIDASE 15 (USP15) SUPPRESSES PROLIFERATION OF HUMAN GLIOBLASTOMA CELL LINES VIA STABILIZATION OF HECTD1 E3 LIGASE. <i>Neuro-Oncology</i> , 2015 , 17, v67.1-v67	1	78
96	Mir-21-Sox2 Axis Delineates Glioblastoma Subtypes with Prognostic Impact. <i>Journal of Neuroscience</i> , 2015 , 35, 15097-112	6.6	41
95	Maintenance Therapy With Tumor-Treating Fields Plus Temozolomide vs Temozolomide Alone for Glioblastoma: A Randomized Clinical Trial. <i>JAMA - Journal of the American Medical Association</i> , 2015 , 314, 2535-43	27.4	696
94	Reply to M.C. Chamberlain. <i>Journal of Clinical Oncology</i> , 2014 , 32, 1634-5	2.2	2
93	Cilengitide combined with standard treatment for patients with newly diagnosed glioblastoma with methylated MGMT promoter (CENTRIC EORTC 26071-22072 study): a multicentre, randomised, open-label, phase 3 trial. <i>Lancet Oncology, The</i> , 2014 , 15, 1100-8	21.7	629
92	MGMT testing--the challenges for biomarker-based glioma treatment. <i>Nature Reviews Neurology</i> , 2014 , 10, 372-85	15	316
91	GENETIC AND EPIGENETIC DEREGLATION ARE ASSOCIATED WITH THE ABERRANT EXPRESSION OF A STEM CELL RELATED HOX GENE SIGNATURE IN GLIOBLASTOMA. <i>Neuro-Oncology</i> , 2014 , 16, iii8-iii8 ¹	1	78
90	EG-05COMBINATION OF GENE COPY GAIN AND EPIGENETIC DEREGLATION ARE ASSOCIATED WITH THE ABERRANT EXPRESSION OF A STEM CELL RELATED HOX-SIGNATURE IN GLIOBLASTOMA. <i>Neuro-Oncology</i> , 2014 , 16, v75-v76	1	78
89	Radiation therapy and concurrent plus adjuvant temsirolimus (CCI-779) versus chemoradiation with temozolomide in newly diagnosed glioblastoma without methylation of the MGMT gene promoter.. <i>Journal of Clinical Oncology</i> , 2014 , 32, 2003-2003	2.2	10

88	Molecular neuro-oncology in clinical practice: a new horizon. <i>Lancet Oncology, The</i> , 2013 , 14, e370-9	21.7	149
87	The DNA repair protein ALKBH2 mediates temozolomide resistance in human glioblastoma cells. <i>Neuro-Oncology</i> , 2013 , 15, 269-78	1	67
86	Dose-dense temozolomide for newly diagnosed glioblastoma: a randomized phase III clinical trial. <i>Journal of Clinical Oncology</i> , 2013 , 31, 4085-91	2.2	623
85	Epigenetics and Brain Cancer 2013 , 21-40		
84	Brain cancer in 2012: Molecular characterization leads the way. <i>Nature Reviews Clinical Oncology</i> , 2013 , 10, 69-70	19.4	11
83	Cilengitide combined with standard treatment for patients with newly diagnosed glioblastoma and methylated O6-methylguanine-DNA methyltransferase (MGMT) gene promoter: Key results of the multicenter, randomized, open-label, controlled, phase III CENTRIC study.. <i>Journal of Clinical Oncology</i> , 2013 , 31, 1512-22	2.2	24
82	EORTC 26083 phase I/II trial of dasatinib in combination with CCNU in patients with recurrent glioblastoma. <i>Neuro-Oncology</i> , 2012 , 14, 1503-10	1	45
81	Temozolomide versus standard 6-week radiotherapy versus hypofractionated radiotherapy in patients older than 60 years with glioblastoma: the Nordic randomised, phase 3 trial. <i>Lancet Oncology, The</i> , 2012 , 13, 916-26	21.7	840
80	New prognostic factors and calculators for outcome prediction in patients with recurrent glioblastoma: a pooled analysis of EORTC Brain Tumour Group phase I and II clinical trials. <i>European Journal of Cancer</i> , 2012 , 48, 1176-84	7.5	137
79	MGMT methylation analysis of glioblastoma on the Infinium methylation BeadChip identifies two distinct CpG regions associated with gene silencing and outcome, yielding a prediction model for comparisons across datasets, tumor grades, and CIMP-status. <i>Acta Neuropathologica</i> , 2012 , 124, 547-60	14.3	213
78	EORTC topics in neurooncology: The long path from a focus on neurological complications of cancer towards molecularly defined trials and therapies in neurooncology. <i>European Journal of Cancer, Supplement</i> , 2012 , 10, 20-26	1.6	
77	Treatment options in elderly patients with glioblastoma [Authors'Reply]. <i>Lancet Oncology, The</i> , 2012 , 13, e461-e462	21.7	
76	A novel volume-age-KPS (VAK) glioblastoma classification identifies a prognostic cognate microRNA-gene signature. <i>PLoS ONE</i> , 2012 , 7, e41522	3.7	70
75	A safety run-in and randomized phase 2 study of cilengitide combined with chemoradiation for newly diagnosed glioblastoma (NABTT 0306). <i>Cancer</i> , 2012 , 118, 5601-7	6.4	95
74	Clinical implications of molecular neuropathology and biomarkers for malignant glioma. <i>Current Neurology and Neuroscience Reports</i> , 2012 , 12, 302-7	6.6	20
73	Presence of an oligodendroglioma-like component in newly diagnosed glioblastoma identifies a pathogenetically heterogeneous subgroup and lacks prognostic value: central pathology review of the EORTC_26981/NCIC_CE.3 trial. <i>Acta Neuropathologica</i> , 2012 , 123, 841-52	14.3	68
72	Individualized targeted therapy for glioblastoma: fact or fiction?. <i>Cancer Journal (Sudbury, Mass)</i> , 2012 , 18, 40-4	2.2	57
71	DNA fingerprinting of glioma cell lines and considerations on similarity measurements. <i>Neuro-Oncology</i> , 2012 , 14, 701-11	1	38

70	Personalized care in neuro-oncology coming of age: why we need MGMT and 1p/19q testing for malignant glioma patients in clinical practice. <i>Neuro-Oncology</i> , 2012 , 14 Suppl 4, iv100-8	1	120
69	Epidermal growth factor receptor: a re-emerging target in glioblastoma. <i>Current Opinion in Neurology</i> , 2012 , 25, 774-9	7.1	66
68	Alkylpurine-DNA-N-glycosylase confers resistance to temozolomide in xenograft models of glioblastoma multiforme and is associated with poor survival in patients. <i>Journal of Clinical Investigation</i> , 2012 , 122, 253-66	15.9	108
67	Expression of O ⁶ -methylguanine-DNA methyltransferase in childhood medulloblastoma. <i>Journal of Neuro-Oncology</i> , 2011 , 103, 59-69	4.8	12
66	Current concepts and management of glioblastoma. <i>Annals of Neurology</i> , 2011 , 70, 9-21	9.4	319
65	The Wnt inhibitory factor 1 (WIF1) is targeted in glioblastoma and has a tumor suppressing function potentially by induction of senescence. <i>Neuro-Oncology</i> , 2011 , 13, 736-47	1	76
64	Extent and patterns of MGMT promoter methylation in glioblastoma- and respective glioblastoma-derived spheres. <i>Clinical Cancer Research</i> , 2011 , 17, 255-66	12.9	67
63	Pathway analysis of glioblastoma tissue after preoperative treatment with the EGFR tyrosine kinase inhibitor gefitinib--a phase II trial. <i>Molecular Cancer Therapeutics</i> , 2011 , 10, 1102-12	6.1	137
62	Marker-independent identification of glioma-initiating cells. <i>Nature Methods</i> , 2010 , 7, 224-8	21.6	66
61	Radiotherapy suppresses angiogenesis in mice through TGF-betaRI/ALK5-dependent inhibition of endothelial cell sprouting. <i>PLoS ONE</i> , 2010 , 5, e11084	3.7	55
60	Phase I/IIa study of cilengitide and temozolomide with concomitant radiotherapy followed by cilengitide and temozolomide maintenance therapy in patients with newly diagnosed glioblastoma. <i>Journal of Clinical Oncology</i> , 2010 , 28, 2712-8	2.2	338
59	Reply to M.C. Chamberlain. <i>Journal of Clinical Oncology</i> , 2010 , 28, e696-e697	2.2	
58	Should biomarkers be used to design personalized medicine for the treatment of glioblastoma?. <i>Future Oncology</i> , 2010 , 6, 1407-14	3.6	20
57	Presence of alternative lengthening of telomeres mechanism in patients with glioblastoma identifies a less aggressive tumor type with longer survival. <i>Journal of Neuropathology and Experimental Neurology</i> , 2010 , 69, 729-36	3.1	66
56	MGMT promoter methylation in malignant gliomas: ready for personalized medicine?. <i>Nature Reviews Neurology</i> , 2010 , 6, 39-51	15	538
55	Molecular diagnostics of gliomas: the clinical perspective. <i>Acta Neuropathologica</i> , 2010 , 120, 585-92	14.3	96
54	MGMT promoter methylation in malignant gliomas. <i>Targeted Oncology</i> , 2010 , 5, 161-5	5	52
53	Combining chromosomal arm status and significantly aberrant genomic locations reveals new cancer subtypes. <i>Cancer Informatics</i> , 2009 , 7, 91-104	2.4	

52	Modulation of angiogenic and inflammatory response in glioblastoma by hypoxia. <i>PLoS ONE</i> , 2009 , 4, e5947	3.7	81
51	MGMT promoter methylation is prognostic but not predictive for outcome to adjuvant PCV chemotherapy in anaplastic oligodendroglial tumors: a report from EORTC Brain Tumor Group Study 26951. <i>Journal of Clinical Oncology</i> , 2009 , 27, 5881-6	2.2	203
50	Tenascin-C is a novel RBPJkappa-induced target gene for Notch signaling in gliomas. <i>Cancer Research</i> , 2009 , 69, 458-65	10.1	102
49	Epigenetic deregulation of DNA repair and its potential for therapy. <i>Clinical Cancer Research</i> , 2009 , 15, 5026-31	12.9	49
48	RNOP-09: pegylated liposomal doxorubicine and prolonged temozolomide in addition to radiotherapy in newly diagnosed glioblastoma--a phase II study. <i>BMC Cancer</i> , 2009 , 9, 308	4.8	61
47	Elevated levels of MIC-1/GDF15 in the cerebrospinal fluid of patients are associated with glioblastoma and worse outcome. <i>International Journal of Cancer</i> , 2009 , 125, 2624-30	7.5	68
46	Effects of radiotherapy with concomitant and adjuvant temozolomide versus radiotherapy alone on survival in glioblastoma in a randomised phase III study: 5-year analysis of the EORTC-NCIC trial. <i>Lancet Oncology, The</i> , 2009 , 10, 459-66	21.7	5111
45	Anti-O6-methylguanine-methyltransferase (MGMT) immunohistochemistry in glioblastoma multiforme: observer variability and lack of association with patient survival impede its use as clinical biomarker. <i>Brain Pathology</i> , 2008 , 18, 520-32	6	157
44	Nomograms for predicting survival of patients with newly diagnosed glioblastoma: prognostic factor analysis of EORTC and NCIC trial 26981-22981/CE.3. <i>Lancet Oncology, The</i> , 2008 , 9, 29-38	21.7	407
43	Validation of real-time methylation-specific PCR to determine O6-methylguanine-DNA methyltransferase gene promoter methylation in glioma. <i>Journal of Molecular Diagnostics</i> , 2008 , 10, 332-7	5.1	145
42	Correlation of O6-methylguanine methyltransferase (MGMT) promoter methylation with clinical outcomes in glioblastoma and clinical strategies to modulate MGMT activity. <i>Journal of Clinical Oncology</i> , 2008 , 26, 4189-99	2.2	625
41	Combined lysophosphatidic acid/platelet-derived growth factor signaling triggers glioma cell migration in a tenascin-C microenvironment. <i>Cancer Research</i> , 2008 , 68, 6942-52	10.1	36
40	Epidermal growth factor receptor inhibitors in neuro-oncology: hopes and disappointments. <i>Clinical Cancer Research</i> , 2008 , 14, 957-60	12.9	110
39	Stem cell-related "self-renewal" signature and high epidermal growth factor receptor expression associated with resistance to concomitant chemoradiotherapy in glioblastoma. <i>Journal of Clinical Oncology</i> , 2008 , 26, 3015-24	2.2	541
38	Infrequent promoter methylation of the MGMT gene in liver metastases from uveal melanoma. <i>International Journal of Cancer</i> , 2008 , 123, 1215-8	7.5	18
37	Chemoradiotherapy in malignant glioma: standard of care and future directions. <i>Journal of Clinical Oncology</i> , 2007 , 25, 4127-36	2.2	385
36	MGMT methylation status: the advent of stratified therapy in glioblastoma?. <i>Disease Markers</i> , 2007 , 23, 97-104	3.2	39
35	Neuro-oncology: oligodendroglioma and molecular markers. <i>Lancet Neurology, The</i> , 2007 , 6, 10-2	24.1	5

34	Endothelin receptor type B counteracts tenascin-C-induced endothelin receptor type A-dependent focal adhesion and actin stress fiber disorganization. <i>Cancer Research</i> , 2007 , 67, 6163-73	10.1	49
33	Correlative studies in neuro-oncology trials: should they influence treatment?. <i>Current Oncology Reports</i> , 2006 , 8, 54-7	6.3	5
32	Histopathologic assessment of hot-spot microvessel density and vascular patterns in glioblastoma: Poor observer agreement limits clinical utility as prognostic factors: a translational research project of the European Organization for Research and Treatment of Cancer Brain Tumor Group. <i>Cancer</i> , 2006 , 107, 162-70	6.4	47
31	New trends in the medical management of glioblastoma multiforme: the role of temozolomide chemotherapy. <i>Neurosurgical Focus</i> , 2006 , 20, E6	4.2	57
30	Changing paradigms--an update on the multidisciplinary management of malignant glioma. <i>Oncologist</i> , 2006 , 11, 165-80	5.7	316
29	Recent developments in the use of chemotherapy in brain tumours. <i>European Journal of Cancer</i> , 2006 , 42, 582-8	7.5	95
28	MGMT gene silencing and benefit from temozolomide in glioblastoma. <i>New England Journal of Medicine</i> , 2005 , 352, 997-1003	59.2	5267
27	Optimal role of temozolomide in the treatment of malignant gliomas. <i>Current Neurology and Neuroscience Reports</i> , 2005 , 5, 198-206	6.6	142
26	Genetic alterations in brain tumors following 1,3-butadiene exposure in B6C3F1 mice. <i>Toxicologic Pathology</i> , 2005 , 33, 307-12	2.1	10
25	Clinical trial substantiates the predictive value of O-6-methylguanine-DNA methyltransferase promoter methylation in glioblastoma patients treated with temozolomide. <i>Clinical Cancer Research</i> , 2004 , 10, 1871-4	12.9	856
24	Growth promoting signaling by tenascin-C [corrected]. <i>Cancer Research</i> , 2004 , 64, 7377-85	10.1	75
23	INK4a/Arf is required for suppression of EGFR/DeltaEGFR(2-7)-dependent ERK activation in mouse astrocytes and glioma. <i>Oncogene</i> , 2004 , 23, 6854-63	9.2	9
22	Prognostic factors for low-grade gliomas. <i>Seminars in Oncology</i> , 2003 , 30, 23-8	5.5	38
21	Classification of human astrocytic gliomas on the basis of gene expression: a correlated group of genes with angiogenic activity emerges as a strong predictor of subtypes. <i>Cancer Research</i> , 2003 , 63, 6613-25	10.1	168
20	Anoxia induces macrophage inhibitory cytokine-1 (MIC-1) in glioblastoma cells independently of p53 and HIF-1. <i>Oncogene</i> , 2002 , 21, 4212-9	9.2	137
19	p73 is not mutated in meningiomas as determined with a functional yeast assay but p73 expression increases with tumor grade. <i>Brain Pathology</i> , 2001 , 11, 296-305	6	22
18	Quantitative real-time PCR does not show selective targeting of p14(ARF) but concomitant inactivation of both p16(INK4A) and p14(ARF) in 105 human primary gliomas. <i>Oncogene</i> , 2001 , 20, 1103-9	9.2	58
17	p53 gene mutation and ink4a-arf deletion appear to be two mutually exclusive events in human glioblastoma. <i>Oncogene</i> , 2000 , 19, 3816-22	9.2	120

16	Reduced latency but no increased brain tumor penetrance in mice with astrocyte specific expression of a human p53 mutant. <i>Oncogene</i> , 2000 , 19, 5329-37	9.2	12
15	No complementation between TP53 or RB-1 and v-src in astrocytomas of GFAP-v-src transgenic mice. <i>Brain Pathology</i> , 1999 , 9, 627-37	6	15
14	Functional analyses of a unique p53 germline mutant (Y236delta) associated with a familial brain tumor syndrome. <i>International Journal of Cancer</i> , 1999 , 82, 17-22	7.5	5
13	The mouse ERG before and after light damage is independent of p53. <i>Documenta Ophthalmologica</i> , 1998 , 96, 311-20	2.2	5
12	Determination of p53 Mutations, EGFR Overexpression, and Loss of p16 Expression in Pediatric Glioblastomas. <i>Journal of Neuropathology and Experimental Neurology</i> , 1997 , 56, 782-789	3.1	69
11	Hemizygous or homozygous deletion of the chromosomal region containing the p16INK4a gene is associated with amplification of the EGF receptor gene in glioblastomas. <i>International Journal of Cancer</i> , 1997 , 73, 57-63	7.5	44
10	Familial brain tumour syndrome associated with a p53 germline deletion of codon 236. <i>Brain Pathology</i> , 1995 , 5, 15-23	6	32
9	Identical mutations of the p53 tumor suppressor gene in the gliomatous and the sarcomatous components of gliosarcomas suggest a common origin from glial cells. <i>Journal of Neuropathology and Experimental Neurology</i> , 1995 , 54, 651-6	3.1	112
8	Hepatic and pulmonary carcinogenicity of methylene chloride in mice: a search for mechanisms. <i>Toxicology</i> , 1995 , 102, 73-81	4.4	14
7	Allelotype analysis of mouse lung carcinomas reveals frequent allelic losses on chromosome 4 and an association between allelic imbalances on chromosome 6 and K-ras activation. <i>Cancer Research</i> , 1994 , 54, 6257-64	10.1	62
6	Characterization of p53 mutations in methylene chloride-induced lung tumors from B6C3F1 mice. <i>Carcinogenesis</i> , 1993 , 14, 803-10	4.6	78
5	Analysis of activated protooncogenes in B6C3F1 mouse liver tumors induced by ciprofibrate, a potent peroxisome proliferator. <i>Carcinogenesis</i> , 1993 , 14, 145-9	4.6	27
4	Comparison of pulmonary O6-methylguanine DNA adduct levels and Ki-ras activation in lung tumors from resistant and susceptible mouse strains. <i>Molecular Carcinogenesis</i> , 1993 , 8, 177-85	5	59
3	Assignment of a locus for familial melanoma, MLM, to chromosome 9p13-p22. <i>Science</i> , 1992 , 258, 1148-52	5.3	456
2	No measurable increase in thymidine glycol or 8-hydroxydeoxyguanosine in liver DNA of rats treated with nafenopin or choline-devoid low-methionine diet. <i>Mutation Research - Reviews in Genetic Toxicology</i> , 1990 , 238, 325-9		32
1	Detection by ³² P-postlabeling of thymidine glycol in gamma-irradiated DNA. <i>Carcinogenesis</i> , 1989 , 10, 43-7	4.6	41