

Lidiane Zito Grund

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/759396/publications.pdf>

Version: 2024-02-01

15
papers

300
citations

840776

11
h-index

996975

15
g-index

15
all docs

15
docs citations

15
times ranked

370
citing authors

#	ARTICLE	IF	CITATIONS
1	Role of interplay between IL-4 and IFN- \hat{I} 3 in the in regulating M1 macrophage polarization induced by Nattectin. <i>International Immunopharmacology</i> , 2012, 14, 513-522.	3.8	46
2	Nattectin a fish C-type lectin drives Th1 responses in vivo: Licenses macrophages to differentiate into cells exhibiting typical DC function. <i>International Immunopharmacology</i> , 2011, 11, 1546-1556.	3.8	36
3	Analysis of the inflammatory reaction induced by the catfish (<i>Cathorops spixii</i>) venoms. <i>Toxicon</i> , 2007, 49, 909-919.	1.6	30
4	Thalassophryne nattereri fish venom: from the envenoming to the understanding of the immune system. <i>Journal of Venomous Animals and Toxins Including Tropical Diseases</i> , 2014, 20, 35.	1.4	30
5	IL-5 and IL-17A are critical for the chronic IgE response and differentiation of long-lived antibody-secreting cells in inflamed tissues. <i>Cytokine</i> , 2012, 59, 335-351.	3.2	25
6	Experimental immunization with Thalassophryne nattereri fish venom: Striking IL-5 production and impaired of B220+ cells. <i>Toxicon</i> , 2006, 48, 499-508.	1.6	22
7	TLR2, TLR4 and the MyD88 Signaling Are Crucial For the In Vivo Generation and the Longevity of Long-Lived Antibody-Secreting Cells. <i>PLoS ONE</i> , 2013, 8, e71185.	2.5	20
8	The Longevity of Th2 Humoral Response Induced by Proteases Natterins Requires the Participation of Long-Lasting Innate-Like B Cells and Plasma Cells in Spleen. <i>PLoS ONE</i> , 2013, 8, e67135.	2.5	20
9	Multiple functional therapeutic effects of TnP: A small stable synthetic peptide derived from fish venom in a mouse model of multiple sclerosis. <i>PLoS ONE</i> , 2017, 12, e0171796.	2.5	15
10	Natterin an aerolysin-like fish toxin drives IL-1 \hat{I} 2-dependent neutrophilic inflammation mediated by caspase-1 and caspase-11 activated by the inflammasome sensor NLRP6. <i>International Immunopharmacology</i> , 2021, 91, 107287.	3.8	13
11	Stingray venom activates IL-33 producing cardiomyocytes, but not mast cell, to promote acute neutrophil-mediated injury. <i>Scientific Reports</i> , 2017, 7, 7912.	3.3	11
12	Neutrophils releasing IL-17A into NETs are essential to plasma cell differentiation in inflamed tissue dependent on IL-1R. <i>Autoimmunity</i> , 2017, 50, 86-101.	2.6	10
13	The Hierarchical Process of Differentiation of Long-Lived Antibody-Secreting Cells Is Dependent on Integrated Signals Derived from Antigen and IL-17A. <i>PLoS ONE</i> , 2013, 8, e74566.	2.5	9
14	Airborne pollutant ROFA enhances the allergic airway inflammation through direct modulation of dendritic cells in an uptake-dependent mechanism. <i>International Immunopharmacology</i> , 2014, 22, 9-20.	3.8	8
15	Anaphylaxis induced by Thalassophryne nattereri venom in mice is an IgE/IgG1-mediated, IL-4-dependent phenomenon. <i>Scientific Reports</i> , 2020, 10, 584.	3.3	5