

Joan Leung

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/7606674/publications.pdf>

Version: 2024-02-01

13
papers

320
citations

1040056

9
h-index

1125743

13
g-index

13
all docs

13
docs citations

13
times ranked

384
citing authors

#	ARTICLE	IF	CITATIONS
1	Impact of Ankle-foot Orthoses on Gait and Leg Muscle Activity in Adults with Hemiplegia. <i>Physiotherapy</i> , 2003, 89, 39-55.	0.4	125
2	Serial casting versus positioning for the treatment of elbow contractures in adults with traumatic brain injury: a randomized controlled trial. <i>Clinical Rehabilitation</i> , 2008, 22, 406-417.	2.2	43
3	Efficacy of a fitness centre-based exercise programme compared with a home-based exercise programme in traumatic brain injury: A randomized controlled trial. <i>Journal of Rehabilitation Medicine</i> , 2009, 41, 247-255.	1.1	39
4	The impact of simulated ankle plantarflexion contracture on the knee joint during stance phase of gait: A within-subject study. <i>Clinical Biomechanics</i> , 2014, 29, 423-428.	1.2	20
5	The prevalence and characteristics of shoulder pain after traumatic brain injury. <i>Clinical Rehabilitation</i> , 2007, 21, 171-181.	2.2	18
6	Functional electrical stimulation cycling does not improve mobility in people with acquired brain injury and its effects on strength are unclear: a randomised trial. <i>Journal of Physiotherapy</i> , 2016, 62, 203-208.	1.7	16
7	Standing with electrical stimulation and splinting is no better than standing alone for management of ankle plantarflexion contractures in people with traumatic brain injury: a randomised trial. <i>Journal of Physiotherapy</i> , 2014, 60, 201-208.	1.7	15
8	Electrical stimulation and splinting were not clearly more effective than splinting alone for contracture management after acquired brain injury: a randomised trial. <i>Journal of Physiotherapy</i> , 2012, 58, 231-240.	1.7	14
9	Effectiveness of a programme comprising serial casting, botulinum toxin, splinting and motor training for contracture management: a randomized controlled trial. <i>Clinical Rehabilitation</i> , 2019, 33, 1035-1044.	2.2	12
10	An Intensive Programme of Passive Stretch and Motor Training to Manage Severe Knee Contractures after Traumatic Brain Injury: A Case Report. <i>Physiotherapy Canada Physiotherapie Canada</i> , 2013, 65, 223-228.	0.6	6
11	Extra practice outside therapy sessions to maximize training opportunity during inpatient rehabilitation after traumatic brain injury. <i>Brain Injury</i> , 2018, 32, 915-925.	1.2	6
12	Long-Term Resolution of Severe Ankle Contractures Using Botulinum Toxin, Serial Casting, Splinting, and Motor Retraining. <i>Physiotherapy Canada Physiotherapie Canada</i> , 2018, 70, 152-159.	0.6	3
13	Factors Influencing the Overall Satisfaction in Patients With Severe Brain Injury With Physiotherapy Services During Inpatient Rehabilitation. <i>Journal of Head Trauma Rehabilitation</i> , 2018, 33, E56-E63.	1.7	3