

Joan Leung

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/7606674/publications.pdf>

Version: 2024-02-01

13
papers

320
citations

1040056

9
h-index

1125743

13
g-index

13
all docs

13
docs citations

13
times ranked

384
citing authors

| # | ARTICLE | IF | CITATIONS |
|----|---|-----|-----------|
| 1 | Effectiveness of a programme comprising serial casting, botulinum toxin, splinting and motor training for contracture management: a randomized controlled trial. <i>Clinical Rehabilitation</i> , 2019, 33, 1035-1044. | 2.2 | 12 |
| 2 | Extra practice outside therapy sessions to maximize training opportunity during inpatient rehabilitation after traumatic brain injury. <i>Brain Injury</i> , 2018, 32, 915-925. | 1.2 | 6 |
| 3 | Long-Term Resolution of Severe Ankle Contractures Using Botulinum Toxin, Serial Casting, Splinting, and Motor Retraining. <i>Physiotherapy Canada Physiotherapie Canada</i> , 2018, 70, 152-159. | 0.6 | 3 |
| 4 | Factors Influencing the Overall Satisfaction in Patients With Severe Brain Injury With Physiotherapy Services During Inpatient Rehabilitation. <i>Journal of Head Trauma Rehabilitation</i> , 2018, 33, E56-E63. | 1.7 | 3 |
| 5 | Functional electrical stimulation cycling does not improve mobility in people with acquired brain injury and its effects on strength are unclear: a randomised trial. <i>Journal of Physiotherapy</i> , 2016, 62, 203-208. | 1.7 | 16 |
| 6 | Standing with electrical stimulation and splinting is no better than standing alone for management of ankle plantarflexion contractures in people with traumatic brain injury: a randomised trial. <i>Journal of Physiotherapy</i> , 2014, 60, 201-208. | 1.7 | 15 |
| 7 | The impact of simulated ankle plantarflexion contracture on the knee joint during stance phase of gait: A within-subject study. <i>Clinical Biomechanics</i> , 2014, 29, 423-428. | 1.2 | 20 |
| 8 | An Intensive Programme of Passive Stretch and Motor Training to Manage Severe Knee Contractures after Traumatic Brain Injury: A Case Report. <i>Physiotherapy Canada Physiotherapie Canada</i> , 2013, 65, 223-228. | 0.6 | 6 |
| 9 | Electrical stimulation and splinting were not clearly more effective than splinting alone for contracture management after acquired brain injury: a randomised trial. <i>Journal of Physiotherapy</i> , 2012, 58, 231-240. | 1.7 | 14 |
| 10 | Efficacy of a fitness centre-based exercise programme compared with a home-based exercise programme in traumatic brain injury: A randomized controlled trial. <i>Journal of Rehabilitation Medicine</i> , 2009, 41, 247-255. | 1.1 | 39 |
| 11 | Serial casting versus positioning for the treatment of elbow contractures in adults with traumatic brain injury: a randomized controlled trial. <i>Clinical Rehabilitation</i> , 2008, 22, 406-417. | 2.2 | 43 |
| 12 | The prevalence and characteristics of shoulder pain after traumatic brain injury. <i>Clinical Rehabilitation</i> , 2007, 21, 171-181. | 2.2 | 18 |
| 13 | Impact of Ankle-foot Orthoses on Gait and Leg Muscle Activity in Adults with Hemiplegia. <i>Physiotherapy</i> , 2003, 89, 39-55. | 0.4 | 125 |