

# Per Svenningsen

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/761166/publications.pdf>

Version: 2024-02-01

53  
papers

1,375  
citations

304368

22  
h-index

360668

35  
g-index

54  
all docs

54  
docs citations

54  
times ranked

1283  
citing authors

#	ARTICLE	IF	CITATIONS
1	The mineralocorticoid receptor blocker spironolactone lowers plasma interferon- $\beta$ and interleukin-6 in patients with type 2 diabetes and treatment-resistant hypertension. <i>Journal of Hypertension</i> , 2022, 40, 153-162.	0.3	4
2	A new transgene mouse model using an extravesicular EGFP tag enables affinity isolation of cell-specific extracellular vesicles. <i>Scientific Reports</i> , 2022, 12, 496.	1.6	10
3	Mineralocorticoid receptor blockade with spironolactone has no direct effect on plasma IL-17A and injury markers in urine from kidney transplant patients. <i>American Journal of Physiology - Renal Physiology</i> , 2022, 322, F138-F149.	1.3	1
4	Proteinuria is accompanied by intratubular complement activation and apical membrane deposition of C3dg and C5b-9 in kidney transplant recipients. <i>American Journal of Physiology - Renal Physiology</i> , 2022, 322, F150-F163.	1.3	9
5	Detection of DZIP1L mutations by whole-exome sequencing in consanguineous families with polycystic kidney disease. <i>Pediatric Nephrology</i> , 2022, 37, 2657-2665.	0.9	5
6	Sodium retention in the nephrotic syndrome and the non-enzymatic function of prostatic prostasin. <i>Pflugers Archiv European Journal of Physiology</i> , 2022, , 1.	1.3	0
7	Interleukin 17A infusion has no acute or long-term hypertensive action in conscious unrestrained male mice. <i>Pflugers Archiv European Journal of Physiology</i> , 2022, 474, 709-719.	1.3	1
8	An estimate of extracellular vesicle secretion rates of human blood cells. , 2022, 1, .		17
9	Localization and regulation of claudin-14 in experimental models of hypercalcemia. <i>American Journal of Physiology - Renal Physiology</i> , 2021, 320, F74-F86.	1.3	17
10	Non-enzymatic function of prostasin and sodium balance. <i>Acta Physiologica</i> , 2021, 232, e13649.	1.8	2
11	Renal claudin-14 expression is not required for regulating Mg <sup>2+</sup> balance in mice. <i>American Journal of Physiology - Renal Physiology</i> , 2021, 320, F897-F907.	1.3	8
12	Adipocyte-Endothelium Crosstalk in Obesity. <i>Frontiers in Endocrinology</i> , 2021, 12, 681290.	1.5	22
13	Urinary extracellular vesicles: Origin, role as intercellular messengers and biomarkers; efficient sorting and potential treatment options. <i>Acta Physiologica</i> , 2020, 228, e13346.	1.8	62
14	Sodium retention by uPA-plasmin-ENaC in nephrotic syndrome”Authors reply. <i>Acta Physiologica</i> , 2020, 228, e13432.	1.8	4
15	Mechanisms of sodium retention in nephrotic syndrome. <i>Current Opinion in Nephrology and Hypertension</i> , 2020, 29, 207-212.	1.0	25
16	Plasminogen Deficiency and Amiloride Mitigate Angiotensin II-Induced Hypertension in Type 1 Diabetic Mice Suggesting Effects Through the Epithelial Sodium Channel. <i>Journal of the American Heart Association</i> , 2020, 9, e016387.	1.6	12
17	Deficiency of T-type voltage-gated calcium channels results in attenuated weight gain and improved endothelium-dependent dilatation of resistance vessels induced by a high-fat diet in mice. <i>Journal of Physiology and Biochemistry</i> , 2020, 76, 135-145.	1.3	5
18	Proteolytic activation of the epithelial sodium channel: role of pro-protein convertases and prostasin. <i>FASEB Journal</i> , 2020, 34, 1-1.	0.2	1

#	ARTICLE	IF	CITATIONS
19	Urokinase-type plasminogen activator contributes to amiloride-sensitive sodium retention in nephrotic range glomerular proteinuria in mice. <i>Acta Physiologica</i> , 2019, 227, e13362.	1.8	30
20	In human nephrectomy specimens, the kidney level of tubular transport proteins does not correlate with their abundance in urinary extracellular vesicles. <i>American Journal of Physiology - Renal Physiology</i> , 2019, 317, F560-F571.	1.3	22
21	Dietary Na <sup>+</sup> intake in healthy humans changes the urine extracellular vesicle prostasin abundance while the vesicle excretion rate, NCC, and ENaC are not altered. <i>American Journal of Physiology - Renal Physiology</i> , 2019, 317, F1612-F1622.	1.3	12
22	Nephrotic syndrome is associated with increased plasma K <sup>+</sup> concentration, intestinal K <sup>+</sup> losses, and attenuated urinary K <sup>+</sup> excretion: a study in rats and humans. <i>American Journal of Physiology - Renal Physiology</i> , 2019, 317, F1549-F1562.	1.3	4
23	The epithelial Na <sup>+</sup> channel $\alpha$ - and $\beta$ -subunits are cleaved at predicted furin-cleavage sites, glycosylated and membrane associated in human kidney. <i>Pflügers Archiv European Journal of Physiology</i> , 2019, 471, 1383-1396.	1.3	10
24	Bacterial Peptide Display for the Selection of Novel Biotinylating Enzymes. <i>Journal of Visualized Experiments</i> , 2019, , .	0.2	0
25	A bacterial display system for effective selection of protein-biotin ligase BirA variants with novel peptide specificity. <i>Scientific Reports</i> , 2019, 9, 4118.	1.6	6
26	The acute blood pressure-lowering effect of amiloride is independent of endothelial ENaC and eNOS in humans and mice. <i>Acta Physiologica</i> , 2019, 225, e13189.	1.8	10
27	SP709ELEVATED URINARY EXTRACELLULAR VESICLE EXCRETION IN PATIENTS WITH DELAYED GRAFT FUNCTION AFTER DECEASED KIDNEY TRANSPLANTATION. <i>Nephrology Dialysis Transplantation</i> , 2018, 33, i585-i585.	0.4	0
28	Aberrant neuronal differentiation is common in glioma but is associated neither with epileptic seizures nor with better survival. <i>Scientific Reports</i> , 2018, 8, 14965.	1.6	6
29	Hydronephrosis is associated with elevated plasmin in urine in pediatric patients and rats and changes in NCC and $\beta$ -ENaC abundance in rat kidney. <i>American Journal of Physiology - Renal Physiology</i> , 2018, 315, F547-F557.	1.3	5
30	H <sup>+</sup> -ATPase B1 subunit localizes to thick ascending limb and distal convoluted tubule of rodent and human kidney. <i>American Journal of Physiology - Renal Physiology</i> , 2018, 315, F429-F444.	1.3	15
31	Albuminuria in kidney transplant recipients is associated with increased urinary serine proteases and activation of the epithelial sodium channel. <i>American Journal of Physiology - Renal Physiology</i> , 2018, 315, F151-F160.	1.3	26
32	Albuminuria is associated with an increased prostasin in urine while aldosterone has no direct effect on urine and kidney tissue abundance of prostasin. <i>Pflügers Archiv European Journal of Physiology</i> , 2017, 469, 655-667.	1.3	8
33	Urine exosomes from healthy and hypertensive pregnancies display elevated level of $\alpha$ -subunit and cleaved $\alpha$ - and $\beta$ -subunits of the epithelial sodium channel "ENaC". <i>Pflügers Archiv European Journal of Physiology</i> , 2017, 469, 1107-1119.	1.3	28
34	Expression of transcellular and paracellular calcium and magnesium transport proteins in renal and intestinal epithelia during lactation. <i>American Journal of Physiology - Renal Physiology</i> , 2017, 313, F629-F640.	1.3	28
35	Physiology and pathophysiology of the plasminogen system in the kidney. <i>Pflügers Archiv European Journal of Physiology</i> , 2017, 469, 1415-1423.	1.3	34
36	Aberrant glomerular filtration of urokinase-type plasminogen activator in nephrotic syndrome leads to amiloride-sensitive plasminogen activation in urine. <i>American Journal of Physiology - Renal Physiology</i> , 2015, 309, F235-F241.	1.3	35

#	ARTICLE	IF	CITATIONS
37	Diabetic nephropathy is associated with increased urine excretion of proteases plasmin, prostatic and urokinase and activation of amiloride-sensitive current in collecting duct cells. <i>Nephrology Dialysis Transplantation</i> , 2015, 30, 781-789.	0.4	63
38	Urinary serine proteases and activation of ENaC in kidney—implications for physiological renal salt handling and hypertensive disorders with albuminuria. <i>Pflügers Archiv European Journal of Physiology</i> , 2015, 467, 531-542.	1.3	53
39	The Epithelial Sodium Channel $\beta$ -Subunit Is Processed Proteolytically in Human Kidney. <i>Journal of the American Society of Nephrology: JASN</i> , 2015, 26, 95-106.	3.0	55
40	Plasmin in urine from patients with type 2 diabetes and treatment-resistant hypertension activates ENaC in vitro. <i>Journal of Hypertension</i> , 2014, 32, 1672-1677.	0.3	44
41	T-type $Ca^{2+}$ channels facilitate NO-formation, vasodilatation and NO-mediated modulation of blood pressure. <i>Pflügers Archiv European Journal of Physiology</i> , 2014, 466, 2205-2214.	1.3	30
42	TMEM16A is a $Ca^{2+}$ -activated $Cl^{-}$ channel expressed in the renal collecting duct. <i>Acta Physiologica</i> , 2014, 212, 166-174.	1.8	18
43	Regulation of renin secretion by renal juxtaglomerular cells. <i>Pflügers Archiv European Journal of Physiology</i> , 2013, 465, 25-37.	1.3	57
44	Remission of nephrotic syndrome diminishes urinary plasmin content and abolishes activation of ENaC. <i>Pediatric Nephrology</i> , 2013, 28, 1227-1234.	0.9	42
45	ATP Releasing Connexin 30 Hemichannels Mediate Flow-Induced Calcium Signaling in the Collecting Duct. <i>Frontiers in Physiology</i> , 2013, 4, 292.	1.3	43
46	Urinary Plasmin Activates Collecting Duct ENaC Current in Preeclampsia. <i>Hypertension</i> , 2012, 60, 1346-1351.	1.3	59
47	Proteinuric diseases with sodium retention: is plasmin the link?. <i>Clinical and Experimental Pharmacology and Physiology</i> , 2012, 39, 117-124.	0.9	22
48	Physiological regulation of epithelial sodium channel by proteolysis. <i>Current Opinion in Nephrology and Hypertension</i> , 2011, 20, 529-533.	1.0	28
49	Development of a renal collecting duct homing peptide using phage display. <i>FASEB Journal</i> , 2011, 25, 665.19.	0.2	0
50	Prostatic-dependent activation of epithelial $Na^{+}$ channels by low plasmin concentrations. <i>American Journal of Physiology - Regulatory Integrative and Comparative Physiology</i> , 2009, 297, R1733-R1741.	0.9	64
51	Hypotonicity-Induced Renin Exocytosis from Juxtaglomerular Cells Requires Aquaporin-1 and Cyclooxygenase-2. <i>Journal of the American Society of Nephrology: JASN</i> , 2009, 20, 2154-2161.	3.0	22
52	Plasmin in Nephrotic Urine Activates the Epithelial Sodium Channel. <i>Journal of the American Society of Nephrology: JASN</i> , 2009, 20, 299-310.	3.0	236
53	Prostaglandin E2 EP2 and EP4 receptor activation mediates cAMP-dependent hyperpolarization and exocytosis of renin in juxtaglomerular cells. <i>American Journal of Physiology - Renal Physiology</i> , 2005, 289, F989-F997.	1.3	55