
## Wei Chen

## List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/7614103/publications.pdf Version: 2024-02-01



WELCHEN

#	Article	lF	CITATIONS
1	Gaming in China Before the COVID-19 Pandemic and After the Lifting of Lockdowns: a Nationwide Online Retrospective Survey. International Journal of Mental Health and Addiction, 2023, 21, 3293-3305.	4.4	10
2	Impaired robust interhemispheric function integration of depressive brain from RESTâ€metaâ€MDD database in China. Bipolar Disorders, 2022, 24, 400-411.	1.1	8
3	Disrupted presynaptic nectin1-based neuronal adhesion in the entorhinal-hippocampal circuit contributes to early-life stress-induced memory deficits. Translational Psychiatry, 2022, 12, 141.	2.4	2
4	A novel joint index based on peripheral blood CD4+/CD8+ T cell ratio, albumin level, and monocyte count to determine the severity of major depressive disorder. BMC Psychiatry, 2022, 22, 248.	1.1	9
5	Reduced nucleus accumbens functional connectivity in reward network and default mode network in patients with recurrent major depressive disorder. Translational Psychiatry, 2022, 12, .	2.4	20
6	Parietal-hippocampal rTMS improves cognitive function in Alzheimer's disease and increases dynamic functional connectivity of default mode network. Psychiatry Research, 2022, 315, 114721.	1.7	11
7	Common and Specific Alterations of Amygdala Subregions in Major Depressive Disorder With and Without Anxiety: A Combined Structural and Resting-State Functional MRI Study. Frontiers in Human Neuroscience, 2021, 15, 634113.	1.0	4
8	A 36-week multicenter, randomized, double-blind, placebo-controlled, parallel-group, phase 3 clinical trial of sodium oligomannate for mild-to-moderate Alzheimer's dementia. Alzheimer's Research and Therapy, 2021, 13, 62.	3.0	99
9	Disrupted hemispheric connectivity specialization in patients with major depressive disorder: Evidence from the REST-meta-MDD Project. Journal of Affective Disorders, 2021, 284, 217-228.	2.0	23
10	Dysfunctional beliefs and attitudes about sleep are associated with regional homogeneity of left inferior occipital gyrus in primary insomnia patients: A preliminary resting state functional magnetic resonance imaging study. Sleep Medicine, 2021, 81, 188-193.	0.8	11
11	Precision Repetitive Transcranial Magnetic Stimulation Over the Left Parietal Cortex Improves Memory in Alzheimer's Disease: A Randomized, Double-Blind, Sham-Controlled Study. Frontiers in Aging Neuroscience, 2021, 13, 693611.	1.7	15
12	Factors affecting the efficacy of repetitive transcranial magnetic stimulation for patients with Alzheimer's disease. Zhejiang Da Xue Xue Bao Yi Xue Ban = Journal of Zhejiang University Medical Sciences, 2021, 50, 383-389.	0.1	2
13	Abnormal Functional Connectivity of the Amygdala in Mild Cognitive Impairment Patients With Depression Symptoms Revealed by Resting-State fMRI. Frontiers in Psychiatry, 2021, 12, 533428.	1.3	6
14	The coupling of BOLD signal variability and degree centrality underlies cognitive functions and psychiatric diseases. Neurolmage, 2021, 237, 118187.	2.1	11
15	Social media use in China before and during COVID-19: Preliminary results from an online retrospective survey. Journal of Psychiatric Research, 2021, 140, 35-38.	1.5	64
16	Disrupted intrinsic functional brain topology in patients with major depressive disorder. Molecular Psychiatry, 2021, 26, 7363-7371.	4.1	82
17	Brain structural alterations in MDD patients with gastrointestinal symptoms: Evidence from the REST-meta-MDD project. Progress in Neuro-Psychopharmacology and Biological Psychiatry, 2021, 111, 110386.	2.5	18
18	Reduced Plasma Estradiol Levels are Associated with Sleep Apnea in Depressed Peri- and Post-Menopausal Women. Neuropsychiatric Disease and Treatment, 2021, Volume 17, 3483-3488.	1.0	1

Wei Chen

#	Article	IF	CITATIONS
19	Altered resting-state dynamic functional brain networks in major depressive disorder: Findings from the REST-meta-MDD consortium. NeuroImage: Clinical, 2020, 26, 102163.	1.4	76
20	Biotypes of major depressive disorder: Neuroimaging evidence from resting-state default mode network patterns. NeuroImage: Clinical, 2020, 28, 102514.	1.4	51
21	A phase II randomized trial of sodium oligomannate in Alzheimer's dementia. Alzheimer's Research and Therapy, 2020, 12, 110.	3.0	54
22	Disrupted Regional Cerebral Blood Flow in Children With Newly-Diagnosed Type 1 Diabetes Mellitus: An Arterial Spin Labeling Perfusion Magnetic Resonance Imaging Study. Frontiers in Neurology, 2020, 11, 572.	1,1	4
23	Safety and Efficacy of Donepezil 10 mg/day in Patients with Mild to Moderate Alzheimer's Disease. Journal of Alzheimer's Disease, 2020, 74, 199-211.	1.2	8
24	Validation of neuropsychological tests for the China Health and Retirement Longitudinal Study Harmonized Cognitive Assessment Protocol. International Psychogeriatrics, 2019, 31, 1709-1719.	0.6	53
25	Connectome analysis of functional and structural hemispheric brain networks in major depressive disorder. Translational Psychiatry, 2019, 9, 136.	2.4	58
26	Reduced default mode network functional connectivity in patients with recurrent major depressive disorder. Proceedings of the National Academy of Sciences of the United States of America, 2019, 116, 9078-9083.	3.3	441
27	The continuum of care for dementia: needs, resources and practice in China. Journal of Global Health, 2019, 9, 020321.	1.2	33
28	Alterations in the Brain Structure and Functional Connectivity in Aquaporin-4 Antibody-Positive Neuromyelitis Optica Spectrum Disorder. Frontiers in Neuroscience, 2019, 13, 1362.	1.4	6
29	Spatiotemporal, metabolic, and therapeutic characterization of altered functional connectivity in major depressive disorder. Human Brain Mapping, 2018, 39, 1957-1971.	1.9	47
30	Multiple Myeloma, Misdiagnosed As Somatic Symptom Disorder: A Case Report. Frontiers in Psychiatry, 2018, 9, 557.	1.3	1
31	Abnormal changes in functional connectivity between the amygdala and frontal regions are associated with depression in Alzheimer's disease. Neuroradiology, 2018, 60, 1315-1322.	1.1	16
32	The Abnormal Functional Connectivity between the Hypothalamus and the Temporal Gyrus Underlying Depression in Alzheimer's Disease Patients. Frontiers in Aging Neuroscience, 2018, 10, 37.	1.7	28
33	Gender differences in the protective effects of green tea against amnestic mild cognitive impairment in the elderly Han population. Neuropsychiatric Disease and Treatment, 2018, Volume 14, 1795-1801.	1.0	17
34	Decreased functional connectivity between the dorsal anterior cingulate cortex and lingual gyrus in Alzheimer's disease patients with depression. Behavioural Brain Research, 2017, 326, 132-138.	1.2	45
35	Personality traits and perception of Müller-Lyer illusion in male Chinese military soldiers and university students. Translational Neuroscience, 2017, 8, 15-20.	0.7	7
36	Evaluation of the risk factors of depressive disorders comorbid with obstructive sleep apnea. Neuropsychiatric Disease and Treatment, 2017, Volume 13, 155-159.	1.0	13

Wei Chen

#	Article	IF	CITATIONS
37	Abnormal functional connectivity of the posterior cingulate cortex is associated with depressive symptoms in patients with Alzheimer's disease. Neuropsychiatric Disease and Treatment, 2017, Volume 13, 2589-2598.	1.0	13
38	Fractional amplitude of low-frequency fluctuations is disrupted in Alzheimer's disease with depression. Clinical Neurophysiology, 2017, 128, 1344-1349.	0.7	20
39	Relationship of Mean Platelet Volume To MDD: A Retrospective Study. Shanghai Archives of Psychiatry, 2017, 29, 21-29.	0.7	21
40	Delayed help seeking behavior in dementia care: preliminary findings from the Clinical Pathway for Alzheimer's Disease in China (CPAD) study. International Psychogeriatrics, 2016, 28, 211-219.	0.6	34
41	1H-MRS asymmetry changes in the anterior and posterior cingulate gyrus in patients with mild cognitive impairment and mild Alzheimer's disease. Comprehensive Psychiatry, 2016, 69, 179-185.	1.5	12
42	Sympathetic skin response and R–R interval variation in the assessment of clinical remission of bipolar disorder. Psychiatry Research, 2016, 237, 279-281.	1.7	6
43	Neurometabolic characteristics in the anterior cingulate gyrus of Alzheimer's disease patients with depression: a 1H magnetic resonance spectroscopy study. BMC Psychiatry, 2015, 15, 306.	1.1	14
44	Regional Coherence Changes in Alzheimer's Disease Patients with Depressive Symptoms: A Resting-State Functional MRI Study. Journal of Alzheimer's Disease, 2015, 48, 603-611.	1.2	16
45	Caspase-8 Mediates Amyloid-β-induced Apoptosis in Differentiated PC12 Cells. Journal of Molecular Neuroscience, 2015, 56, 491-499.	1.1	16
46	Sub-hubs of baseline functional brain networks are related to early improvement following two-week pharmacological therapy for major depressive disorder. Human Brain Mapping, 2015, 36, 2915-2927.	1.9	72
47	Relationship between Personality Disorder Functioning Styles and the Emotional States in Bipolar I and II Disorders. PLoS ONE, 2015, 10, e0117353.	1.1	9
48	Comparison of the effectiveness of duloxetine in depressed patients with and without a family history of affective disorders in first-degree relatives. Shanghai Archives of Psychiatry, 2015, 27, 237-45.	0.7	2
49	Proteomic analysis of ischemic rat brain after human mesenchymal stem cell transplantation. Tissue Engineering and Regenerative Medicine, 2014, 11, 333-339.	1.6	3
50	Methodology of China's national study on the evaluation, early recognition, and treatment of psychological problems in the elderly: the China Longitudinal Aging Study (CLAS). Shanghai Archives of Psychiatry, 2013, 25, 91-8.	0.7	40
51	Triptolide Downâ€regulates COXâ€2 Expression and PGE2 Release by Suppressing the Activity of NFâ€î®B and MAP kinases in Lipopolysaccharideâ€treated PC12 Cells. Phytotherapy Research, 2012, 26, 337-343.	2.8	40
52	Roles of the prostaglandin E2 receptors EP subtypes in Alzheimer's disease. Neuroscience Bulletin, 2010, 26, 77-84.	1.5	12
53	Adjectival Descriptors for Antisocial Personality Trait in Chinese University Students. Journal of Personality Disorders, 2009, 23, 661-668.	0.8	6