

Leif Thuesen

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/7620303/leif-thuesen-publications-by-year.pdf>

Version: 2024-04-28

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

145
papers

11,218
citations

51
h-index

105
g-index

159
ext. papers

12,616
ext. citations

8.1
avg, IF

5.18
L-index

#	Paper	IF	Citations
145	Age-Stratified Outcome in Treatment of Left Main Coronary Artery Stenosis: A NOBLE Trial Substudy. <i>Cardiology</i> , 2021 , 146, 409-418	1.6	1
144	Randomized Clinical Comparison of the Dual-Therapy CD34 Antibody-Covered Sirolimus-Eluting Combo Stent With the Sirolimus-Eluting Orsiro Stent in Patients Treated With Percutaneous Coronary Intervention: The SORT OUT X Trial. <i>Circulation</i> , 2021 , 143, 2155-2165	16.7	6
143	16-year follow-up of the Danish Acute Myocardial Infarction 2 (DANAMI-2) trial: primary percutaneous coronary intervention vs. fibrinolysis in ST-segment elevation myocardial infarction. <i>European Heart Journal</i> , 2020 , 41, 847-854	9.5	17
142	Randomized Comparison of the Polymer-Free Biolimus-Coated BioFreedom Stent With the Ultrathin Strut Biodegradable Polymer Sirolimus-Eluting Orsiro Stent in an All-Comers Population Treated With Percutaneous Coronary Intervention: The SORT OUT IX Trial. <i>Circulation</i> , 2020 , 141, 2052-2063	16.7	23
141	Randomised comparison of provisional side branch stenting versus a two-stent strategy for treatment of true coronary bifurcation lesions involving a large side branch: the Nordic-Baltic Bifurcation Study IV. <i>Open Heart</i> , 2020 , 7, e000947	3	5
140	Percutaneous coronary angioplasty versus coronary artery bypass grafting in the treatment of unprotected left main stenosis: updated 5-year outcomes from the randomised, non-inferiority NOBLE trial. <i>Lancet, The</i> , 2020 , 395, 191-199	40	136
139	Ten-Year Outcomes of Sirolimus-Eluting Versus Zotarolimus-Eluting Coronary Stents in Patients With Versus Without Diabetes Mellitus (SORT OUT III). <i>American Journal of Cardiology</i> , 2020 , 125, 349-353	3	5
138	Lower ST-elevation myocardial infarction incidence during COVID-19 epidemic in Northern Europe. <i>Scandinavian Cardiovascular Journal</i> , 2020 , 54, 358-360	2	6
137	Derivation and Validation of a Chronic Total Coronary Occlusion Intervention Procedural Success Score From the 20,000-Patient EuroCTO Registry: The EuroCTO (CASTLE) Score. <i>JACC: Cardiovascular Interventions</i> , 2019 , 12, 335-342	5	59
136	Ten-year clinical outcome of patients treated with a drug-eluting stent in the proximal left anterior descending artery segment compared with patients stented in other non-left main coronary segments. <i>EuroIntervention</i> , 2018 , 14, 764-771	3.1	5
135	Comparison of Durable-Polymer Zotarolimus-Eluting and Biodegradable-Polymer Biolimus-Eluting Coronary Stents in Patients With Coronary Artery Disease: 3-Year Clinical Outcomes in the Randomized SORT OUT VI Trial. <i>JACC: Cardiovascular Interventions</i> , 2017 , 10, 255-264	5	33
134	10-Year Clinical Outcome After Randomization to Treatment by Sirolimus- or Paclitaxel-Eluting Coronary Stents. <i>Journal of the American College of Cardiology</i> , 2017 , 69, 616-624	15.1	38
133	Chronic total coronary occlusion: treatment results. <i>Scandinavian Cardiovascular Journal</i> , 2017 , 51, 197-201		6
132	Stents versus bypass surgery for left main stem stenosis - AuthorsTreply. <i>Lancet, The</i> , 2017 , 389, 1609	40	3
131	Clinical outcomes with percutaneous coronary revascularization vs coronary artery bypass grafting surgery in patients with unprotected left main coronary artery disease: A meta-analysis of 6 randomized trials and 4,686 patients. <i>American Heart Journal</i> , 2017 , 190, 54-63	4.9	62
130	Intravascular ultrasound assessment of minimum lumen area and intimal hyperplasia in in-stent restenosis after drug-eluting or bare-metal stent implantation. The Nordic Intravascular Ultrasound Study (NIVUS). <i>Cardiovascular Revascularization Medicine</i> , 2017 , 18, 577-582	1.6	11
129	All-cause mortality and major cardiovascular outcomes comparing percutaneous coronary angioplasty versus coronary artery bypass grafting in the treatment of unprotected left main stenosis: a meta-analysis of short-term and long-term randomised trials. <i>Open Heart</i> , 2017 , 4, e000638	3	8

128	Percutaneous coronary angioplasty versus coronary artery bypass grafting in treatment of unprotected left main stenosis (NOBLE): a prospective, randomised, open-label, non-inferiority trial. <i>Lancet, The</i> , 2016 , 388, 2743-2752	40	446
127	Impact of the Everolimus-eluting Bioresorbable Scaffold in Coronary Atherosclerosis. <i>Revista Espanola De Cardiologia (English Ed)</i> , 2016 , 69, 109-16	0.7	3
126	Efecto del armazn bioabsorbible liberador de everolimus en la aterosclerosis coronaria. <i>Revista Espanola De Cardiologia</i> , 2016 , 69, 109-116	1.5	6
125	Worsening Clinical Outcome with Increasing Number of So-Called Off-Label or Unapproved Indications for Use of Drug Eluting Stents. <i>World Journal of Cardiovascular Diseases</i> , 2016 , 06, 224-234	0	
124	Coronary bifurcation lesions treated with simple or complex stenting: 5-year survival from patient-level pooled analysis of the Nordic Bifurcation Study and the British Bifurcation Coronary Study. <i>European Heart Journal</i> , 2016 , 37, 1923-8	9.5	67
123	Zotarolimus-eluting durable-polymer-coated stent versus a biolimus-eluting biodegradable-polymer-coated stent in unselected patients undergoing percutaneous coronary intervention (SORT OUT VI): a randomised non-inferiority trial. <i>Lancet, The</i> , 2015 , 385, 1527-35	40	92
122	Influence of multivessel disease with or without additional revascularization on mortality in patients with ST-segment elevation myocardial infarction. <i>American Heart Journal</i> , 2015 , 170, 70-8	4.9	16
121	Coronary bifurcation lesions: Present status and future perspectives. <i>International Journal of Cardiology</i> , 2015 , 187, 48-57	3.2	18
120	One-year clinical and angiographic results of hybrid coronary revascularization. <i>Journal of Thoracic and Cardiovascular Surgery</i> , 2015 , 150, 1181-6	1.5	13
119	The impact of distal embolization and distal protection on long-term outcome in patients with ST elevation myocardial infarction randomized to primary percutaneous coronary intervention--results from a randomized study. <i>European Heart Journal: Acute Cardiovascular Care</i> , 2015 , 4, 180-8	4.3	14
118	Retrograde Recanalization of Chronic Total Occlusions in Europe: Procedural, In-Hospital, and Long-Term Outcomes From the Multicenter ERCTO Registry. <i>Journal of the American College of Cardiology</i> , 2015 , 65, 2388-400	15.1	160
117	Feasibility and early safety of hybrid coronary revascularisation combining off-pump coronary surgery through J-hemisternotomy with percutaneous coronary intervention. <i>EuroIntervention</i> , 2015 , 10, e1-6	3.1	8
116	Clopidogrel discontinuation within the first year after coronary drug-eluting stent implantation: an observational study. <i>BMC Cardiovascular Disorders</i> , 2014 , 14, 100	2.3	22
115	Stent thrombosis is the primary cause of ST-segment elevation myocardial infarction following coronary stent implantation: a five year follow-up of the SORT OUT II study. <i>PLoS ONE</i> , 2014 , 9, e113399 ³⁻⁷	3.7	8
114	Similar five-year outcome with paclitaxel- and sirolimus-eluting coronary stents. <i>Scandinavian Cardiovascular Journal</i> , 2014 , 48, 148-55	2	8
113	Serial optical frequency domain imaging in STEMI patients: the follow-up report of TROFI study. <i>European Heart Journal Cardiovascular Imaging</i> , 2014 , 15, 987-95	4.1	28
112	Differential clinical outcomes after 1 year versus 5 years in a randomised comparison of zotarolimus-eluting and sirolimus-eluting coronary stents (the SORT OUT III study): a multicentre, open-label, randomised superiority trial. <i>Lancet, The</i> , 2014 , 383, 2047-2056	40	80
111	Long-term outcome after drug-eluting versus bare-metal stent implantation in patients with ST-segment elevation myocardial infarction: 5 years follow-up from the randomized DEDICATION trial (Drug Elution and Distal Protection in Acute Myocardial Infarction). <i>JACC: Cardiovascular Interventions</i> , 2013 , 6, 548-59	5	32

110	Five-year clinical and functional multislice computed tomography angiographic results after coronary implantation of the fully resorbable polymeric everolimus-eluting scaffold in patients with de novo coronary artery disease: the ABSORB cohort A trial. <i>JACC: Cardiovascular Interventions</i> , 2013 , 6, 999-1009	5	164
109	In vivo three dimensional optical coherence tomography. A novel imaging modality to visualize the edge vascular response. <i>International Journal of Cardiology</i> , 2013 , 164, e35-7	3.2	5
108	Biolimus-eluting biodegradable polymer-coated stent versus durable polymer-coated sirolimus-eluting stent in unselected patients receiving percutaneous coronary intervention (SORT OUT V): a randomised non-inferiority trial. <i>Lancet, The</i> , 2013 , 381, 661-9	4.0	152
107	Intravascular ultrasound assessed incomplete stent apposition and stent fracture in stent thrombosis after bare metal versus drug-eluting stent treatment the Nordic Intravascular Ultrasound Study (NIVUS). <i>International Journal of Cardiology</i> , 2013 , 168, 1010-6	3.2	24
106	Diagnosis and outcome in a prehospital cohort of patients with bundle branch block and suspected acute myocardial infarction. <i>European Heart Journal: Acute Cardiovascular Care</i> , 2013 , 2, 176-81	4.3	7
105	Clinical and intravascular imaging outcomes at 1 and 2 years after implantation of absorb everolimus eluting bioresorbable vascular scaffolds in small vessels. Late lumen enlargement: does bioresorption matter with small vessel size? Insight from the ABSORB cohort B trial. <i>Heart</i> , 2013 , 89, 106-107	5.1	60
104	Moderate overweight is beneficial and severe obesity detrimental for patients with documented atherosclerotic heart disease. <i>Heart</i> , 2013 , 99, 655-60	5.1	51
103	Outcomes after primary percutaneous coronary intervention in octogenarians and nonagenarians with ST-segment elevation myocardial infarction: from the Western Denmark heart registry. <i>Catheterization and Cardiovascular Interventions</i> , 2013 , 81, 912-9	2.7	48
102	Incidence of definite stent thrombosis or in-stent restenosis after drug-eluting stent implantation for treatment of coronary in-stent restenosis: from Western Denmark Heart Registry. <i>Catheterization and Cardiovascular Interventions</i> , 2013 , 81, 260-5	2.7	11
101	Event detection using population-based health care databases in randomized clinical trials: a novel research tool in interventional cardiology. <i>Clinical Epidemiology</i> , 2013 , 5, 357-61	5.9	21
100	Clinical outcomes after treatment of multiple lesions with zotarolimus-eluting versus sirolimus-eluting coronary stents (a SORT OUT III substudy). <i>BMC Cardiovascular Disorders</i> , 2012 , 12, 18	2.3	0
99	Analysis of 1 year virtual histology changes in coronary plaque located behind the struts of the everolimus eluting bioresorbable vascular scaffold. <i>International Journal of Cardiovascular Imaging</i> , 2012 , 28, 1307-14	2.5	12
98	Comparison of zotarolimus-eluting and sirolimus-eluting coronary stents: a study from the Western Denmark Heart Registry. <i>BMC Cardiovascular Disorders</i> , 2012 , 12, 84	2.3	2
97	2-year patient-related versus stent-related outcomes: the SORT OUT IV (Scandinavian Organization for Randomized Trials With Clinical Outcome IV) Trial. <i>Journal of the American College of Cardiology</i> , 2012 , 60, 1140-7	15.1	41
96	Vascular response of the segments adjacent to the proximal and distal edges of the ABSORB everolimus-eluting bioresorbable vascular scaffold: 6-month and 1-year follow-up assessment: a virtual histology intravascular ultrasound study from the first-in-man ABSORB cohort B trial. <i>JACC: Cardiovascular Interventions</i> , 2012 , 5, 656-65	5	29
95	3-Year clinical outcomes in the randomized SORT OUT III superiority trial comparing zotarolimus- and sirolimus-eluting coronary stents. <i>JACC: Cardiovascular Interventions</i> , 2012 , 5, 812-8	5	42
94	Influence of diabetes mellitus on clinical outcomes following primary percutaneous coronary intervention in patients with ST-segment elevation myocardial infarction. <i>American Journal of Cardiology</i> , 2012 , 109, 629-35	3	43
93	Dimensions of socioeconomic status and clinical outcome after primary percutaneous coronary intervention. <i>Circulation: Cardiovascular Interventions</i> , 2012 , 5, 641-8	6	30

92	Randomized comparison of everolimus-eluting and sirolimus-eluting stents in patients treated with percutaneous coronary intervention: the Scandinavian Organization for Randomized Trials with Clinical Outcome IV (SORT OUT IV). <i>Circulation</i> , 2012 , 125, 1246-55	16.7	131
91	Influence of pre-infarction angina, collateral flow, and pre-procedural TIMI flow on myocardial salvage index by cardiac magnetic resonance in patients with ST-segment elevation myocardial infarction. <i>European Heart Journal Cardiovascular Imaging</i> , 2012 , 13, 433-43	4.1	41
90	Side branch fractional flow reserve measurements after main vessel stenting: a Nordic-Baltic Bifurcation Study III substudy. <i>EuroIntervention</i> , 2012 , 7, 1155-61	3.1	41
89	Four-year clinical follow-up of the ABSORB everolimus-eluting bioresorbable vascular scaffold in patients with de novo coronary artery disease: the ABSORB trial. <i>EuroIntervention</i> , 2012 , 7, 1060-1	3.1	94
88	Culprit only or multivessel percutaneous coronary interventions in patients with ST-segment elevation myocardial infarction and multivessel disease. <i>EuroIntervention</i> , 2012 , 8, 456-64	3.1	32
87	The risk and prognostic impact of definite stent thrombosis or in-stent restenosis after coronary stent implantation. <i>EuroIntervention</i> , 2012 , 8, 591-8	3.1	17
86	Commentary: Deliver the drug and disappear: is the bioabsorbable magnesium stent growing up or still shrinking?. <i>Journal of Endovascular Therapy</i> , 2011 , 18, 416-7	2.5	
85	Health care system delay and heart failure in patients with ST-segment elevation myocardial infarction treated with primary percutaneous coronary intervention: follow-up of population-based medical registry data. <i>Annals of Internal Medicine</i> , 2011 , 155, 361-7	8	69
84	System delay and timing of intervention in acute myocardial infarction (from the Danish Acute Myocardial Infarction-2 [DANAMI-2] trial). <i>American Journal of Cardiology</i> , 2011 , 108, 776-81	3	41
83	Transatrial stent-valve implantation in a stenotic tricuspid valve bioprosthesis. <i>Annals of Thoracic Surgery</i> , 2011 , 91, e74-6	2.7	7
82	Temporal changes of coronary artery plaque located behind the struts of the everolimus eluting bioresorbable vascular scaffold. <i>International Journal of Cardiovascular Imaging</i> , 2011 , 27, 859-66	2.5	17
81	Randomized comparison of final kissing balloon dilatation versus no final kissing balloon dilatation in patients with coronary bifurcation lesions treated with main vessel stenting: the Nordic-Baltic Bifurcation Study III. <i>Circulation</i> , 2011 , 123, 79-86	16.7	219
80	Primary percutaneous coronary intervention as a national reperfusion strategy in patients with ST-segment elevation myocardial infarction. <i>Circulation: Cardiovascular Interventions</i> , 2011 , 4, 570-6	6	31
79	Influence of distance from home to invasive centre on invasive treatment after acute coronary syndrome: a nationwide study of 24 910 patients. <i>Heart</i> , 2011 , 97, 27-32	5.1	13
78	Existing data sources for clinical epidemiology: The Western Denmark Heart Registry. <i>Clinical Epidemiology</i> , 2010 , 2, 137-44	5.9	131
77	System delay and mortality among patients with STEMI treated with primary percutaneous coronary intervention. <i>JAMA - Journal of the American Medical Association</i> , 2010 , 304, 763-71	27.4	431
76	Increased rate of stent thrombosis and target lesion revascularization after filter protection in primary percutaneous coronary intervention for ST-segment elevation myocardial infarction: 15-month follow-up of the DEDICATION (Drug Elution and Distal Protection in ST Elevation Myocardial Infarction) trial. <i>Journal of the American College of Cardiology</i> , 2010 , 55, 867-71	15.1	30
75	Sirolimus-eluting versus bare-metal stent implantation in patients with ostial lesions. <i>International Journal of Cardiology</i> , 2010 , 145, 162-3	3.2	1

74	Thrombus Aspiration in ST-Elevation myocardial infarction in Scandinavia (TASTE trial). A multicenter, prospective, randomized, controlled clinical registry trial based on the Swedish angiography and angioplasty registry (SCAAR) platform. Study design and rationale. <i>American Heart Journal</i> , 2010 , 160, 1042-8	4.9	111
73	Remote ischaemic conditioning before hospital admission, as a complement to angioplasty, and effect on myocardial salvage in patients with acute myocardial infarction: a randomised trial. <i>Lancet, The</i> , 2010 , 375, 727-34	4.0	773
72	Efficacy and safety of zotarolimus-eluting and sirolimus-eluting coronary stents in routine clinical care (SORT OUT III): a randomised controlled superiority trial. <i>Lancet, The</i> , 2010 , 375, 1090-9	4.0	174
71	Zotarolimus-eluting versus sirolimus-eluting coronary stent implantation. <i>Interventional Cardiology</i> , 2010 , 2, 807-812	3	1
70	Long-term outcomes after percutaneous coronary intervention in patients with and without diabetes mellitus in Western Denmark. <i>American Journal of Cardiology</i> , 2010 , 105, 1513-9	3	35
69	IVUS radiofrequency analysis in the evaluation of the polymeric struts of the bioabsorbable everolimus-eluting device during the bioabsorption process. <i>Catheterization and Cardiovascular Interventions</i> , 2010 , 75, 914-8	2.7	16
68	Five-year long-term clinical follow-up of the XIENCE V everolimus eluting coronary stent system in the treatment of patients with de novo coronary artery lesions: the SPIRIT FIRST trial. <i>Catheterization and Cardiovascular Interventions</i> , 2010 , 75, 997-1003	2.7	42
67	Three-year results of clinical follow-up after a bioresorbable everolimus-eluting scaffold in patients with de novo coronary artery disease: the ABSORB trial. <i>EuroIntervention</i> , 2010 , 6, 447-53	3.1	100
66	Paclitaxel and sirolimus eluting stents versus bare metal stents: long-term risk of stent thrombosis and other outcomes. From the Western Denmark Heart Registry. <i>EuroIntervention</i> , 2010 , 5, 898-905	3.1	39
65	Infarct size and myocardial salvage after primary angioplasty in patients presenting with symptoms for . <i>European Heart Journal</i> , 2009 , 30, 1322-30	9.5	65
64	Randomized comparison of coronary bifurcation stenting with the crush versus the culotte technique using sirolimus eluting stents: the Nordic stent technique study. <i>Circulation: Cardiovascular Interventions</i> , 2009 , 2, 27-34	6	133
63	Clinical reinfarction according to infarct location and reperfusion modality in patients with ST elevation myocardial infarction. A DANAMI-2 long-term follow-up substudy. <i>Cardiology</i> , 2009 , 113, 72-80 ^{1.6}		3
62	ST changes before and during primary percutaneous coronary intervention predict final infarct size in patients with ST elevation myocardial infarction. <i>Journal of Electrocardiology</i> , 2009 , 42, 64-72	1.4	22
61	Comparison of the sirolimus-eluting versus paclitaxel-eluting coronary stent in patients with diabetes mellitus: the diabetes and drug-eluting stent (DiabeDES) randomized angiography trial. <i>American Journal of Cardiology</i> , 2009 , 103, 345-9	3	48
60	Timing, causes, and predictors of death after three years follow-up in the Danish Multicenter Randomized Study of Fibrinolysis versus Primary Angioplasty in Acute Myocardial Infarction (DANAMI-2) trial. <i>American Journal of Cardiology</i> , 2009 , 104, 210-5	3	15
59	Prevalence and significance of accelerated idioventricular rhythm in patients with ST-elevation myocardial infarction treated with primary percutaneous coronary intervention. <i>American Journal of Cardiology</i> , 2009 , 104, 1641-6	3	39
58	Transapical transcatheter treatment of a stenosed aortic valve bioprosthesis using the Edwards SAPIEN Transcatheter Heart Valve. <i>Annals of Thoracic Surgery</i> , 2009 , 87, 1943-6	2.7	26
57	2-year clinical outcomes after implantation of sirolimus-eluting, paclitaxel-eluting, and bare-metal coronary stents: results from the WDHR (Western Denmark Heart Registry). <i>Journal of the American College of Cardiology</i> , 2009 , 53, 658-64	15.1	268

56	A bioabsorbable everolimus-eluting coronary stent system (ABSORB): 2-year outcomes and results from multiple imaging methods. <i>Lancet, The</i> , 2009 , 373, 897-910	40	642
55	Assessment of the absorption process following bioabsorbable everolimus-eluting stent implantation: temporal changes in strain values and tissue composition using intravascular ultrasound radiofrequency data analysis. A substudy of the ABSORB clinical trial. <i>EuroIntervention</i> , 2009 , 4, 117-22	3.1	52
54	Randomized comparison of distal protection versus conventional treatment in primary percutaneous coronary intervention: the drug elution and distal protection in ST-elevation myocardial infarction (DEDICATION) trial. <i>Journal of the American College of Cardiology</i> , 2008 , 51, 899-905	15.1	117
53	Long-term outcome in patients treated with sirolimus-eluting stents in complex coronary artery lesions: 3-year results of the SCANDSTENT (Stenting Coronary Arteries in Non-Stress/Benestent Disease) trial. <i>Journal of the American College of Cardiology</i> , 2008 , 51, 2011-6	15.1	44
52	A bioabsorbable everolimus-eluting coronary stent system for patients with single de-novo coronary artery lesions (ABSORB): a prospective open-label trial. <i>Lancet, The</i> , 2008 , 371, 899-907	40	552
51	Diastolic dysfunction after an acute myocardial infarction in patients with antecedent hypertension. <i>Journal of the American Society of Echocardiography</i> , 2008 , 21, 171-7	5.8	3
50	Neointimal hyperplasia after sirolimus-eluting and paclitaxel-eluting stent implantation in diabetic patients: the Randomized Diabetes and Drug-Eluting Stent (DiabeDES) Intravascular Ultrasound Trial. <i>European Heart Journal</i> , 2008 , 29, 2733-41	9.5	32
49	Comparison of paclitaxel- and sirolimus-eluting stents in everyday clinical practice: the SORT OUT II randomized trial. <i>JAMA - Journal of the American Medical Association</i> , 2008 , 299, 409-16	27.4	116
48	Clinical outcome after primary percutaneous coronary intervention with drug-eluting and bare metal stents in patients with ST-segment elevation myocardial infarction. <i>Circulation: Cardiovascular Interventions</i> , 2008 , 1, 176-84	6	26
47	Drug-eluting versus bare metal stents in patients with st-segment-elevation myocardial infarction: eight-month follow-up in the Drug Elution and Distal Protection in Acute Myocardial Infarction (DEDICATION) trial. <i>Circulation</i> , 2008 , 118, 1155-62	16.7	58
46	The Danish multicentre randomized study of fibrinolytic therapy vs. primary angioplasty in acute myocardial infarction (the DANAMI-2 trial): outcome after 3 years follow-up. <i>European Heart Journal</i> , 2008 , 29, 1259-66	9.5	58
45	Safety in simple versus complex stenting of coronary artery bifurcation lesions. The nordic bifurcation study 14-month follow-up results. <i>EuroIntervention</i> , 2008 , 4, 229-33	3.1	53
44	Abstract from the Bern-Rotterdam registry. <i>EuroIntervention</i> , 2008 , 4 Suppl C, C52-4	3.1	
43	Stent thrombosis, myocardial infarction, and death after drug-eluting and bare-metal stent coronary interventions. <i>Journal of the American College of Cardiology</i> , 2007 , 50, 463-70	15.1	195
42	Comparison of in vivo acute stent recoil between the bioabsorbable everolimus-eluting coronary stent and the everolimus-eluting cobalt chromium coronary stent: insights from the ABSORB and SPIRIT trials. <i>Catheterization and Cardiovascular Interventions</i> , 2007 , 70, 515-23	2.7	124
41	Influence of a pressure gradient distal to implanted bare-metal stent on in-stent restenosis after percutaneous coronary intervention. <i>Circulation</i> , 2007 , 116, 2802-8	16.7	17
40	Two-year results of a durable polymer everolimus-eluting stent in de novo coronary artery stenosis (The SPIRIT FIRST Trial). <i>EuroIntervention</i> , 2007 , 3, 206-12	3.1	24
39	Safety and efficacy of multiple, overlapping polymer-based paclitaxel-eluting stents. <i>EuroIntervention</i> , 2007 , 3, 213-21	3.1	11

38	Effectiveness of "direct" stenting without balloon predilatation (from the Multilink Tetra Randomised European Direct Stent Study [TRENDS]). <i>American Journal of Cardiology</i> , 2006 , 97, 316-21	3	14
37	Potential significance of spontaneous and interventional ST-changes in patients transferred for primary percutaneous coronary intervention: observations from the ST-MONitoring in Acute Myocardial Infarction study (The MONAMI study). <i>European Heart Journal</i> , 2006 , 27, 267-75	9.5	57
36	Randomized study on simple versus complex stenting of coronary artery bifurcation lesions: the Nordic bifurcation study. <i>Circulation</i> , 2006 , 114, 1955-61	16.7	574
35	Routine thrombectomy in percutaneous coronary intervention for acute ST-segment-elevation myocardial infarction: a randomized, controlled trial. <i>Circulation</i> , 2006 , 114, 40-7	16.7	216
34	Distal embolic protection during percutaneous coronary intervention in patients with acute coronary syndromes: the RUBY study. <i>Acute Cardiac Care</i> , 2006 , 8, 148-54		8
33	The Stenting Coronary Arteries in Non-stress/benestent Disease (SCANDSTENT) trial. <i>Journal of the American College of Cardiology</i> , 2006 , 47, 449-55	15.1	95
32	Comparison of sirolimus-eluting and bare metal stents in coronary bifurcation lesions: subgroup analysis of the Stenting Coronary Arteries in Non-Stress/Benestent Disease Trial (SCANDSTENT). <i>American Heart Journal</i> , 2006 , 152, 1140-5	4.9	67
31	Target vessel revascularization following percutaneous coronary intervention. A 10-year report from the Danish Percutaneous Transluminal Coronary Angioplasty Registry. <i>Scandinavian Cardiovascular Journal</i> , 2005 , 39, 30-5	2	1
30	Direct intramyocardial plasmid vascular endothelial growth factor-A165 gene therapy in patients with stable severe angina pectoris A randomized double-blind placebo-controlled study: the Euroinject One trial. <i>Journal of the American College of Cardiology</i> , 2005 , 45, 982-8	15.1	389
29	Clinical efficacy of polymer-based paclitaxel-eluting stents in the treatment of complex, long coronary artery lesions from a multicenter, randomized trial: support for the use of drug-eluting stents in contemporary clinical practice. <i>Circulation</i> , 2005 , 112, 3306-13	16.7	266
28	In-laboratory femoral sheath removal after heparin reversal by protamine after percutaneous coronary intervention. <i>EuroIntervention</i> , 2005 , 1, 66-9	3.1	6
27	Randomized comparison of deliverability and in-hospital complications in implantation of BxSonic(R), Express(R), and Flexmaster(R) coronary stents. <i>EuroIntervention</i> , 2005 , 1, 273-6	3.1	
26	Arterial concentration of 99mTc-sestamibi at rest, during peak exercise and after dipyridamole infusion. <i>Clinical Physiology and Functional Imaging</i> , 2004 , 24, 394-7	2.4	2
25	The NUGGET study: NIR ultra gold-gilded equivalency trial. <i>Catheterization and Cardiovascular Interventions</i> , 2004 , 62, 18-25	2.7	20
24	Randomized comparison of the coil-design Crossflex and the tubular NIR stent. <i>Catheterization and Cardiovascular Interventions</i> , 2003 , 59, 8-12	2.7	6
23	Catheter-based ³² P beta-radiation after stent implantation in porcine coronary arteries: role of source-centering and geographical miss. <i>Catheterization and Cardiovascular Interventions</i> , 2003 , 60, 247-57	2.7	4
22	Electromechanical mapping versus positron emission tomography and single photon emission computed tomography for the detection of myocardial viability in patients with ischemic cardiomyopathy. <i>Journal of the American College of Cardiology</i> , 2003 , 41, 843-8	15.1	35
21	A comparison of coronary angioplasty with fibrinolytic therapy in acute myocardial infarction. <i>New England Journal of Medicine</i> , 2003 , 349, 733-42	59.2	1005

20	Long genuine coronary artery lesions treated with stiff tubular or flexible coiled stents. A randomized angiographic follow-up study. <i>Scandinavian Cardiovascular Journal</i> , 2002 , 36, 91-4	2	5
19	Establishing primary angioplasty as the preferred treatment for acute myocardial infarction. <i>Scandinavian Cardiovascular Journal</i> , 2002 , 36, 215-20	2	1
18	Electromechanical mapping for detection of myocardial viability in patients with ischemic cardiomyopathy. <i>Circulation</i> , 2001 , 103, 1631-7	16.7	68
17	Myocardial laser revascularization--the end of a new therapy?. <i>Scandinavian Cardiovascular Journal</i> , 2001 , 35, 6-7	2	0
16	Growth Hormone Replacement and Cardiac Function in the Growth Hormone Deficient Adults. <i>Growth Hormone</i> , 2001 , 29-32		
15	Functional significance of recruitable collaterals during temporary coronary occlusion evaluated by 99mTc-sestamibi single-photon emission computerized tomography. <i>Journal of the American College of Cardiology</i> , 2000 , 35, 624-32	15.1	26
14	Atrioventricular conduction during long-term follow-up of patients with sick sinus syndrome. <i>Circulation</i> , 1998 , 98, 1315-21	16.7	90
13	Heart failure and echocardiographic changes during long-term follow-up of patients with sick sinus syndrome randomized to single-chamber atrial or ventricular pacing. <i>Circulation</i> , 1998 , 97, 987-95	16.7	182
12	Growth hormone versus placebo treatment for one year in growth hormone deficient adults: increase in exercise capacity and normalization of body composition. <i>Clinical Endocrinology</i> , 1996 , 45, 681-8	3.4	80
11	Three years of growth hormone treatment in growth hormone-deficient adults: near normalization of body composition and physical performance. <i>European Journal of Endocrinology</i> , 1994 , 130, 224-8	6.5	157
10	Short and long-term cardiovascular effects of growth hormone therapy in growth hormone deficient adults. <i>Clinical Endocrinology</i> , 1994 , 41, 615-20	3.4	100
9	The Abnormal Albuminuria Syndrome in Diabetes. <i>Frontiers in Diabetes</i> , 1993 , 12, 86-121	0.6	5
8	The Heart in Diabetic Nephropathy 1992 , 205-217		
7	The cardiovascular effects of octreotide treatment in acromegaly: an echocardiographic study. <i>Clinical Endocrinology</i> , 1989 , 30, 619-25	3.4	76
6	A hyperkinetic heart in uncomplicated active acromegaly. Explanation of hypertension in acromegalic patients?. <i>Acta Medica Scandinavica</i> , 1988 , 223, 337-43		52
5	Echocardiographic-determined left ventricular wall characteristics in insulin-dependent diabetic patients. <i>Acta Medica Scandinavica</i> , 1988 , 224, 343-8		18
4	The Heart in Diabetes in Early and Advanced Nephropathy 1988 , 303-311		
3	Beneficial effect of a low-fat low-calorie diet on myocardial energy metabolism in patients with angina pectoris. <i>Lancet, The</i> , 1984 , 2, 59-62	40	9

2 Bifurcation Lesions298-311

1 The Nordic Experience21-33