Ali Rza Trkoglu

List of Publications by Year in Descending Order

Source: https://exaly.com/author-pdf/7625959/ali-riza-turkoglu-publications-by-year.pdf

Version: 2024-04-28

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

9	30	3	5
papers	citations	h-index	g-index
10	35 ext. citations	1.9	1.13
ext. papers		avg, IF	L-index

#	Paper	IF	Citations
9	Tofacitinib monotherapy and erectile dysfunction in rheumatoid arthritis: a pilot observational study <i>Rheumatology International</i> , 2022 , 1	3.6	
8	Geriatric giants in women over 65 years living in a rural area in Turkey. <i>Journal of Women and Aging</i> , 2021 , 33, 676-682	1.4	
7	Comparison of the effects of trospium and solifenacin on intraocular pressure and tear secretion in patients with overactive bladder. <i>International Urogynecology Journal</i> , 2021 , 32, 173-178	2	
6	Can continuous positive airway pressure improve lower urinary tract symptoms and erectile dysfunction in male patients with severe obstructive sleep apnea syndrome?. <i>Investigative and Clinical Urology</i> , 2020 , 61, 607-612	1.9	2
5	Thiol-disulphide balance in infertility secondary to varicocele. <i>Andrologia</i> , 2019 , 51, e13300	2.4	1
4	Effect of cigarette smoking on sexual functions, psychological factors, and disease activity in male patients with ankylosing spondylitis. <i>Aging Male</i> , 2019 , 22, 109-115	2.1	6
3	Evaluation of the relationship between compliance with the follow-up and treatment protocol and health literacy in bladder tumor patients. <i>Aging Male</i> , 2019 , 22, 266-271	2.1	11
2	Effect of trospium chloride therapy on intraocular pressure and tear secretion in overactive bladder patients. <i>Cutaneous and Ocular Toxicology</i> , 2017 , 36, 331-335	1.8	3
1	Changes in intraocular pressure and tear secretion in patients given 5 mg solifenacin for the treatment of overactive bladder. <i>International Urogynecology Journal</i> , 2017 , 28, 777-781	2	6