
## Kirsti Skavberg Roaldsen

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/7628185/publications.pdf Version: 2024-02-01



#	Article	IF	CITATIONS
1	Intraclass correlation $\hat{a} \in A$ discussion and demonstration of basic features. PLoS ONE, 2019, 14, e0219854.	2.5	375
2	High incidence of falls and fall-related injuries in wheelchair users with spinal cord injury: A prospective study of risk indicators. Journal of Rehabilitation Medicine, 2017, 49, 144-151.	1.1	44
3	Factors Associated With Recurrent Falls in Individuals With Traumatic Spinal Cord Injury: A Multicenter Study. Archives of Physical Medicine and Rehabilitation, 2016, 97, 1908-1916.	0.9	40
4	Comparison of the Berg Balance Scale and the Mini-BESTest for Assessing Balance in Ambulatory People With Spinal Cord Injury: Validation Study. Physical Therapy, 2017, 97, 677-687.	2.4	39
5	Cognitive Performance in Multiple Trauma Patients 3 Years After Injury. Psychosomatic Medicine, 1999, 61, 576-583.	2.0	35
6	Functional ability in female leg ulcer patients — a challenge for physiotherapy. Physiotherapy Research International, 2006, 11, 191-203.	1.5	32
7	Physical activity in patients with venous leg ulcer – between engagement and avoidance. A patient perspective. Clinical Rehabilitation, 2011, 25, 275-286.	2.2	31
8	Fearâ€avoidance beliefs and pain as predictors for low physical activity in patients with leg ulcer. Physiotherapy Research International, 2009, 14, 167-180.	1.5	29
9	Negotiating identity and self-image: perceptions of falls in ambulatory individuals with spinal cord injury – a qualitative study. Clinical Rehabilitation, 2017, 31, 544-554.	2.2	25
10	"Pushing the Limits― Rethinking Motor and Cognitive Resources After a Highly Challenging Balance Training Program for Parkinson Disease. Physical Therapy, 2017, 97, 81-89.	2.4	20
11	Task-specific balance training improves self-assessed function in community-dwelling older adults with balance deficits and fear of falling: a randomized controlled trial. Clinical Rehabilitation, 2014, 28, 1189-1197.	2.2	17
12	Self-reported function and disability in late life – cross-cultural adaptation and validation of the Swedish version of the late-life function and disability instrument. Disability and Rehabilitation, 2014, 36, 813-817.	1.8	17
13	Bridging the gap between research-based knowledge and clinical practice: a qualitative examination of patients and physiotherapists' views on the Otago exercise Programme. BMC Geriatrics, 2019, 19, 278.	2.7	15
14	"You Have to Keep Moving, Be Active― Perceptions and Experiences of Habitual Physical Activity in Older Women With Osteoporosis. Physical Therapy, 2016, 96, 361-370.	2.4	13
15	Predictors of falls in persons with spinal cord injury—a prospective study using the Downton fall risk index and a single question of previous falls. Spinal Cord, 2019, 57, 91-99.	1.9	11
16	"Fear of falling serves as protection and signifies potential danger― a qualitative study to conceptualise the phrase "fear of falling―in women with osteoporosis. Osteoporosis International, 2021, 32, 2563-2570.	3.1	10
17	"Better safe than sorry†a qualitative content analysis of participant's perspectives of fall-related concerns and balance in older women with osteoporosis after balance training. Disability and Rehabilitation, 2016, 38, 796-802.	1.8	9
18	Otago exercise programme—from evidence to practice: a qualitative study of physiotherapists' perceptions of the importance of organisational factors of leadership, context and culture for knowledge translation in Norway. BMC Health Services Research, 2020, 20, 985.	2.2	9

#	Article	IF	CITATIONS
19	Test-retest reliability at the item level and total score level of the Norwegian version of the Spinal Cord Injury Falls Concern Scale (SCI-FCS). Journal of Spinal Cord Medicine, 2016, 39, 317-326.	1.4	8
20	Controlling the Uncontrollable: Perceptions of Balance in People With Parkinson Disease. Physical Therapy, 2019, 99, 1501-1510.	2.4	8
21	Physiotherapists' perceptions of challenges facing evidence-based practice and the importance of environmental empowerment in fall prevention in the municipality – a qualitative study. BMC Geriatrics, 2020, 20, 432.	2.7	7
22	"Stay balanced―– effectiveness of evidence-based balance training for older adults transferred into a physical therapy primary care setting – a pilot study. Disability and Rehabilitation, 2020, 42, 1797-1802.	1.8	4
23	Reflections of older people about their experience of fall prevention exercise in the community- a qualitative study exploring evidence-based practice. BMC Public Health, 2020, 20, 1671.	2.9	4
24	Perceptions of balance and falls following a supervised training intervention – a qualitative study of people with Parkinson's disease. Disability and Rehabilitation, 2019, 41, 934-940.	1.8	3
25	Clinical supervisors' experience of giving feedback to students during clinical integrated learning. Physiotherapy Theory and Practice, 2022, 38, 122-131.	1.3	3
26	Pediatric spinal cord injury rehabilitation: A protocol for an international multicenter project (SINpedSCI). Journal of Pediatric Rehabilitation Medicine, 2022, 15, 395-403.	0.5	3
27	Carboxy terminal collagen crosslinks as a prognostic risk factor for fall-related fractures in individuals with established spinal cord injury. Spinal Cord, 2019, 57, 985-991.	1.9	2
28	Patient-specific functioning related to dizziness and balance problems after traumatic brain injury – A cross sectional study using an ICF perspective. Cogent Medicine, 2021, 8, .	0.7	2
29	"A manageable and challenging fall prevention intervention with impact on societyâ€⊷ older women's perspectives on participation in the stayBalanced training programme. Physiotherapy Theory and Practice, 2022, 38, 2806-2816.	1.3	1
30	Organisation of services and systems of care in paediatric spinal cord injury rehabilitation in seven countries: a survey with a descriptive cross-sectional design. Spinal Cord, 2022, 60, 339-347.	1.9	1