

Kirsti Skavberg Roaldsen

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/7628185/publications.pdf>

Version: 2024-02-01

30
papers

817
citations

759233

12
h-index

526287

27
g-index

30
all docs

30
docs citations

30
times ranked

954
citing authors

| # | ARTICLE | IF | CITATIONS |
|----|---|-----|-----------|
| 1 | Clinical supervisorsâ€™ experience of giving feedback to students during clinical integrated learning. <i>Physiotherapy Theory and Practice</i> , 2022, 38, 122-131. | 1.3 | 3 |
| 2 | â€œA manageable and challenging fall prevention intervention with impact on societyâ€• older womenâ€™s perspectives on participation in the stayBalanced training programme. <i>Physiotherapy Theory and Practice</i> , 2022, 38, 2806-2816. | 1.3 | 1 |
| 3 | Organisation of services and systems of care in paediatric spinal cord injury rehabilitation in seven countries: a survey with a descriptive cross-sectional design. <i>Spinal Cord</i> , 2022, 60, 339-347. | 1.9 | 1 |
| 4 | Pediatric spinal cord injury rehabilitation: A protocol for an international multicenter project (SINpedSCI). <i>Journal of Pediatric Rehabilitation Medicine</i> , 2022, 15, 395-403. | 0.5 | 3 |
| 5 | â€œFear of falling serves as protection and signifies potential dangerâ€• a qualitative study to conceptualise the phrase â€œfear of fallingâ€• in women with osteoporosis. <i>Osteoporosis International</i> , 2021, 32, 2563-2570. | 3.1 | 10 |
| 6 | Patient-specific functioning related to dizziness and balance problems after traumatic brain injury â€• A cross sectional study using an ICF perspective. <i>Cogent Medicine</i> , 2021, 8, . | 0.7 | 2 |
| 7 | â€œStay balancedâ€• effectiveness of evidence-based balance training for older adults transferred into a physical therapy primary care setting â€• a pilot study. <i>Disability and Rehabilitation</i> , 2020, 42, 1797-1802. | 1.8 | 4 |
| 8 | Reflections of older people about their experience of fall prevention exercise in the community- a qualitative study exploring evidence-based practice. <i>BMC Public Health</i> , 2020, 20, 1671. | 2.9 | 4 |
| 9 | Physiotherapistsâ€™ perceptions of challenges facing evidence-based practice and the importance of environmental empowerment in fall prevention in the municipality â€• a qualitative study. <i>BMC Geriatrics</i> , 2020, 20, 432. | 2.7 | 7 |
| 10 | Otago exercise programmeâ€™from evidence to practice: a qualitative study of physiotherapistsâ€™ perceptions of the importance of organisational factors of leadership, context and culture for knowledge translation in Norway. <i>BMC Health Services Research</i> , 2020, 20, 985. | 2.2 | 9 |
| 11 | Predictors of falls in persons with spinal cord injuryâ€™a prospective study using the Downton fall risk index and a single question of previous falls. <i>Spinal Cord</i> , 2019, 57, 91-99. | 1.9 | 11 |
| 12 | Intraclass correlation â€• A discussion and demonstration of basic features. <i>PLoS ONE</i> , 2019, 14, e0219854. | 2.5 | 375 |
| 13 | Bridging the gap between research-based knowledge and clinical practice: a qualitative examination of patients and physiotherapistsâ€™ views on the Otago exercise Programme. <i>BMC Geriatrics</i> , 2019, 19, 278. | 2.7 | 15 |
| 14 | Carboxy terminal collagen crosslinks as a prognostic risk factor for fall-related fractures in individuals with established spinal cord injury. <i>Spinal Cord</i> , 2019, 57, 985-991. | 1.9 | 2 |
| 15 | Controlling the Uncontrollable: Perceptions of Balance in People With Parkinson Disease. <i>Physical Therapy</i> , 2019, 99, 1501-1510. | 2.4 | 8 |
| 16 | Perceptions of balance and falls following a supervised training intervention â€• a qualitative study of people with Parkinsonâ€™s disease. <i>Disability and Rehabilitation</i> , 2019, 41, 934-940. | 1.8 | 3 |
| 17 | â€œPushing the Limitsâ€• Rethinking Motor and Cognitive Resources After a Highly Challenging Balance Training Program for Parkinson Disease. <i>Physical Therapy</i> , 2017, 97, 81-89. | 2.4 | 20 |
| 18 | Negotiating identity and self-image: perceptions of falls in ambulatory individuals with spinal cord injury â€• a qualitative study. <i>Clinical Rehabilitation</i> , 2017, 31, 544-554. | 2.2 | 25 |

| # | ARTICLE | IF | CITATIONS |
|----|---|-----|-----------|
| 19 | Comparison of the Berg Balance Scale and the Mini-BESTest for Assessing Balance in Ambulatory People With Spinal Cord Injury: Validation Study. <i>Physical Therapy</i> , 2017, 97, 677-687. | 2.4 | 39 |
| 20 | High incidence of falls and fall-related injuries in wheelchair users with spinal cord injury: A prospective study of risk indicators. <i>Journal of Rehabilitation Medicine</i> , 2017, 49, 144-151. | 1.1 | 44 |
| 21 | Test-retest reliability at the item level and total score level of the Norwegian version of the Spinal Cord Injury Falls Concern Scale (SCI-FCS). <i>Journal of Spinal Cord Medicine</i> , 2016, 39, 317-326. | 1.4 | 8 |
| 22 | Factors Associated With Recurrent Falls in Individuals With Traumatic Spinal Cord Injury: A Multicenter Study. <i>Archives of Physical Medicine and Rehabilitation</i> , 2016, 97, 1908-1916. | 0.9 | 40 |
| 23 | “Better safe than sorry” a qualitative content analysis of participant's perspectives of fall-related concerns and balance in older women with osteoporosis after balance training. <i>Disability and Rehabilitation</i> , 2016, 38, 796-802. | 1.8 | 9 |
| 24 | “You Have to Keep Moving, Be Active” Perceptions and Experiences of Habitual Physical Activity in Older Women With Osteoporosis. <i>Physical Therapy</i> , 2016, 96, 361-370. | 2.4 | 13 |
| 25 | Task-specific balance training improves self-assessed function in community-dwelling older adults with balance deficits and fear of falling: a randomized controlled trial. <i>Clinical Rehabilitation</i> , 2014, 28, 1189-1197. | 2.2 | 17 |
| 26 | Self-reported function and disability in late life “ cross-cultural adaptation and validation of the Swedish version of the late-life function and disability instrument. <i>Disability and Rehabilitation</i> , 2014, 36, 813-817. | 1.8 | 17 |
| 27 | Physical activity in patients with venous leg ulcer “ between engagement and avoidance. A patient perspective. <i>Clinical Rehabilitation</i> , 2011, 25, 275-286. | 2.2 | 31 |
| 28 | Fear avoidance beliefs and pain as predictors for low physical activity in patients with leg ulcer. <i>Physiotherapy Research International</i> , 2009, 14, 167-180. | 1.5 | 29 |
| 29 | Functional ability in female leg ulcer patients “ a challenge for physiotherapy. <i>Physiotherapy Research International</i> , 2006, 11, 191-203. | 1.5 | 32 |
| 30 | Cognitive Performance in Multiple Trauma Patients 3 Years After Injury. <i>Psychosomatic Medicine</i> , 1999, 61, 576-583. | 2.0 | 35 |