

Maria Oliveira-Souza

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/7634646/publications.pdf>

Version: 2024-02-01

29
papers

437
citations

706676

14
h-index

843174

20
g-index

29
all docs

29
docs citations

29
times ranked

683
citing authors

#	ARTICLE	IF	CITATIONS
1	The Role of Tamm-Horsfall Protein in the Pathogenesis of Crystalline Nephropathy. <i>FASEB Journal</i> , 2022, 36, .	0.2	0
2	Long-Term Angiotensin II Infusion Induces Oxidative and Endoplasmic Reticulum Stress and Modulates Na ⁺ Transporters Through the Nephron. <i>Frontiers in Physiology</i> , 2021, 12, 642752.	1.3	9
3	SRT1720 Administration Attenuates Albuminuria and Renal Fibrosis on Adriamycin-Induced Glomerulosclerosis. <i>FASEB Journal</i> , 2021, 35, .	0.2	0
4	Early type 1 diabetes aggravates renal ischemia/reperfusion-induced acute kidney injury. <i>Scientific Reports</i> , 2021, 11, 19028.	1.6	11
5	Sodium Oxalate-Induced Acute Kidney Injury Associated With Glomerular and Tubulointerstitial Damage in Rats. <i>Frontiers in Physiology</i> , 2020, 11, 1076.	1.3	8
6	Signaling pathways involved in the rapid biphasic effect of aldosterone on Na ⁺ /H ⁺ exchanger in rat proximal tubule cells. <i>Journal of Steroid Biochemistry and Molecular Biology</i> , 2018, 182, 87-94.	1.2	2
7	Intracellular albumin overload elicits endoplasmic reticulum stress and PKC- δ /p38 MAPK pathway activation to induce podocyte apoptosis. <i>Scientific Reports</i> , 2018, 8, 18012.	1.6	27
8	Angiotensin II-induced podocyte apoptosis is mediated by endoplasmic reticulum stress/PKC- δ /p38 MAPK pathway activation and through increased Na ⁺ /H ⁺ exchanger isoform 1 activity. <i>BMC Nephrology</i> , 2018, 19, 179.	0.8	43
9	The Effect of Albumin on Podocyte Apoptosis: the Role of PKC δ , p38 MAPK and Endoplasmic Reticulum Stress. <i>FASEB Journal</i> , 2018, 32, .	0.2	0
10	The Role of β -Adrenergic Overstimulation in the Early Stages of Renal Injury. <i>Kidney and Blood Pressure Research</i> , 2017, 42, 1277-1289.	0.9	13
11	Beta-2-microglobulin (B2M) expression in the urinary sediment correlates with clinical markers of kidney disease in patients with type 1 diabetes. <i>Metabolism: Clinical and Experimental</i> , 2016, 65, 816-824.	1.5	24
12	N-Acetyl Cysteine Attenuated the Deleterious Effects of Advanced Glycation End-Products on the Kidney of Non-Diabetic Rats. <i>Cellular Physiology and Biochemistry</i> , 2016, 40, 608-620.	1.1	9
13	Renovascular remodeling and renal injury after extended angiotensin II infusion. <i>American Journal of Physiology - Renal Physiology</i> , 2016, 310, F1295-F1307.	1.3	27
14	Thioredoxin interacting protein expression in the urinary sediment associates with renal function decline in type 1 diabetes. <i>Free Radical Research</i> , 2016, 50, 101-110.	1.5	23
15	Proximal tubule NHE3 activity is inhibited by beta-arrestin-biased angiotensin II type 1 receptor signaling. <i>American Journal of Physiology - Cell Physiology</i> , 2015, 309, C541-C550.	2.1	15
16	Renal Hemodynamic and Morphological Changes after 7 and 28 Days of Leptin Treatment: The Participation of Angiotensin II via the AT1 Receptor. <i>PLoS ONE</i> , 2015, 10, e0122265.	1.1	15
17	High Glucose Concentration Stimulates NHE-1 Activity in Distal Nephron Cells: the Role of the Mek/Erk1/2/p90 ^{RSK} and p38MAPK Signaling Pathways. <i>Cellular Physiology and Biochemistry</i> , 2014, 33, 333-343.	1.1	25
18	The regulation of NHE1 and NHE3 activity by angiotensin II is mediated by the activation of the angiotensin II type I receptor/phospholipase C/calcium/calmodulin pathway in distal nephron cells. <i>European Journal of Pharmacology</i> , 2013, 721, 322-331.	1.7	26

#	ARTICLE	IF	CITATIONS
19	Glucose-Induced Regulation of NHEs Activity and SGLTs Expression Involves the PKA Signaling Pathway. Journal of Membrane Biology, 2011, 239, 157-165.	1.0	37
20	Regulation of Na ⁺ /H ⁺ Exchanger Isoform 1 (NHE1) by Calmodulin-binding Sites: Role of Angiotensin II. Cellular Physiology and Biochemistry, 2010, 26, 541-552.	1.1	16
21	Renal Function of Rats with Isoproterenol-induced Cardiac Hypertrophy. FASEB Journal, 2010, 24, 1059-20.	0.2	0
22	The role of kidney on sodium balance in Angiotensin II-induced hypertension.. FASEB Journal, 2010, 24, 605-9.	0.2	0
23	The effect of angiotensin II on intracellular pH is mediated by AT1 receptor translocation. American Journal of Physiology - Cell Physiology, 2008, 295, C138-C145.	2.1	8
24	Insulin but Not Phlorizin Treatment Induces a Transient Increase in GLUT2 Gene Expression in the Kidney of Diabetic Rats. Nephron Physiology, 2007, 105, p42-p51.	1.5	23
25	Signaling Pathways in the Biphasic Effect of ANG II on Na ⁺ /H ⁺ Exchanger in T84 Cells. Journal of Membrane Biology, 2005, 205, 49-60.	1.0	6
26	Arginine vasopressin stimulates H ⁺ -ATPase in MDCK cells via V1 (cell Ca ²⁺) and V2 (cAMP) receptors. American Journal of Physiology - Renal Physiology, 2004, 286, F402-F408.	1.3	21
27	Atrial natriuretic peptide impairs the stimulatory effect of angiotensin II on H ⁺ -ATPase. Kidney International, 2002, 62, 1693-1699.	2.6	23
28	Effect of arginine vasopressin and ANP on intracellular pH and cytosolic free [Ca ²⁺] regulation in MDCK cells. Kidney International, 2001, 60, 1800-1808.	2.6	11
29	Na ⁺ -independent proton secretion in MDCK-C11 cells. Pflugers Archiv European Journal of Physiology, 2000, 441, 287-293.	1.3	15