

Xiao Qun Wang

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/7639320/publications.pdf>

Version: 2024-02-01

20
papers

575
citations

623699

14
h-index

752679

20
g-index

27
all docs

27
docs citations

27
times ranked

842
citing authors

#	ARTICLE	IF	CITATIONS
1	C1q/TNF-related protein-1: an adipokine marking and promoting atherosclerosis. <i>European Heart Journal</i> , 2016, 37, 1762-1771.	2.2	75
2	Advanced glycation end products induce chemokine/cytokine production via activation of p38 pathway and inhibit proliferation and migration of bone marrow mesenchymal stem cells. <i>Cardiovascular Diabetology</i> , 2010, 9, 66.	6.8	56
3	Reduced coronary collateralization in type 2 diabetic patients with chronic total occlusion. <i>Cardiovascular Diabetology</i> , 2018, 17, 26.	6.8	55
4	Deletion of BACH1 Attenuates Atherosclerosis by Reducing Endothelial Inflammation. <i>Circulation Research</i> , 2022, 130, 1038-1055.	4.5	55
5	C1q/TNF-related protein 1 links macrophage lipid metabolism to inflammation and atherosclerosis. <i>Atherosclerosis</i> , 2016, 250, 38-45.	0.8	43
6	CTRP5 promotes transcytosis and oxidative modification of low-density lipoprotein and the development of atherosclerosis. <i>Atherosclerosis</i> , 2018, 278, 197-209.	0.8	41
7	The Role of Monocyte to High-Density Lipoprotein Cholesterol Ratio in Prediction of Carotid Intima-Media Thickness in Patients With Type 2 Diabetes. <i>Frontiers in Endocrinology</i> , 2019, 10, 191.	3.5	37
8	Receptor Mediated Elevation in FABP4 Levels by Advanced Glycation End Products Induces Cholesterol and Triacylglycerol Accumulation in THP- α 1 Macrophages. <i>Lipids</i> , 2011, 46, 479-486.	1.7	28
9	Glycated Apolipoprotein A-IV Induces Atherogenesis in Patients With CAD in Type 2 Diabetes. <i>Journal of the American College of Cardiology</i> , 2017, 70, 2006-2019.	2.8	28
10	Increased serum TREM-1 level is associated with in-stent restenosis, and activation of TREM-1 promotes inflammation, proliferation and migration in vascular smooth muscle cells. <i>Atherosclerosis</i> , 2017, 267, 10-18.	0.8	24
11	C1q/TNF-related protein 1 promotes endothelial barrier dysfunction under disturbed flow. <i>Biochemical and Biophysical Research Communications</i> , 2017, 490, 580-586.	2.1	23
12	Insulin resistance and dysglycemia are associated with left ventricular remodeling after myocardial infarction in non-diabetic patients. <i>Cardiovascular Diabetology</i> , 2019, 18, 100.	6.8	22
13	Visit-to-visit HbA1c variability is associated with in-stent restenosis in patients with type 2 diabetes after percutaneous coronary intervention. <i>Cardiovascular Diabetology</i> , 2020, 19, 133.	6.8	21
14	Lipoprotein (a) interactions with cholesterol-containing lipids on angiographic coronary collateralization in type 2 diabetic patients with chronic total occlusion. <i>Cardiovascular Diabetology</i> , 2019, 18, 82.	6.8	17
15	Visit-to-visit fasting plasma glucose variability is associated with left ventricular adverse remodeling in diabetic patients with STEMI. <i>Cardiovascular Diabetology</i> , 2020, 19, 131.	6.8	15
16	Searching for optimal blood pressure targets in type 2 diabetic patients with coronary artery disease. <i>Cardiovascular Diabetology</i> , 2019, 18, 160.	6.8	14
17	Relationship of High-Density Lipoprotein-Associated Arylesterase Activity to Systolic Heart Failure in Patients with and without Type 2 Diabetes. <i>Scientific Reports</i> , 2019, 9, 5979.	3.3	8
18	Donor artery stenosis interactions with diastolic blood pressure on coronary collateral flow in type 2 diabetic patients with chronic total occlusion. <i>Cardiovascular Diabetology</i> , 2018, 17, 76.	6.8	7

#	ARTICLE	IF	CITATIONS
19	Data on the expression and role of TREM-1 in the development of in-stent restenosis. Data in Brief, 2018, 16, 604-607.	1.0	3
20	Using En Face Immunofluorescence Staining to Observe Vascular Endothelial Cells Directly. Journal of Visualized Experiments, 2019, , .	0.3	3