

Xiao Qun Wang

List of Publications by Citations

Source: <https://exaly.com/author-pdf/7639320/xiao-qun-wang-publications-by-citations.pdf>

Version: 2024-04-28

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

19
papers

343
citations

11
h-index

18
g-index

27
ext. papers

476
ext. citations

5.9
avg, IF

2.98
L-index

| # | Paper | IF | Citations |
|----|---|------|-----------|
| 19 | C1q/TNF-related protein-1: an adipokine marking and promoting atherosclerosis. <i>European Heart Journal</i> , 2016 , 37, 1762-71 | 9.5 | 55 |
| 18 | Advanced glycation end products induce chemokine/cytokine production via activation of p38 pathway and inhibit proliferation and migration of bone marrow mesenchymal stem cells. <i>Cardiovascular Diabetology</i> , 2010 , 9, 66 | 8.7 | 48 |
| 17 | Reduced coronary collateralization in type 2 diabetic patients with chronic total occlusion. <i>Cardiovascular Diabetology</i> , 2018 , 17, 26 | 8.7 | 38 |
| 16 | C1q/TNF-related protein 1 links macrophage lipid metabolism to inflammation and atherosclerosis. <i>Atherosclerosis</i> , 2016 , 250, 38-45 | 3.1 | 31 |
| 15 | CTRP5 promotes transcytosis and oxidative modification of low-density lipoprotein and the development of atherosclerosis. <i>Atherosclerosis</i> , 2018 , 278, 197-209 | 3.1 | 24 |
| 14 | Receptor mediated elevation in FABP4 levels by advanced glycation end products induces cholesterol and triacylglycerol accumulation in THP-1 macrophages. <i>Lipids</i> , 2011 , 46, 479-86 | 1.6 | 21 |
| 13 | Glycated Apolipoprotein A-IV Induces Atherogenesis in Patients With CAD in Type 2 Diabetes. <i>Journal of the American College of Cardiology</i> , 2017 , 70, 2006-2019 | 15.1 | 20 |
| 12 | C1q/TNF-related protein 1 promotes endothelial barrier dysfunction under disturbed flow. <i>Biochemical and Biophysical Research Communications</i> , 2017 , 490, 580-586 | 3.4 | 18 |
| 11 | Increased serum TREM-1 level is associated with in-stent restenosis, and activation of TREM-1 promotes inflammation, proliferation and migration in vascular smooth muscle cells. <i>Atherosclerosis</i> , 2017 , 267, 10-18 | 3.1 | 16 |
| 10 | The Role of Monocyte to High-Density Lipoprotein Cholesterol Ratio in Prediction of Carotid Intima-Media Thickness in Patients With Type 2 Diabetes. <i>Frontiers in Endocrinology</i> , 2019 , 10, 191 | 5.7 | 14 |
| 9 | Insulin resistance and dysglycemia are associated with left ventricular remodeling after myocardial infarction in non-diabetic patients. <i>Cardiovascular Diabetology</i> , 2019 , 18, 100 | 8.7 | 11 |
| 8 | Lipoprotein (a) interactions with cholesterol-containing lipids on angiographic coronary collateralization in type 2 diabetic patients with chronic total occlusion. <i>Cardiovascular Diabetology</i> , 2019 , 18, 82 | 8.7 | 9 |
| 7 | Visit-to-visit HbA variability is associated with in-stent restenosis in patients with type 2 diabetes after percutaneous coronary intervention. <i>Cardiovascular Diabetology</i> , 2020 , 19, 133 | 8.7 | 9 |
| 6 | Visit-to-visit fasting plasma glucose variability is associated with left ventricular adverse remodeling in diabetic patients with STEMI. <i>Cardiovascular Diabetology</i> , 2020 , 19, 131 | 8.7 | 7 |
| 5 | Searching for optimal blood pressure targets in type 2 diabetic patients with coronary artery disease. <i>Cardiovascular Diabetology</i> , 2019 , 18, 160 | 8.7 | 7 |
| 4 | Relationship of High-Density Lipoprotein-Associated Arylesterase Activity to Systolic Heart Failure in Patients with and without Type 2 Diabetes. <i>Scientific Reports</i> , 2019 , 9, 5979 | 4.9 | 6 |
| 3 | Donor artery stenosis interactions with diastolic blood pressure on coronary collateral flow in type 2 diabetic patients with chronic total occlusion. <i>Cardiovascular Diabetology</i> , 2018 , 17, 76 | 8.7 | 6 |

- | | | | |
|---|---|------|---|
| 2 | Data on the expression and role of TREM-1 in the development of in-stent restenosis. <i>Data in Brief</i> , 2018 , 16, 604-607 | 1.2 | 2 |
| 1 | Deletion of BACH1 Attenuates Atherosclerosis by Reducing Endothelial Inflammation.. <i>Circulation Research</i> , 2022 , CIRCRESAHA121319540 | 15.7 | 1 |