
Maciej Boleslaw Olszewski

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/7641398/publications.pdf Version: 2024-02-01



#	Article	IF	CITATIONS
1	Chemotherapy of HER2- and MDM2-Enriched Breast Cancer Subtypes Induces Homologous Recombination DNA Repair and Chemoresistance. Cancers, 2021, 13, 4501.	1.7	2
2	Gain-of-Function Mutations in p53 in Cancer Invasiveness and Metastasis. International Journal of Molecular Sciences, 2020, 21, 1334.	1.8	20
3	Zebrafish Microenvironment Elevates EMT and CSC-Like Phenotype of Engrafted Prostate Cancer Cells. Cells, 2020, 9, 797.	1.8	12
4	Diverse and cancer type‑specific roles of the p53 R248Q gain‑of‑function mutation in cancer migration and invasiveness. International Journal of Oncology, 2019, 54, 1168-1182.	1.4	12
5	Molecular chaperones in the acquisition of cancer cell chemoresistance with mutated <i>TP53</i> and MDM2 up-regulation. Oncotarget, 2017, 8, 82123-82143.	0.8	29
6	The clathrin-binding and J-domains of GAK support the uncoating and chaperoning of clathrin by Hsc70 in the brain. Journal of Cell Science, 2015, 128, 3811-21.	1.2	23
7	Unbiased screen for interactors of leucine-rich repeat kinase 2 supports a common pathway for sporadic and familial Parkinson disease. Proceedings of the National Academy of Sciences of the United States of America, 2014, 111, 2626-2631.	3.3	342
8	Disruption of Clathrinâ€Mediated Trafficking Causes Centrosome Overduplication and Senescence. Traffic, 2014, 15, 60-77.	1.3	16
9	Molecular Mechanism of Mutant p53 Stabilization: The Role of HSP70 and MDM2. PLoS ONE, 2012, 7, e51426.	1.1	105
10	Basophils and mast cells: Underdog in immune regulation?. Immunology Letters, 2011, 138, 28-31.	1.1	19
11	Hsp70 molecular chaperones are required to support p53 tumor suppressor activity under stress conditions. Oncogene, 2009, 28, 4284-4294.	2.6	75
12	TNF Trafficking to Human Mast Cell Granules: Mature Chain-Dependent Endocytosis. Journal of Immunology, 2007, 178, 5701-5709.	0.4	132
13	Efficient sorting of TNF-alpha to rodent mast cell granules is dependent on N-linked glycosylation. European Journal of Immunology, 2006, 36, 997-1008.	1.6	26
14	Development of the ?Cell Chip?: a new in vitro alternative technique for immunotoxicity testing. Toxicology, 2005, 206, 245-256.	2.0	19
15	"Fluorescent Cell Chip―for immunotoxicity testing: Development of the c- expression reporter cell lines. Toxicology and Applied Pharmacology, 2005, 207, 133-141.	1.3	3
16	Detection of immunotoxicity using T-cell based cytokine reporter cell lines (?Cell Chip?). Toxicology, 2005, 206, 257-272.	2.0	23