

# Lie Dai

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/764194/publications.pdf>

Version: 2024-02-01

39  
papers

664  
citations

706676

14  
h-index

685536

24  
g-index

43  
all docs

43  
docs citations

43  
times ranked

1028  
citing authors

#	ARTICLE	IF	CITATIONS
1	Euphoesulatin A prevents osteoclast differentiation and bone loss via inhibiting RANKL-induced ROS production and NF- $\kappa$ B and MAPK signal pathways. <i>Bioorganic Chemistry</i> , 2022, 119, 105511.	2.0	5
2	Fine Comparison of the Efficacy and Safety Between GB242 and Infliximab in Patients with Rheumatoid Arthritis: A Phase III Study. <i>Rheumatology and Therapy</i> , 2022, 9, 175-189.	1.1	2
3	A Matrix Prediction Model for the 6-Month Mortality Risk in Patients With Anti-Melanoma Differentiation-Associated Protein-5-Positive Dermatomyositis. <i>Frontiers in Medicine</i> , 2022, 9, 860798.	1.2	8
4	Association Between Metabolic Dysfunction-Associated Fatty Liver Disease and Cardiovascular Risk in Patients With Rheumatoid Arthritis: A Cross-Sectional Study of Chinese Cohort. <i>Frontiers in Cardiovascular Medicine</i> , 2022, 9, .	1.1	6
5	Low expression of estrogen receptor $\beta$ in renal tubular epithelial cells may cause hyperuricemia in premenopausal patients with systemic lupus erythematosus. <i>Lupus</i> , 2021, 30, 560-567.	0.8	13
6	Ultrasonography predicts the results of labial salivary gland biopsy in patients with suspected Sjögren's syndrome: a matrix risk model. <i>Therapeutic Advances in Musculoskeletal Disease</i> , 2021, 13, 1759720X2110105.	1.2	6
7	LncRNA THRIL is involved in the proliferation, migration, and invasion of rheumatoid fibroblast-like synoviocytes. <i>Annals of Translational Medicine</i> , 2021, 9, 1368-1368.	0.7	11
8	Addition of Fibroblast-Stromal Cell Markers to Immune Synovium Pathotypes Better Predicts Radiographic Progression at 1 Year in Active Rheumatoid Arthritis. <i>Frontiers in Immunology</i> , 2021, 12, 778480.	2.2	3
9	Gasdermin-E Mediated Pyroptosis: A Novel Mechanism Regulating Migration, Invasion and Release of Inflammatory Cytokines in Rheumatoid Arthritis Fibroblast-like Synoviocytes. <i>Frontiers in Cell and Developmental Biology</i> , 2021, 9, 810635.	1.8	12
10	Reduced skeletal muscle independently predicts 1-year aggravated joint destruction in patients with rheumatoid arthritis. <i>Therapeutic Advances in Musculoskeletal Disease</i> , 2020, 12, 1759720X2094622.	1.2	7
11	Neglected extra-articular manifestations in rheumatoid arthritis patients with normal body mass index: reduced skeletal muscle overlapping overfat. <i>Therapeutic Advances in Chronic Disease</i> , 2020, 11, 204062232097524.	1.1	3
12	Identification of Specific Joint-Inflammatogenic Cell-Free DNA Molecules From Synovial Fluids of Patients With Rheumatoid Arthritis. <i>Frontiers in Immunology</i> , 2020, 11, 662.	2.2	24
13	Myopenia is associated with joint damage in rheumatoid arthritis: a cross-sectional study. <i>Journal of Cachexia, Sarcopenia and Muscle</i> , 2019, 10, 355-367.	2.9	37
14	A novel function of artesunate on inhibiting migration and invasion of fibroblast-like synoviocytes from rheumatoid arthritis patients. <i>Arthritis Research and Therapy</i> , 2019, 21, 153.	1.6	71
15	Impaired Tip60-mediated Foxp3 acetylation attenuates regulatory T cell development in rheumatoid arthritis. <i>Journal of Autoimmunity</i> , 2019, 100, 27-39.	3.0	31
16	Activation of the Peroxisome Proliferator-Activated Receptor $\gamma$ Coactivator 1/NFATc1 Pathway in Circulating Osteoclast Precursors Associated With Bone Destruction in Rheumatoid Arthritis. <i>Arthritis and Rheumatology</i> , 2019, 71, 1252-1264.	2.9	18
17	FRI0226...LOW EXPRESSION OF ESTROGEN RECEPTOR BETA IN RENAL TUBULAR EPITHELIAL CELL MAY CONTRIBUTE TO HYPERURICEMIA IN PREMENOPAUNITED STATES OF AMERICAL FEMALE SYSTEMIC LUPUS ERYTHEMATOSUS PATIENTS. , 2019, , .		0
18	The value of MRI examination on bilateral hands including proximal interphalangeal joints for disease assessment in patients with early rheumatoid arthritis: a cross-sectional cohort study. <i>Arthritis Research and Therapy</i> , 2019, 21, 279.	1.6	7

#	ARTICLE	IF	CITATIONS
19	Genetic predictors of efficacy and toxicity of iguratimod in patients with rheumatoid arthritis. <i>Pharmacogenomics</i> , 2018, 19, 383-392.	0.6	8
20	Magnetic Resonance Imaging of Bilateral Hands Is More Optimal Than MRI of Unilateral Hands for Rheumatoid Arthritis. <i>Journal of Rheumatology</i> , 2018, 45, 895-904.	1.0	5
21	Elevated uric acid levels in premenopausal female systemic lupus erythematosus patients: Association with potential or existing renal damage. <i>European Journal of Inflammation</i> , 2018, 16, 205873921880244.	0.2	5
22	Presence of hepatitis B virus in synovium and its clinical significance in rheumatoid arthritis. <i>Arthritis Research and Therapy</i> , 2018, 20, 130.	1.6	14
23	Deleterious role of hepatitis B virus infection in therapeutic response among patients with rheumatoid arthritis in a clinical practice setting: a case-control study. <i>Arthritis Research and Therapy</i> , 2018, 20, 81.	1.6	8
24	Joint damage is amplified in rheumatoid arthritis patients with positive thyroid autoantibodies. <i>PeerJ</i> , 2018, 6, e4216.	0.9	10
25	A Novel Hypothesis on Excessive Activation of Residual B Lymphocytes in Common Variable Immunodeficiency Concurrent with Aseptic, Erosive Polyarthritis. <i>Medical Science Monitor</i> , 2018, 24, 4952-4960.	0.5	0
26	Short-course tocilizumab increases risk of hepatitis B virus reactivation in patients with rheumatoid arthritis: a prospective clinical observation. <i>International Journal of Rheumatic Diseases</i> , 2017, 20, 859-869.	0.9	89
27	Association Analysis of the MHC in Lupus Nephritis. <i>Journal of the American Society of Nephrology: JASN</i> , 2017, 28, 3383-3394.	3.0	21
28	Patients with Coexistence of Circulating Hepatitis B Surface Antigen and Its Antibody May Have a Strong Predisposition to Virus Reactivation During Immunosuppressive Therapy: A Hypothesis. <i>Medical Science Monitor</i> , 2017, 23, 5980-5985.	0.5	1
29	Continuously elevated serum matrix metalloproteinase-3 for 36 months predict one-year radiographic progression in rheumatoid arthritis: a prospective cohort study. <i>Arthritis Research and Therapy</i> , 2015, 17, 289.	1.6	34
30	The Smad3/Smad4/CDK9 complex promotes renal fibrosis in mice with unilateral ureteral obstruction. <i>Kidney International</i> , 2015, 88, 1323-1335.	2.6	18
31	Smad3 deficiency protects mice from obesity-induced podocyte injury that precedes insulin resistance. <i>Kidney International</i> , 2015, 88, 286-298.	2.6	39
32	Serum Matrix Metalloproteinase-3 as a Noninvasive Biomarker of Histological Synovitis for Diagnosis of Rheumatoid Arthritis. <i>Mediators of Inflammation</i> , 2014, 2014, 1-10.	1.4	57
33	Elevated Serum IgG4 Defines Specific Clinical Phenotype of Rheumatoid Arthritis. <i>Mediators of Inflammation</i> , 2014, 2014, 1-10.	1.4	30
34	Discontinuation of antiviral prophylaxis correlates with high prevalence of hepatitis B virus (HBV) reactivation in rheumatoid arthritis patients with HBV carrier state: a real-world clinical practice. <i>BMC Musculoskeletal Disorders</i> , 2014, 15, 449.	0.8	23
35	Effective treatment of Kimura's disease with leflunomide in combination with glucocorticoids. <i>Clinical Rheumatology</i> , 2011, 30, 859-865.	1.0	15
36	Elevated Serum Glucose-6-Phosphate Isomerase Correlates with Histological Disease Activity and Clinical Improvement After Initiation of Therapy in Patients with Rheumatoid Arthritis. <i>Journal of Rheumatology</i> , 2010, 37, 2452-2461.	1.0	16

#	ARTICLE	IF	CITATIONS
37	Cerebrospinal fluid and serum prolactin in systemic lupus erythematosus with and without central nervous system involvement. <i>APLAR Journal of Rheumatology</i> , 2007, 10, 37-42.	0.2	3
38	Cerebrospinal fluid and serum prolactin in systemic lupus erythematosus with and without central nervous system involvement. <i>APLAR Journal of Rheumatology</i> , 2006, 9, 232-237.	0.2	1
39	Comparison of the efficacy and safety of the adalimumab biosimilar TQ-Z2301 and adalimumab for the treatment of Chinese patients with active ankylosing spondylitis: a multi-center, randomized, double-blind, phase III clinical trial. <i>Clinical Rheumatology</i> , 0, , .	1.0	0