Lie Dai

List of Publications by Year in Descending Order

Source: https://exaly.com/author-pdf/764194/lie-dai-publications-by-year.pdf

Version: 2024-04-28

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

36	395	12	19
papers	citations	h-index	g-index
43	532	5.3	3.2
ext. papers	ext. citations	avg, IF	L-index

#	Paper	IF	Citations
36	A Matrix Prediction Model for the 6-Month Mortality Risk in Patients With Anti-Melanoma Differentiation-Associated Protein-5-Positive Dermatomyositis <i>Frontiers in Medicine</i> , 2022 , 9, 860798	4.9	O
35	Euphoesulatin A prevents osteoclast differentiation and bone loss via inhibiting RANKL-induced ROS production and NF- B and MAPK signal pathways. <i>Bioorganic Chemistry</i> , 2021 , 119, 105511	5.1	1
34	Addition of Fibroblast-Stromal Cell Markers to Immune Synovium Pathotypes Better Predicts Radiographic Progression at 1 Year in Active Rheumatoid Arthritis. <i>Frontiers in Immunology</i> , 2021 , 12, 778480	8.4	O
33	Fine Comparison of the Efficacy and Safety Between GB242 and Infliximab in Patients with Rheumatoid Arthritis: A Phase III Study. <i>Rheumatology and Therapy</i> , 2021 , 1	4.4	О
32	Low expression of estrogen receptor In renal tubular epithelial cells may cause hyperuricemia in premenopausal patients with systemic lupus erythematosus. <i>Lupus</i> , 2021 , 30, 560-567	2.6	2
31	Ultrasonography predicts the results of labial salivary gland biopsy in patients with suspected Sjgren's syndrome: a matrix risk model. <i>Therapeutic Advances in Musculoskeletal Disease</i> , 2021 , 13, 1759	7 ³ 28X2	19010592
30	LncRNA is involved in the proliferation, migration, and invasion of rheumatoid fibroblast-like synoviocytes. <i>Annals of Translational Medicine</i> , 2021 , 9, 1368	3.2	3
29	Gasdermin-E Mediated Pyroptosis-A Novel Mechanism Regulating Migration, Invasion and Release of Inflammatory Cytokines in Rheumatoid Arthritis Fibroblast-like Synoviocytes <i>Frontiers in Cell and Developmental Biology</i> , 2021 , 9, 810635	5.7	2
28	Neglected extra-articular manifestations in rheumatoid arthritis patients with normal body mass index: reduced skeletal muscle overlapping overfat. <i>Therapeutic Advances in Chronic Disease</i> , 2020 , 11, 2040622320975241	4.9	O
27	Identification of Specific Joint-Inflammatogenic Cell-Free DNA Molecules From Synovial Fluids of Patients With Rheumatoid Arthritis. <i>Frontiers in Immunology</i> , 2020 , 11, 662	8.4	14
26	Reduced skeletal muscle independently predicts 1-year aggravated joint destruction in patients with rheumatoid arthritis. <i>Therapeutic Advances in Musculoskeletal Disease</i> , 2020 , 12, 1759720X2094622	2 ∂ .8	3
25	Myopenia is associated with joint damage in rheumatoid arthritis: a cross-sectional study. <i>Journal of Cachexia, Sarcopenia and Muscle</i> , 2019 , 10, 355-367	10.3	20
24	A novel function of artesunate on inhibiting migration and invasion of fibroblast-like synoviocytes from rheumatoid arthritis patients. <i>Arthritis Research and Therapy</i> , 2019 , 21, 153	5.7	31
23	Impaired Tip60-mediated Foxp3 acetylation attenuates regulatory T cell development in rheumatoid arthritis. <i>Journal of Autoimmunity</i> , 2019 , 100, 27-39	15.5	19
22	Activation of the Peroxisome Proliferator-Activated Receptor Coactivator 1/NFATc1 Pathway in Circulating Osteoclast Precursors Associated With Bone Destruction in Rheumatoid Arthritis. <i>Arthritis and Rheumatology</i> , 2019 , 71, 1252-1264	9.5	9
21	The value of MRI examination on bilateral hands including proximal interphalangeal joints for disease assessment in patients with early rheumatoid arthritis: a cross-sectional cohort study. <i>Arthritis Research and Therapy</i> , 2019 , 21, 279	5.7	3
20	Genetic predictors of efficacy and toxicity of iguratimod in patients with rheumatoid arthritis. <i>Pharmacogenomics</i> , 2018 , 19, 383-392	2.6	8

19	Magnetic Resonance Imaging of Bilateral Hands Is More Optimal Than MRI of Unilateral Hands for Rheumatoid Arthritis. <i>Journal of Rheumatology</i> , 2018 , 45, 895-904	4.1	3
18	Presence of hepatitis B virus in synovium and its clinical significance in rheumatoid arthritis. <i>Arthritis Research and Therapy</i> , 2018 , 20, 130	5.7	7
17	Deleterious role of hepatitis B virus infection in therapeutic response among patients with rheumatoid arthritis in a clinical practice setting: a case-control study. <i>Arthritis Research and Therapy</i> , 2018 , 20, 81	5.7	3
16	Joint damage is amplified in rheumatoid arthritis patients with positive thyroid autoantibodies. <i>PeerJ</i> , 2018 , 6, e4216	3.1	1
15	A Novel Hypothesis on Excessive Activation of Residual B Lymphocytes in Common Variable Immunodeficiency Concurrent with Aseptic, Erosive Polyarthritis. <i>Medical Science Monitor</i> , 2018 , 24, 495	5 2:2 96	0
14	Elevated uric acid levels in premenopausal female systemic lupus erythematosus patients: Association with potential or existing renal damage. <i>European Journal of Inflammation</i> , 2018 , 16, 20587	3 ⁹ 218	80244
13	Short-course tocilizumab increases risk of hepatitis B virus reactivation in patients with rheumatoid arthritis: a prospective clinical observation. <i>International Journal of Rheumatic Diseases</i> , 2017 , 20, 859-86	6 3 .3	62
12	Association Analysis of the MHC in Lupus Nephritis. <i>Journal of the American Society of Nephrology: JASN</i> , 2017 , 28, 3383-3394	12.7	13
11	Patients with Coexistence of Circulating Hepatitis B Surface Antigen and Its Antibody May Have a Strong Predisposition to Virus Reactivation During Immunosuppressive Therapy: A Hypothesis. <i>Medical Science Monitor</i> , 2017 , 23, 5980-5985	3.2	1
10	The Smad3/Smad4/CDK9 complex promotes renal fibrosis in mice with unilateral ureteral obstruction. <i>Kidney International</i> , 2015 , 88, 1323-1335	9.9	9
9	Smad3 deficiency protects mice from obesity-induced podocyte injury that precedes insulin resistance. <i>Kidney International</i> , 2015 , 88, 286-98	9.9	33
8	Continuously elevated serum matrix metalloproteinase-3 for 3 ~ 6 months predict one-year radiographic progression in rheumatoid arthritis: a prospective cohort study. <i>Arthritis Research and Therapy</i> , 2015 , 17, 289	5.7	29
7	Serum matrix metalloproteinase-3 as a noninvasive biomarker of histological synovitis for diagnosis of rheumatoid arthritis. <i>Mediators of Inflammation</i> , 2014 , 2014, 179284	4.3	45
6	Elevated serum IgG4 defines specific clinical phenotype of rheumatoid arthritis. <i>Mediators of Inflammation</i> , 2014 , 2014, 635293	4.3	20
5	Discontinuation of antiviral prophylaxis correlates with high prevalence of hepatitis B virus (HBV) reactivation in rheumatoid arthritis patients with HBV carrier state: a real-world clinical practice. <i>BMC Musculoskeletal Disorders</i> , 2014 , 15, 449	2.8	20
4	Effective treatment of Kimurald disease with leflunomide in combination with glucocorticoids. <i>Clinical Rheumatology</i> , 2011 , 30, 859-65	3.9	12
3	Elevated serum glucose-6-phosphate isomerase correlates with histological disease activity and clinical improvement after initiation of therapy in patients with rheumatoid arthritis. <i>Journal of Rheumatology</i> , 2010 , 37, 2452-61	4.1	14
2	Cerebrospinal fluid and serum prolactin in systemic lupus erythematosus with and without central nervous system involvement. APLAR Journal of Rheumatology. 2007. 10, 37-42		3

Cerebrospinal fluid and serum prolactin in systemic lupus erythematosus with and without central nervous system involvement. *APLAR Journal of Rheumatology*, **2006**, 9, 232-237