

# Jean-Marie Casillas

## List of Publications by Year in Descending Order

**Source:** <https://exaly.com/author-pdf/7642060/jean-marie-casillas-publications-by-year.pdf>

**Version:** 2024-04-28

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

31  
papers

746  
citations

16  
h-index

27  
g-index

32  
ext. papers

846  
ext. citations

2.4  
avg, IF

3.3  
L-index

| #  | Paper  | IF  | Citations |
|----|--|-----|-----------|
| 31 | Assessing Gait After Changing Prosthetic Foot: Contribution of Subjective and Objective Measurements. A Pilot Study. <i>Journal of Prosthetics and Orthotics</i> , <b>2021</b> , 33, 146-150   | 0.7 |           |
| 30 | Training with FES-assisted cycling in a subject with spinal cord injury: Psychological, physical and physiological considerations. <i>Journal of Spinal Cord Medicine</i> , <b>2020</b> , 43, 402-413  | 1.9 | 5         |
| 29 | A tool to improve functional outcome assessment of a multimodal program for patients with chronic low back pain: A study on walk tests (at comfortable and fast speed). <i>Journal of Back and Musculoskeletal Rehabilitation</i> , <b>2020</b> , 33, 485-494  | 1.4 |           |
| 28 | Walking Speed as an Alternative Measure of Functional Status in Patients with Lumbar Spinal Stenosis. <i>World Neurosurgery</i> , <b>2019</b> , 122, e591-e597   | 2.1 | 8         |
| 27 | One year of training with FES has impressive beneficial effects in a 36-year-old woman with spinal cord injury. <i>Journal of Spinal Cord Medicine</i> , <b>2017</b> , 40, 107-112   | 1.9 | 18        |
| 26 | Assessment tools for personalizing training intensity during cardiac rehabilitation: Literature review and practical proposals. <i>Annals of Physical and Rehabilitation Medicine</i> , <b>2017</b> , 60, 43-49  | 3.8 | 16        |
| 25 | Fixed-distance walk tests at comfortable and fast speed: Potential tools for the functional assessment of coronary patients?. <i>Annals of Physical and Rehabilitation Medicine</i> , <b>2017</b> , 60, 13-19  | 3.8 | 7         |
| 24 | Microcirculatory Assessment of Arterial Below-Knee Stumps: Near-Infrared Spectroscopy Versus Transcutaneous Oxygen Tension-A Preliminary Study in Prosthesis Users. <i>Archives of Physical Medicine and Rehabilitation</i> , <b>2017</b> , 98, 1187-1194  | 2.8 | 4         |
| 23 | A study of the 200-metre fast walk test as a possible new assessment tool to predict maximal heart rate and define target heart rate for exercise training of coronary heart disease patients. <i>Clinical Rehabilitation</i> , <b>2015</b> , 29, 175-83   | 3.3 | 4         |
| 22 | The Log book for the secondary prevention of coronary artery disease: A pilot study. <i>Presse Medicale</i> , <b>2015</b> , 44, e301-9   | 2.2 |           |
| 21 | Transcultural validation of the SIGAM mobility grades in French: The SIGAM-Fr. <i>Annals of Physical and Rehabilitation Medicine</i> , <b>2015</b> , 58, 161-6   | 3.8 | 3         |
| 20 | Cross-talk correction method for knee kinematics in gait analysis using principal component analysis (PCA): a new proposal. <i>PLoS ONE</i> , <b>2014</b> , 9, e102098   | 3.7 | 25        |
| 19 | Is it possible to individualize intensity of eccentric cycling exercise from perceived exertion on concentric test?. <i>Archives of Physical Medicine and Rehabilitation</i> , <b>2013</b> , 94, 1621-1627.e1  | 2.8 | 12        |
| 18 | Selecting a test for the clinical assessment of balance and walking capacity at the definitive fitting state after unilateral amputation: a comparative study. <i>Prosthetics and Orthotics International</i> , <b>2012</b> , 36, 415-22   | 1.5 | 36        |
| 17 | Nurse-led educative consultation setting personalized tertiary prevention goals after cardiovascular rehabilitation: evaluation of patient satisfaction and long-term effects. <i>Rehabilitation Nursing</i> , <b>2012</b> , 37, 105-13  | 1.3 | 7         |
| 16 | Poststroke disposition and associated factors in a population-based study: the Dijon Stroke Registry. <i>Stroke</i> , <b>2012</b> , 43, 2071-7   | 6.7 | 45        |
| 15 | Determining the minimal clinically important difference for the six-minute walk test and the 200-meter fast-walk test during cardiac rehabilitation program in coronary artery disease patients after acute coronary syndrome. <i>Archives of Physical Medicine and Rehabilitation</i> , <b>2011</b> , 92, 611-9 | 2.8 | 125       |

|    |  |     |    |
|----|--|-----|----|
| 14 | How do scoliotic women shrink throughout life?. <i>Spine</i> , <b>2009</b> , 34, 598-602   | 3.3 | 2  |
| 13 | Low-frequency electric muscle stimulation combined with physical therapy after total hip arthroplasty for hip osteoarthritis in elderly patients: a randomized controlled trial. <i>Archives of Physical Medicine and Rehabilitation</i> , <b>2008</b> , 89, 2265-73 | 2.8 | 26 |
| 12 | Comparative analysis of oxygen uptake in elderly subjects performing two walk tests: the six-minute walk test and the 200-m fast walk test. <i>Clinical Rehabilitation</i> , <b>2008</b> , 22, 162-8   | 3.3 | 26 |
| 11 | Do low-frequency electrical myostimulation and aerobic training similarly improve performance in chronic heart failure patients with different exercise capacities?. <i>Acta Dermato-Venereologica</i> , <b>2008</b> , 40, 219-24                                    | 2.2 | 23 |
| 10 | Analysis of low back pain in adults with scoliosis. <i>Spine</i> , <b>2008</b> , 33, 402-5   | 3.3 | 17 |
| 9  | Neuromuscular adaptations to low-frequency stimulation training in a patient with chronic heart failure. <i>American Journal of Physical Medicine and Rehabilitation</i> , <b>2008</b> , 87, 502-9   | 2.6 | 9  |
| 8  | Does proprioception contribute to the sense of verticality?. <i>Experimental Brain Research</i> , <b>2008</b> , 185, 545-53  | 5.3 | 83 |
| 7  | Effects of a one-year exercise training program in adults over 70 years old: a study with a control group. <i>Aging Clinical and Experimental Research</i> , <b>2007</b> , 19, 310-5   | 4.8 | 49 |
| 6  | Comparison of low-frequency electrical myostimulation and conventional aerobic exercise training in patients with chronic heart failure. <i>European Journal of Cardiovascular Prevention and Rehabilitation</i> , <b>2005</b> , 12, 226-33                          |     | 30 |
| 5  | Comparison of low-frequency electrical myostimulation and conventional aerobic exercise training in patients with chronic heart failure. <i>European Journal of Cardiovascular Prevention and Rehabilitation</i> , <b>2005</b> , 12, 226-233                         |     | 50 |
| 4  | Healing of open stump wounds after vascular below-knee amputation: plaster cast socket with silicone sleeve versus elastic compression. <i>Archives of Physical Medicine and Rehabilitation</i> , <b>1999</b> , 80, 1327-30  | 2.8 | 33 |
| 3  | Nuclear magnetic resonance evidence of different muscular adaptations after resistance training. <i>Archives of Physical Medicine and Rehabilitation</i> , <b>1998</b> , 79, 1391-8  | 2.8 | 15 |
| 2  | Muscle metabolism changes with training in the nonamputated limb after vascular amputation: interest of phosphorus 31 NMR spectroscopy. <i>Archives of Physical Medicine and Rehabilitation</i> , <b>1997</b> , 78, 867-71   | 2.8 | 9  |
| 1  | Bioenergetic comparison of a new energy-storing foot and SACH foot in traumatic below-knee vascular amputations. <i>Archives of Physical Medicine and Rehabilitation</i> , <b>1995</b> , 76, 39-44   | 2.8 | 59 |