

Carles Justicia

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/7645453/publications.pdf>

Version: 2024-02-01

48
papers

3,124
citations

168829

31
h-index

286692

43
g-index

48
all docs

48
docs citations

48
times ranked

4660
citing authors

#	ARTICLE	IF	CITATIONS
1	Dendritic Cells and Microglia Have Non-redundant Functions in the Inflamed Brain with Protective Effects of Type 1 cDCs. <i>Cell Reports</i> , 2020, 33, 108291.	2.9	39
2	CCR2 deficiency in monocytes impairs angiogenesis and functional recovery after ischemic stroke in mice. <i>Journal of Cerebral Blood Flow and Metabolism</i> , 2020, 40, S98-S116.	2.4	57
3	Role of the S1P pathway and inhibition by fingolimod in preventing hemorrhagic transformation after stroke. <i>Scientific Reports</i> , 2019, 9, 8309.	1.6	39
4	Location of Neutrophils in Different Compartments of the Damaged Mouse Brain After Severe Ischemia/Reperfusion. <i>Stroke</i> , 2019, 50, 1548-1557.	1.0	61
5	Microglial cell loss after ischemic stroke favors brain neutrophil accumulation. <i>Acta Neuropathologica</i> , 2019, 137, 321-341.	3.9	177
6	High-density speckle contrast optical tomography of cerebral blood flow response to functional stimuli in the rodent brain. <i>Neurophotonics</i> , 2019, 6, 1.	1.7	14
7	CNS-border associated macrophages respond to acute ischemic stroke attracting granulocytes and promoting vascular leakage. <i>Acta Neuropathologica Communications</i> , 2018, 6, 76.	2.4	78
8	T Cells Prevent Hemorrhagic Transformation in Ischemic Stroke by P-Selectin Binding. <i>Arteriosclerosis, Thrombosis, and Vascular Biology</i> , 2018, 38, 1761-1771.	1.1	38
9	High-density speckle contrast optical tomography (SCOT) for three dimensional tomographic imaging of the small animal brain. <i>NeuroImage</i> , 2017, 153, 283-292.	2.1	21
10	Uric Acid Is Protective After Cerebral Ischemia/Reperfusion in Hyperglycemic Mice. <i>Translational Stroke Research</i> , 2017, 8, 294-305.	2.3	45
11	A CNS-permeable Hsp90 inhibitor rescues synaptic dysfunction and memory loss in APP-overexpressing Alzheimer's mouse model via an HSF1-mediated mechanism. <i>Molecular Psychiatry</i> , 2017, 22, 990-1001.	4.1	40
12	Longitudinal, transcranial measurement of functional activation in the rat brain by diffuse correlation spectroscopy. <i>Neurophotonics</i> , 2017, 4, 1.	1.7	3
13	Immature monocytes recruited to the ischemic mouse brain differentiate into macrophages with features of alternative activation. <i>Brain, Behavior, and Immunity</i> , 2016, 53, 18-33.	2.0	111
14	A new method utilizing novel single-photon avalanche diode arrays for multi-exposure laser speckle flowmetry. , 2016, , .		0
15	Latest developments in speckle contrast optical tomography (SCOT) for deep tissue blood flow imaging. , 2016, , .		0
16	Neutrophil recruitment to the brain in mouse and human ischemic stroke. <i>Acta Neuropathologica</i> , 2015, 129, 239-257.	3.9	307
17	High-speed multi-exposure laser speckle contrast imaging with a single-photon counting camera. <i>Biomedical Optics Express</i> , 2015, 6, 2865.	1.5	46
18	Mannose-Binding Lectin Promotes Local Microvascular Thrombosis After Transient Brain Ischemia in Mice. <i>Stroke</i> , 2014, 45, 1453-1459.	1.0	45

#	ARTICLE	IF	CITATIONS
19	Hypoxia and P1 receptor activation regulate the high-affinity concentrative adenosine transporter CNT2 in differentiated neuronal PC12 cells. <i>Biochemical Journal</i> , 2013, 454, 437-445.	1.7	26
20	In vivo imaging of induction of heat-shock protein-70 gene expression with fluorescence reflectance imaging and intravital confocal microscopy following brain ischaemia in reporter mice. <i>European Journal of Nuclear Medicine and Molecular Imaging</i> , 2013, 40, 426-438.	3.3	15
21	Improving image quality in small animal diffusion tensor imaging at 7T. , 2012, , .		0
22	Stem Cell Mediation of Functional Recovery after Stroke in the Rat. <i>PLoS ONE</i> , 2010, 5, e12779.	1.1	69
23	Genetically-Defined Deficiency of Mannose-Binding Lectin Is Associated with Protection after Experimental Stroke in Mice and Outcome in Human Stroke. <i>PLoS ONE</i> , 2010, 5, e8433.	1.1	128
24	Reproducible imaging of rat corticothalamic pathway by longitudinal manganese-enhanced MRI (L-MEMRI). <i>NeuroImage</i> , 2008, 41, 668-674.	2.1	25
25	MRI Detection of Secondary Damage After Stroke. <i>Stroke</i> , 2008, 39, 1541-1547.	1.0	65
26	Transient middle cerebral artery occlusion causes different structural, mechanical, and myogenic alterations in normotensive and hypertensive rats. <i>American Journal of Physiology - Heart and Circulatory Physiology</i> , 2007, 293, H628-H635.	1.5	34
27	Cell tracking using magnetic resonance imaging. <i>Journal of Physiology</i> , 2007, 584, 25-30.	1.3	80
28	Anti-VCAM-1 Antibodies did not Protect against Ischemic Damage Either in Rats Or in Mice. <i>Journal of Cerebral Blood Flow and Metabolism</i> , 2006, 26, 421-432.	2.4	104
29	Modest MRI Signal Intensity Changes Precede Delayed Cortical Necrosis After Transient Focal Ischemia in the Rat. <i>Stroke</i> , 2006, 37, 1525-1532.	1.0	31
30	Steady plasma concentration of unfractionated heparin reduces infarct volume and prevents inflammatory damage after transient focal cerebral ischemia in the rat. <i>Journal of Neuroscience Research</i> , 2004, 77, 565-572.	1.3	38
31	Caspase-dependent and caspase-independent signalling of apoptosis in the penumbra following middle cerebral artery occlusion in the adult rat. <i>Neuropathology and Applied Neurobiology</i> , 2003, 29, 472-481.	1.8	94
32	Neutrophil Infiltration Increases Matrix Metalloproteinase-9 in the Ischemic Brain after Occlusion/Reperfusion of the Middle Cerebral Artery in Rats. <i>Journal of Cerebral Blood Flow and Metabolism</i> , 2003, 23, 1430-1440.	2.4	221
33	Activation of ERK and Akt Signaling in Focal Cerebral Ischemia: Modulation by TGF- β and Involvement of NMDA Receptor. <i>Neurobiology of Disease</i> , 2002, 11, 443-456.	2.1	40
34	Certain Forms of Matrix Metalloproteinase-9 Accumulate in the Extracellular Space after Microdialysis Probe Implantation and Middle Cerebral Artery Occlusion/Reperfusion. <i>Journal of Cerebral Blood Flow and Metabolism</i> , 2002, 22, 918-925.	2.4	24
35	Expression and Activation of Matrix Metalloproteinase-2 and -9 in Rat Brain after Transient Focal Cerebral Ischemia. <i>Neurobiology of Disease</i> , 2001, 8, 834-846.	2.1	215
36	Focal cerebral ischemia causes two temporal waves of Akt activation. <i>NeuroReport</i> , 2001, 12, 3381-3384.	0.6	34

#	ARTICLE	IF	CITATIONS
37	Temporospatial expression of HSP72 and c-JUN, and DNA fragmentation in goat hippocampus after global cerebral ischemia. <i>Hippocampus</i> , 2001, 11, 146-156.	0.9	9
38	Administration of Transforming Growth Factor- β Reduces Infarct Volume after Transient Focal Cerebral Ischemia in the Rat. <i>Journal of Cerebral Blood Flow and Metabolism</i> , 2001, 21, 1097-1104.	2.4	61
39	Activation of the JAK/STAT pathway following transient focal cerebral ischemia: Signaling through Jak1 and Stat3 in astrocytes. , 2000, 30, 253-270.		181
40	Estimation of Gelatinase Content in Rat Brain: Effect of Focal Ischemia. <i>Biochemical and Biophysical Research Communications</i> , 2000, 278, 803-807.	1.0	36
41	Transforming Growth Factor- β Acting at the Epidermal Growth Factor Receptor Reduces Infarct Volume after Permanent Middle Cerebral Artery Occlusion in Rats. <i>Journal of Cerebral Blood Flow and Metabolism</i> , 1999, 19, 128-132.	2.4	46
42	Induction of cyclooxygenase-2 in the rat brain after a mild episode of focal ischemia without tissue inflammation or neural cell damage. <i>Neuroscience Letters</i> , 1999, 275, 141-144.	1.0	38
43	Activation of nuclear factor- κ B in the rat brain after transient focal ischemia. <i>Molecular Brain Research</i> , 1999, 65, 61-69.	2.5	116
44	Epidermal growth factor receptor in proliferating reactive glia following transient focal ischemia in the rat brain. , 1998, 23, 120-129.		87
45	Stat1 in developing and adult rat brain. Induction after transient focal ischemia. <i>NeuroReport</i> , 1997, 8, 1359-1362.	0.6	22
46	Striatal Infarction in the Rat Causes a Transient Reduction of Tyrosine Hydroxylase Immunoreactivity in the Ipsilateral Substantia Nigra. <i>Neurobiology of Disease</i> , 1997, 4, 376-385.	2.1	29
47	Stat3 Is Present in the Developing and Adult Rat Cerebellum and Participates in the Formation of Transcription Complexes Binding DNA at the γ -Inducible Element. <i>Journal of Neurochemistry</i> , 1997, 68, 1345-1351.	2.1	35
48	Induction of Stat3, a Signal Transducer and Transcription Factor, in Reactive Microglia following Transient Focal Cerebral Ischaemia. <i>European Journal of Neuroscience</i> , 1996, 8, 2612-2618.	1.2	100