

# Warwick D Ngan Kee

## List of Publications by Year in Descending Order

**Source:** <https://exaly.com/author-pdf/7660274/warwick-d-ngan-kee-publications-by-year.pdf>

**Version:** 2024-04-27

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

56  
papers

2,815  
citations

26  
h-index

53  
g-index

59  
ext. papers

3,291  
ext. citations

4.5  
avg, IF

5.31  
L-index

#	Paper	IF	Citations
56	Intravenous lidocaine attenuates response to cervical dilation for hysteroscopy: a randomised controlled trial. <i>British Journal of Anaesthesia</i> , <b>2021</b> , 127, e166-e168	5.4	0
55	Dose-Response Study of 4 Weight-Based Phenylephrine Infusion Regimens for Preventing Hypotension During Cesarean Delivery Under Combined Spinal-Epidural Anesthesia. <i>Anesthesia and Analgesia</i> , <b>2020</b> , 130, 187-193	3.9	8
54	Norepinephrine or phenylephrine during spinal anaesthesia for Caesarean delivery: a randomised double-blind pragmatic non-inferiority study of neonatal outcome. <i>British Journal of Anaesthesia</i> , <b>2020</b> , 125, 588-595	5.4	10
53	Performance of a closed-loop feedback computer-controlled infusion system for maintaining blood pressure during spinal anaesthesia for caesarean section: a randomized controlled comparison of norepinephrine versus phenylephrine. <i>Journal of Clinical Monitoring and Computing</i> , <b>2017</b> , 31, 617-623	2	31
52	Norepinephrine Intermittent Intravenous Boluses to Prevent Hypotension During Spinal Anesthesia for Cesarean Delivery: A Sequential Allocation Dose-Finding Study. <i>Anesthesia and Analgesia</i> , <b>2017</b> , 125, 212-218	3.9	54
51	Closed-Loop Feedback Computer-Controlled Phenylephrine for Maintenance of Blood Pressure During Spinal Anesthesia for Cesarean Delivery: A Randomized Trial Comparing Automated Boluses Versus Infusion. <i>Anesthesia and Analgesia</i> , <b>2017</b> , 125, 117-123	3.9	21
50	The use of vasopressors during spinal anaesthesia for caesarean section. <i>Current Opinion in Anaesthesiology</i> , <b>2017</b> , 30, 319-325	2.9	36
49	A Random-allocation Graded Dose-Response Study of Norepinephrine and Phenylephrine for Treating Hypotension during Spinal Anesthesia for Cesarean Delivery. <i>Anesthesiology</i> , <b>2017</b> , 127, 934-941	4.3	60
48	Randomized double-blinded comparison of norepinephrine and phenylephrine for maintenance of blood pressure during spinal anesthesia for cesarean delivery. <i>Anesthesiology</i> , <b>2015</b> , 122, 736-45	4.3	138
47	Synergistic interaction between fentanyl and bupivacaine given intrathecally for labor analgesia. <i>Anesthesiology</i> , <b>2014</b> , 120, 1126-36	4.3	36
46	The effect of maternal and fetal $\beta$ -adrenoceptor and nitric oxide synthase genotype on vasopressor requirement and fetal acid-base status during spinal anesthesia for cesarean delivery. <i>Anesthesia and Analgesia</i> , <b>2011</b> , 112, 1432-7	3.9	23
45	Pulmonary Disease in Pregnancy <b>2010</b> , 1-47		2
44	Prevention of maternal hypotension after regional anaesthesia for caesarean section. <i>Current Opinion in Anaesthesiology</i> , <b>2010</b> , 23, 304-9	2.9	90
43	Determination and comparison of graded dose-response curves for epidural bupivacaine and ropivacaine for analgesia in laboring nulliparous women. <i>Anesthesiology</i> , <b>2010</b> , 113, 445-53	4.3	23
42	Ephedrine and Phenylephrine Use during Cesarean Delivery. <i>Anesthesiology</i> , <b>2010</b> , 112, 1290-1294	4.3	1
41	The median effective dose of bupivacaine, levobupivacaine, and ropivacaine after intrathecal injection in lower limb surgery. <i>Anesthesia and Analgesia</i> , <b>2009</b> , 109, 1331-4	3.9	43
40	Placental transfer and fetal metabolic effects of phenylephrine and ephedrine during spinal anesthesia for cesarean delivery. <i>Anesthesiology</i> , <b>2009</b> , 111, 506-12	4.3	189

39	Low-dose spinal anesthesia with low-dose phenylephrine infusions for cesarean delivery: better but not necessarily best. <i>Anesthesiology</i> , <b>2009</b> , 111, 210-1; author reply 213-6	4.3	1
38	A randomized double-blinded comparison of phenylephrine and ephedrine infusion combinations to maintain blood pressure during spinal anesthesia for cesarean delivery: the effects on fetal acid-base status and hemodynamic control. <i>Anesthesia and Analgesia</i> , <b>2008</b> , 107, 1295-302	3.9	148
37	A prospective comparison of vasopressor requirement and hemodynamic changes during spinal anesthesia for cesarean delivery in patients with multiple gestation versus singleton pregnancy. <i>Anesthesia and Analgesia</i> , <b>2007</b> , 104, 407-11	3.9	11
36	Spinal ropivacaine for lower limb surgery: a dose response study. <i>Anesthesia and Analgesia</i> , <b>2007</b> , 105, 520-3	3.9	18
35	A systematic review (meta-analysis) of the accuracy of the Mallampati tests to predict the difficult airway. <i>Anesthesia and Analgesia</i> , <b>2006</b> , 102, 1867-78	3.9	191
34	HYPOTENSION DURING SPINAL ANAESTHESIA FOR CAESAREAN SECTION: IMPLICATIONS, DETECTION PREVENTION AND TREATMENT. <i>Fetal and Maternal Medicine Review</i> , <b>2006</b> , 17, 157-183		15
33	Maternal and neonatal effects of remifentanyl at induction of general anesthesia for cesarean delivery: a randomized, double-blind, controlled trial. <i>Anesthesiology</i> , <b>2006</b> , 104, 14-20	4.3	127
32	Vasopressors in obstetrics: what should we be using?. <i>Current Opinion in Anaesthesiology</i> , <b>2006</b> , 19, 238-43		86
31	Prevention of hypotension during spinal anesthesia for cesarean delivery: an effective technique using combination phenylephrine infusion and crystalloid cohydration. <i>Anesthesiology</i> , <b>2005</b> , 103, 744-50	4.3	237
30	A double blinded randomised placebo-controlled study of intramuscular pethidine for pain relief in the first stage of labour. <i>BJOG: an International Journal of Obstetrics and Gynaecology</i> , <b>2004</b> , 111, 648-55	3.7	89
29	Epidural infusions of ropivacaine and bupivacaine for labor analgesia: a randomized, double-blind study of obstetric outcome. <i>Anesthesia and Analgesia</i> , <b>2004</b> , 98, 1145-1152	3.9	29
28	Prophylactic phenylephrine infusion for preventing hypotension during spinal anesthesia for cesarean delivery. <i>Anesthesia and Analgesia</i> , <b>2004</b> , 98, 815-21, table of contents	3.9	110
27	A dose-response meta-analysis of prophylactic intravenous ephedrine for the prevention of hypotension during spinal anesthesia for elective cesarean delivery. <i>Anesthesia and Analgesia</i> , <b>2004</b> , 98, 483-490	3.9	58
26	Intrathecal meperidine and shivering in obstetric anesthesia. <i>Anesthesia and Analgesia</i> , <b>2004</b> , 99, 1272-1273	3.7	4
25	Obstetric neuraxial anaesthesia: which vasopressor should we be using?. <i>International Journal of Obstetric Anesthesia</i> , <b>2003</b> , 12, 55-6; author reply 56-7	2.1	5
24	Prophylactic ephedrine prevents hypotension during spinal anesthesia for Cesarean delivery but does not improve neonatal outcome: a quantitative systematic review. <i>Canadian Journal of Anaesthesia</i> , <b>2002</b> , 49, 588-99	3	62
23	Spinal ropivacaine for cesarean delivery: a comparison of hyperbaric and plain solutions. <i>Anesthesia and Analgesia</i> , <b>2002</b> , 94, 680-5; table of contents	3.9	76
22	The effect of the addition of epinephrine on early systemic absorption of epidural ropivacaine in humans. <i>Anesthesia and Analgesia</i> , <b>2002</b> , 95, 1402-7, table of contents	3.9	30

21	A quantitative, systematic review of randomized controlled trials of ephedrine versus phenylephrine for the management of hypotension during spinal anesthesia for cesarean delivery. <i>Anesthesia and Analgesia</i> , <b>2002</b> , 94, 920-6, table of contents	3.9	305
20	Epidural infusions for labor analgesia: a comparison of 0.2% ropivacaine, 0.1% ropivacaine, and 0.1% ropivacaine with fentanyl. <i>Regional Anesthesia and Pain Medicine</i> , <b>2002</b> , 27, 31-6	3.4	22
19	Vertebral osteomyelitis and psoas abscess occurring after obstetric epidural anesthesia. <i>Regional Anesthesia and Pain Medicine</i> , <b>2002</b> , 27, 220-4	3.4	10
18	The limitations of ropivacaine with epinephrine as an epidural test dose in parturients. <i>Anesthesia and Analgesia</i> , <b>2001</b> , 92, 1529-31	3.9	9
17	Metaraminol infusion for maintenance of arterial blood pressure during spinal anesthesia for cesarean delivery: the effect of a crystalloid bolus. <i>Anesthesia and Analgesia</i> , <b>2001</b> , 93, 703-8	3.9	31
16	Anesthetic management of renal cell carcinoma with inferior vena caval extension. <i>Journal of Clinical Anesthesia</i> , <b>2001</b> , 13, 585-7	1.9	9
15	A dose-response study of prophylactic intravenous ephedrine for the prevention of hypotension during spinal anesthesia for cesarean delivery. <i>Anesthesia and Analgesia</i> , <b>2000</b> , 90, 1390-5	3.9	138
14	Idiopathic dilated cardiomyopathy presenting in pregnancy. <i>Canadian Journal of Anaesthesia</i> , <b>1999</b> , 46, 1146-9	3	19
13	Postdural-puncture headache. <i>Lancet, The</i> , <b>1999</b> , 354, 680	4.0	0
12	Perioperative documentation: are we doing enough?. <i>Australian and New Zealand Journal of Obstetrics and Gynaecology</i> , <b>1998</b> , 38, 166-9	1.7	3
11	Metabolic and inflammatory responses after laparoscopic and abdominal hysterectomy. <i>American Journal of Obstetrics and Gynecology</i> , <b>1998</b> , 179, 1-5	6.4	61
10	Postoperative epidural opioid analgesia: what are the choices?. <i>Anesthesia and Analgesia</i> , <b>1997</b> , 84, 1390-3	3.9	1
9	Epidural meperidine after cesarean section: the effect of diluent volume. <i>Anesthesia and Analgesia</i> , <b>1997</b> , 85, 380-4	3.9	5
8	Postoperative analgesic requirement after cesarean section: a comparison of anesthetic induction with ketamine or thiopental. <i>Anesthesia and Analgesia</i> , <b>1997</b> , 85, 1294-8	3.9	10
7	Stroke following central venous cannulation. <i>Lancet, The</i> , <b>1997</b> , 349, 921	4.0	64
6	How accurate are references in the Australian and New Zealand Journal of Surgery?. <i>ANZ Journal of Surgery</i> , <b>1997</b> , 67, 417-9	1	13
5	A survey of factors influencing patients' choice of anaesthesia for caesarean section. <i>Australian and New Zealand Journal of Obstetrics and Gynaecology</i> , <b>1997</b> , 37, 300-3	1.7	9
4	Patient-Controlled Epidural Analgesia After Caesarean Section Using a Disposable Device. <i>Australian and New Zealand Journal of Obstetrics and Gynaecology</i> , <b>1997</b> , 37, 304-307	1.7	5

- |   |  |   |    |
|---|--|---|----|
| 3 | Patient-controlled epidural analgesia after caesarean section using meperidine. <i>Canadian Journal of Anaesthesia</i> , <b>1997</b> , 44, 702-6 | 3 | 10 |
| 2 | Respiratory depression associated with patient-controlled analgesia. <i>Canadian Journal of Anaesthesia</i> , <b>1995</b> , 42, 953-4            | 3 | 1  |
| 1 | Pharmacology, pharmacokinetics, and management of the patient after overdose150-159  |   |    |