
Helen Sjöland

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/7666479/publications.pdf Version: 2024-02-01



HELEN SIÃTIAND

#	Article	IF	CITATIONS
1	Prognostic differences in longâ€standing vs. recentâ€onset dilated cardiomyopathy. ESC Heart Failure, 2022, , .	3.1	2
2	BMI, sex and outcomes in hospitalised patients in western Sweden during the COVID-19 pandemic. Scientific Reports, 2022, 12, 4918.	3.3	5
3	Temporal trends in outcome and patient characteristics in dilated cardiomyopathy, data from the Swedish Heart Failure Registry 2003–2015. BMC Cardiovascular Disorders, 2021, 21, 307.	1.7	7
4	Prognostic impact over time of ischaemic heart disease vs. nonâ€ i schaemic heart disease in heart failure. ESC Heart Failure, 2020, 7, 265-274.	3.1	16
5	A negative Tâ€wave in electrocardiogram at 50 years predicted lifetime mortality in a random populationâ€based cohort. Clinical Cardiology, 2020, 43, 1279-1285.	1.8	5
6	Revealed by degrees: Patients' experience of receiving information after inâ€hospital cardiac arrest. Journal of Clinical Nursing, 2019, 28, 1517-1527.	3.0	5
7	Trends in Gender Differences in Cardiac Care and Outcome After Acute Myocardial Infarction in Western Sweden: A Report From the Swedish Web System for Enhancement of Evidenceâ€Based Care in Heart Disease Evaluated According to Recommended Therapies (SWEDEHEART). Journal of the American Heart Association. 2015. 4.	3.7	79
8	Lack of Very Strong Association between Pre-Treatment Fibrinogen and PAI-1 with Long-Term Mortality after Coronary Bypass Surgery. Cardiology, 2007, 108, 82-89.	1.4	7
9	Relief of symptoms and improvement of healthâ€related quality of life five years after coronary artery bypass graft in women and men. Clinical Cardiology, 2001, 24, 385-392.	1.8	32
10	Impact of a History of Hypertension on Symptoms and Quality of Life Prior to and at Five Years after Coronary Artery Bypass Grafting. Blood Pressure, 2000, 9, 52-63.	1.5	3
11	Mortality, risk indicators for death, and mode of death in younger and elderly patients during five years after coronary artery bypass graft. Clinical Cardiology, 2000, 23, 421-426.	1.8	4