

Gunther Hartmann

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/7670135/gunther-hartmann-publications-by-year.pdf>

Version: 2024-04-28

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

195
papers

25,609
citations

70
h-index

159
g-index

215
ext. papers

28,510
ext. citations

10.7
avg, IF

6.53
L-index

#	Paper	IF	Citations
195	Correlation between a quantitative anti-SARS-CoV-2 IgG ELISA and neutralization activity. <i>Journal of Medical Virology</i> , 2022 , 94, 388-392	19.7	16
194	Deficiency in coatamer complex I causes aberrant activation of STING signalling.. <i>Nature Communications</i> , 2022 , 13, 2321	17.4	4
193	Interferon-driven brain phenotype in a mouse model of RNaseT2 deficient leukoencephalopathy. <i>Nature Communications</i> , 2021 , 12, 6530	17.4	1
192	Recessive NLRC4-Autoinflammatory Disease Reveals an Ulcerative Colitis Locus. <i>Journal of Clinical Immunology</i> , 2021 , 1	5.7	2
191	BIOM-24. PROTEIN SURFACE SIGNATURE ON SERUM EXTRACELLULAR VESICLES FOR NON-INVASIVE DETECTION OF TUMOR PROGRESSION IN GLIOBLASTOMA PATIENTS. <i>Neuro-Oncology</i> , 2021 , 23, vi15-vi16	1	
190	X-chromosomale TLR7-Expression bei Frauen und Pr disposition zu Lupus-assoziiierter Autoimmunit. <i>Trillium Immunologie</i> , 2021 , 48-53	0	
189	Memory B cells targeting SARS-CoV-2 spike protein and their dependence on CD4 T cell help. <i>Cell Reports</i> , 2021 , 35, 109320	10.6	17
188	Monocyte-dependent co-stimulation of cytokine induction in human T cells by TLR8 RNA ligands. <i>Scientific Reports</i> , 2021 , 11, 15231	4.9	2
187	Detectable SARS-CoV-2 RNAemia in Critically Ill Patients, but Not in Mild and Asymptomatic Infections. <i>Transfusion Medicine and Hemotherapy</i> , 2021 , 48, 154-160	4.2	0
186	Extracellular Vesicle Separation Techniques Impact Results from Human Blood Samples: Considerations for Diagnostic Applications. <i>International Journal of Molecular Sciences</i> , 2021 , 22,	6.3	1
185	Malaria parasites both repress host CXCL10 and use it as a cue for growth acceleration. <i>Nature Communications</i> , 2021 , 12, 4851	17.4	2
184	MAPK-pathway inhibition mediates inflammatory reprogramming and sensitizes tumors to targeted activation of innate immunity sensor RIG-I. <i>Nature Communications</i> , 2021 , 12, 5505	17.4	2
183	Animal models of SARS-CoV-2 and COVID-19 for the development of prophylactic and therapeutic interventions. <i>Pharmacology & Therapeutics</i> , 2021 , 228, 107931	13.9	7
182	The coffee ingredients caffeic acid and caffeic acid phenylethyl ester protect against irinotecan-induced leukopenia and oxidative stress response. <i>British Journal of Pharmacology</i> , 2020 , 177, 4193-4208	8.6	6
181	BIOM-40. ANALYSIS OF SERUM MIRNA IN GLIOBLASTOMA PATIENTS: TARGETED ENRICHMENT OF EXTRACELLULAR VESICLES ENHANCES SPECIFICITY FOR PROGNOSTIC SIGNATURE. <i>Neuro-Oncology</i> , 2020 , 22, ii10-ii10	1	
180	Targeting the innate immunoreceptor RIG-I overcomes melanoma-intrinsic resistance to T cell immunotherapy. <i>Journal of Clinical Investigation</i> , 2020 , 130, 4266-4281	15.9	15
179	Infektionsletalit von SARS-CoV-2: Eine immunologische Fragestellung. <i>Trillium Immunologie</i> , 2020 , 4, 224-230	0	

178	High RIG-I expression in ovarian cancer associates with an immune-escape signature and poor clinical outcome. <i>International Journal of Cancer</i> , 2020 , 146, 2007-2018	7.5	21
177	Analysis of Serum miRNA in Glioblastoma Patients: CD44-Based Enrichment of Extracellular Vesicles Enhances Specificity for the Prognostic Signature. <i>International Journal of Molecular Sciences</i> , 2020 , 21,	6.3	7
176	Immune Sensing Mechanisms that Discriminate Self from Altered Self and Foreign Nucleic Acids. <i>Immunity</i> , 2020 , 53, 54-77	32.3	48
175	Infection fatality rate of SARS-CoV2 in a super-spreading event in Germany. <i>Nature Communications</i> , 2020 , 11, 5829	17.4	111
174	Absence of cGAS-mediated type I IFN responses in HIV-1-infected T cells. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2020 , 117, 19475-19486	11.5	9
173	A continuous responder algorithm to optimize clinical management of small-cell lung cancer with progastrin-releasing peptide as a simple blood test. <i>Tumor Biology</i> , 2020 , 42, 1010428320958603	2.9	2
172	U-DCS: characterization of the first permanent human dendritic sarcoma cell line. <i>Scientific Reports</i> , 2020 , 10, 21221	4.9	1
171	Immune Sensing of Synthetic, Bacterial, and Protozoan RNA by Toll-like Receptor 8 Requires Coordinated Processing by RNase T2 and RNase 2. <i>Immunity</i> , 2020 , 52, 591-605.e6	32.3	34
170	Human TLR8 Senses RNA From -Infected Red Blood Cells Which Is Uniquely Required for the IFN- γ Response in NK Cells. <i>Frontiers in Immunology</i> , 2019 , 10, 371	8.4	15
169	The PNPLA3 I148M variant promotes lipid-induced hepatocyte secretion of CXC chemokines establishing a tumorigenic milieu. <i>Journal of Molecular Medicine</i> , 2019 , 97, 1589-1600	5.5	3
168	Phenprocoumon Dose Requirements, Dose Stability and Time in Therapeutic Range in Elderly Patients With and Polymorphisms. <i>Frontiers in Pharmacology</i> , 2019 , 10, 1620	5.6	1
167	Targeted Nanoparticle Delivery of Bifunctional RIG-I Agonists to Pancreatic Cancer. <i>Molecular Therapy</i> , 2019 , 27, 491-492	11.7	5
166	Interferon-beta-induced changes in neuroimaging phenotypes of appetitive motivation and reactivity to emotional salience. <i>NeuroImage: Clinical</i> , 2019 , 24, 102020	5.3	1
165	Direct RIG-I activation in human NK cells induces TRAIL-dependent cytotoxicity toward autologous melanoma cells. <i>International Journal of Cancer</i> , 2019 , 144, 1645-1656	7.5	14
164	Structural Alterations of MET Trigger Response to MET Kinase Inhibition in Lung Adenocarcinoma Patients. <i>Clinical Cancer Research</i> , 2018 , 24, 1337-1343	12.9	44
163	Improved sensitivity for detection of breast cancer by combination of miR-34a and tumor markers CA 15-3 or CEA. <i>Oncotarget</i> , 2018 , 9, 22523-22536	3.3	28
162	ATG16L1 orchestrates interleukin-22 signaling in the intestinal epithelium via cGAS-STING. <i>Journal of Experimental Medicine</i> , 2018 , 215, 2868-2886	16.6	83
161	Amplification of N-Myc is associated with a T-cell-poor microenvironment in metastatic neuroblastoma restraining interferon pathway activity and chemokine expression. <i>Oncolmmunology</i> , 2017 , 6, e1320626	7.2	47

160	RIG-I Resists Hypoxia-Induced Immunosuppression and Dedifferentiation. <i>Cancer Immunology Research</i> , 2017 , 5, 455-467	12.5	19
159	Diagnostic relevance of a novel multiplex immunoassay panel in breast cancer. <i>Tumor Biology</i> , 2017 , 39, 1010428317711381	2.9	5
158	Free-Circulating Methylated DNA in Blood for Diagnosis, Staging, Prognosis, and Monitoring of Head and Neck Squamous Cell Carcinoma Patients: An Observational Prospective Cohort Study. <i>Clinical Chemistry</i> , 2017 , 63, 1288-1296	5.5	60
157	Nucleic Acid Immunity. <i>Advances in Immunology</i> , 2017 , 133, 121-169	5.6	118
156	Clinical performance of LOCIbased tumor marker assays for tumor markers CA 15-3, CA 125, CEA, CA 19-9 and AFP in gynecological cancers. <i>Tumor Biology</i> , 2017 , 39, 1010428317730246	2.9	13
155	Analysis of integrated clinical trial protocols in early phases of medicinal product development. <i>European Journal of Clinical Pharmacology</i> , 2017 , 73, 1565-1577	2.8	11
154	RIG-I Activation Protects and Rescues from Lethal Influenza Virus Infection and Bacterial Superinfection. <i>Molecular Therapy</i> , 2017 , 25, 2093-2103	11.7	14
153	Type I interferon-mediated autoinflammation due to DNase II deficiency. <i>Nature Communications</i> , 2017 , 8, 2176	17.4	111
152	Abstract B44: Selective stimulation of RIG-I with a novel synthetic RNA induces strong anti-tumor immunity in mouse tumor models 2017 ,		6
151	Clinical Performance of CEA, CA19-9, CA15-3, CA125 and AFP in Gastrointestinal Cancer Using LOCIbased Assays. <i>Anticancer Research</i> , 2017 , 37, 353-359	2.3	23
150	Diagnostic Performance of a Novel Multiplex Immunoassay in Colorectal Cancer. <i>Anticancer Research</i> , 2017 , 37, 2477-2486	2.3	14
149	RIG-I activation induces the release of extracellular vesicles with antitumor activity. <i>Oncolmmunology</i> , 2016 , 5, e1219827	7.2	35
148	Discriminating self from non-self in nucleic acid sensing. <i>Nature Reviews Immunology</i> , 2016 , 16, 566-80	36.5	253
147	Inflammasome-Dependent Induction of Adaptive NK Cell Memory. <i>Immunity</i> , 2016 , 44, 1406-21	32.3	55
146	Individualized versus standardized risk assessment in patients at high risk for adverse drug reactions (IDrug) - study protocol for a pragmatic randomized controlled trial. <i>BMC Family Practice</i> , 2016 , 17, 49	2.6	8
145	Cutting Edge: The RIG-I Ligand 3pRNA Potently Improves CTL Cross-Priming and Facilitates Antiviral Vaccination. <i>Journal of Immunology</i> , 2016 , 196, 2439-43	5.3	30
144	G-rich DNA-induced stress response blocks type-I-IFN but not CXCL10 secretion in monocytes. <i>Scientific Reports</i> , 2016 , 6, 38405	4.9	2
143	MDA-5 activation by cytoplasmic double-stranded RNA impairs endothelial function and aggravates atherosclerosis. <i>Journal of Cellular and Molecular Medicine</i> , 2016 , 20, 1696-705	5.6	11

142	A Conserved Histidine in the RNA Sensor RIG-I Controls Immune Tolerance to N1-2O-Methylated Self RNA. <i>Immunity</i> , 2015 , 43, 41-51	32.3	154
141	Sequence-specific activation of the DNA sensor cGAS by Y-form DNA structures as found in primary HIV-1 cDNA. <i>Nature Immunology</i> , 2015 , 16, 1025-33	19.1	145
140	Immune- and miRNA-response to recombinant interferon beta-1a: a biomarker evaluation study to guide the development of novel type I interferon- based therapies. <i>BMC Pharmacology & Toxicology</i> , 2015 , 16, 25	2.6	5
139	Where Failure Is Not an Option -Personalized Medicine in Astronauts. <i>PLoS ONE</i> , 2015 , 10, e0140764	3.7	19
138	ATP hydrolysis by the viral RNA sensor RIG-I prevents unintentional recognition of self-RNA. <i>ELife</i> , 2015 , 4,	8.9	63
137	Efficient solid-phase synthesis of pppRNA by using product-specific labeling. <i>Angewandte Chemie - International Edition</i> , 2014 , 53, 4694-8	16.4	22
136	Yeast virus-derived stimulator of the innate immune system augments the efficacy of virus vector-based immunotherapy. <i>Journal of Virology</i> , 2014 , 88, 5242-55	6.6	11
135	Calponin-h2: a potential serum marker for the early detection of human breast cancer?. <i>Tumor Biology</i> , 2014 , 35, 11121-7	2.9	5
134	Characterizing the genetic basis of innate immune response in TLR4-activated human monocytes. <i>Nature Communications</i> , 2014 , 5, 5236	17.4	48
133	Therapeutic tissue regeneration by a macrophage colony-stimulating factor Fc conjugate. <i>Molecular Therapy</i> , 2014 , 22, 1577-9	11.7	1
132	Antiviral immunity via RIG-I-mediated recognition of RNA bearing 5Rdiphosphates. <i>Nature</i> , 2014 , 514, 372-375	50.4	359
131	Self-priming determines high type I IFN production by plasmacytoid dendritic cells. <i>European Journal of Immunology</i> , 2014 , 44, 807-818	6.1	47
130	Binding-pocket and lid-region substitutions render human STING sensitive to the species-specific drug DMXAA. <i>Cell Reports</i> , 2014 , 8, 1668-1676	10.6	58
129	AChE and RACK1 promote the anti-inflammatory properties of fluoxetine. <i>Journal of Molecular Neuroscience</i> , 2014 , 53, 306-15	3.3	29
128	Enzymatic synthesis and purification of a defined RIG-I ligand. <i>Methods in Molecular Biology</i> , 2014 , 1169, 15-25	1.4	9
127	VKORC1-dependent pharmacokinetics of intravenous and oral phylloquinone (vitamin K1) mixed micelles formulation. <i>European Journal of Clinical Pharmacology</i> , 2013 , 69, 467-75	2.8	9
126	Structure-function analysis of STING activation by c[G(2R5R)pA(3R5R)p] and targeting by antiviral DMXAA. <i>Cell</i> , 2013 , 154, 748-62	56.2	322
125	Turning tumors into vaccines: co-opting the innate immune system. <i>Immunity</i> , 2013 , 39, 27-37	32.3	72

124	Specific expression of k63-linked ubiquitination of calmodulin-like protein 5 in breast cancer of premenopausal patients. <i>Journal of Cancer Research and Clinical Oncology</i> , 2013 , 139, 2125-32	4.9	15
123	Oxidative damage of DNA confers resistance to cytosolic nuclease TREX1 degradation and potentiates STING-dependent immune sensing. <i>Immunity</i> , 2013 , 39, 482-95	32.3	251
122	Exosomes as nucleic acid nanocarriers. <i>Advanced Drug Delivery Reviews</i> , 2013 , 65, 331-5	18.5	166
121	Cyclic [G(2',5'-BpA(3',5'-Bp)] is the metazoan second messenger produced by DNA-activated cyclic GMP-AMP synthase. <i>Cell</i> , 2013 , 153, 1094-107	56.2	562
120	Therapeutic efficacy of bifunctional siRNA combining TGF- β silencing with RIG-I activation in pancreatic cancer. <i>Cancer Research</i> , 2013 , 73, 1709-20	10.1	103
119	Targeting the cytosolic innate immune receptors RIG-I and MDA5 effectively counteracts cancer cell heterogeneity in glioblastoma. <i>Stem Cells</i> , 2013 , 31, 1064-74	5.8	55
118	RIG-I detects triphosphorylated RNA of <i>Listeria monocytogenes</i> during infection in non-immune cells. <i>PLoS ONE</i> , 2013 , 8, e62872	3.7	62
117	A human in vitro whole blood assay to predict the systemic cytokine response to therapeutic oligonucleotides including siRNA. <i>PLoS ONE</i> , 2013 , 8, e71057	3.7	35
116	Immunohistological analysis of in-transit metastasis in a patient with advanced melanoma treated with combination therapy of cytosine guanine dinucleotide oligodeoxynucleotide, dacarbazine and beta-interferon: a case report. <i>Journal of Dermatology</i> , 2012 , 39, 1035-7	1.6	1
115	Endothelial RIG-I activation impairs endothelial function. <i>Biochemical and Biophysical Research Communications</i> , 2012 , 420, 66-71	3.4	18
114	RIG-I detects infection with live <i>Listeria</i> by sensing secreted bacterial nucleic acids. <i>EMBO Journal</i> , 2012 , 31, 4153-64	13	132
113	Nucleic acid adjuvants: toward an educated vaccine. <i>Advances in Immunology</i> , 2012 , 114, 1-32	5.6	8
112	Stressing hematopoiesis and immunity: an acetylcholinesterase window into nervous and immune system interactions. <i>Frontiers in Molecular Neuroscience</i> , 2012 , 5, 30	6.1	27
111	Influence of acute exposure to high altitude on basal and postprandial plasma levels of gastroenteropancreatic peptides. <i>PLoS ONE</i> , 2012 , 7, e44445	3.7	22
110	Cytosolic RIG-I-like helicases act as negative regulators of sterile inflammation in the CNS. <i>Nature Neuroscience</i> , 2011 , 15, 98-106	25.5	54
109	5'Triphosphorylated small interfering RNAs control replication of hepatitis B virus and induce an interferon response in human liver cells and mice. <i>Gastroenterology</i> , 2011 , 141, 696-706, 706.e1-3	13.3	57
108	Delivery with polycations extends the immunostimulant Ribomunyl into a potent antiviral Toll-like receptor 7/8 agonist. <i>Antiviral Therapy</i> , 2011 , 16, 751-8	1.6	4
107	Stimulation of TLR7 prior to polymicrobial sepsis improves the immune control of the inflammatory response in adult mice. <i>Inflammation Research</i> , 2011 , 60, 271-9	7.2	11

106	Immunogenic cell death of human ovarian cancer cells induced by cytosolic poly(I:C) leads to myeloid cell maturation and activates NK cells. <i>European Journal of Immunology</i> , 2011 , 41, 3028-39	6.1	35
105	Identification of specific nuclear structural protein alterations in human breast cancer. <i>Journal of Cellular Biochemistry</i> , 2011 , 112, 3176-84	4.7	6
104	Activation of endothelial toll-like receptor 3 impairs endothelial function. <i>Circulation Research</i> , 2011 , 108, 1358-66	15.7	87
103	Structural and functional insights into 5Rppp RNA pattern recognition by the innate immune receptor RIG-I. <i>Nature Structural and Molecular Biology</i> , 2010 , 17, 781-7	17.6	196
102	Recognition of RNA virus by RIG-I results in activation of CARD9 and inflammasome signaling for interleukin 1 beta production. <i>Nature Immunology</i> , 2010 , 11, 63-9	19.1	407
101	Immunostimulatory RNA blocks suppression by regulatory T cells. <i>Journal of Immunology</i> , 2010 , 184, 939-46	5.3	47
100	Human plasmacytoid dendritic cells support Th17 cell effector function in response to TLR7 ligation. <i>Journal of Immunology</i> , 2010 , 184, 1159-67	5.3	81
99	Monocyte-mediated inhibition of TLR9-dependent IFN- β induction in plasmacytoid dendritic cells questions bacterial DNA as the active ingredient of bacterial lysates. <i>Journal of Immunology</i> , 2010 , 185, 7367-73	5.3	19
98	Targeted activation of RNA helicase retinoic acid-inducible gene-I induces proimmunogenic apoptosis of human ovarian cancer cells. <i>Cancer Research</i> , 2010 , 70, 5293-304	10.1	65
97	Virally infected mouse liver endothelial cells trigger CD8+ T-cell immunity. <i>Gastroenterology</i> , 2010 , 138, 336-46	13.3	57
96	Effects of an active immunization on the immune response of laying Japanese quail (<i>Coturnix coturnix japonica</i>) fed with or without genetically modified <i>Bacillus thuringiensis</i> -maize. <i>Poultry Science</i> , 2010 , 89, 1122-8	3.9	9
95	Mitf silencing cooperates with IL-12 gene transfer to inhibit melanoma in mice. <i>International Immunopharmacology</i> , 2010 , 10, 540-5	5.8	7
94	The chase for the RIG-I ligand--recent advances. <i>Molecular Therapy</i> , 2010 , 18, 1254-62	11.7	70
93	Dendritic cell vaccination in human melanoma: relationships between clinical effects and vaccine parameters. <i>Pigment Cell and Melanoma Research</i> , 2010 , 23, 607-19	4.5	40
92	Sorafenib in combination with carboplatin and paclitaxel as neoadjuvant chemotherapy in patients with advanced ovarian cancer. <i>Cancer Chemotherapy and Pharmacology</i> , 2010 , 66, 203-7	3.5	46
91	<i>Listeria monocytogenes</i> is sensed by the NLRP3 and AIM2 inflammasome. <i>European Journal of Immunology</i> , 2010 , 40, 1545-51	6.1	199
90	Immunostimulatory RNA oligonucleotides induce an effective antitumoral NK cell response through the TLR7. <i>Journal of Immunology</i> , 2009 , 183, 6078-86	5.3	35
89	Higher activation of TLR9 in plasmacytoid dendritic cells by microbial DNA compared with self-DNA based on CpG-specific recognition of phosphodiester DNA. <i>Journal of Leukocyte Biology</i> , 2009 , 86, 663-70	6.5	27

88	Complete regression of advanced primary and metastatic mouse melanomas following combination chemoimmunotherapy. <i>Cancer Research</i> , 2009 , 69, 6265-74	10.1	43
87	Regulation and function of the cytosolic viral RNA sensor RIG-I in pancreatic beta cells. <i>Biochimica Et Biophysica Acta - Molecular Cell Research</i> , 2009 , 1793, 1768-75	4.9	15
86	Tumour-derived prostaglandin E and transforming growth factor-beta synergize to inhibit plasmacytoid dendritic cell-derived interferon-alpha. <i>Immunology</i> , 2009 , 128, 439-50	7.8	73
85	Syk kinase signalling couples to the Nlrp3 inflammasome for anti-fungal host defence. <i>Nature</i> , 2009 , 459, 433-6	50.4	675
84	RIG-I-dependent sensing of poly(dA:dT) through the induction of an RNA polymerase III-transcribed RNA intermediate. <i>Nature Immunology</i> , 2009 , 10, 1065-72	19.1	645
83	Approaching the RNA ligand for RIG-I?. <i>Immunological Reviews</i> , 2009 , 227, 66-74	11.3	66
82	Recognition of 5'Triphosphate by RIG-I helicase requires short blunt double-stranded RNA as contained in panhandle of negative-strand virus. <i>Immunity</i> , 2009 , 31, 25-34	32.3	564
81	Selection of molecular structure and delivery of RNA oligonucleotides to activate TLR7 versus TLR8 and to induce high amounts of IL-12p70 in primary human monocytes. <i>Journal of Immunology</i> , 2009 , 182, 6824-33	5.3	71
80	Selective and direct activation of human neutrophils but not eosinophils by Toll-like receptor 8. <i>Journal of Allergy and Clinical Immunology</i> , 2009 , 123, 1026-33	11.5	53
79	TLR8-driven IL-12-dependent reciprocal and synergistic activation of NK cells and monocytes by immunostimulatory RNA. <i>Journal of Immunotherapy</i> , 2009 , 32, 262-71	5	20
78	Proapoptotic signaling induced by RIG-I and MDA-5 results in type I interferon-independent apoptosis in human melanoma cells. <i>Journal of Clinical Investigation</i> , 2009 , 119, 2399-411	15.9	270
77	Gene silencing below the immune radar. <i>Journal of Clinical Investigation</i> , 2009 , 119, 438-42	15.9	7
76	5'Triphosphate-siRNA: turning gene silencing and Rig-I activation against melanoma. <i>Nature Medicine</i> , 2008 , 14, 1256-63	50.5	307
75	TRADD protein is an essential component of the RIG-like helicase antiviral pathway. <i>Immunity</i> , 2008 , 28, 651-61	32.3	242
74	RNA recognition via TLR7 and TLR8. <i>Handbook of Experimental Pharmacology</i> , 2008 , 71-86	3.2	63
73	Delivery by cationic gelatin nanoparticles strongly increases the immunostimulatory effects of CpG oligonucleotides. <i>Pharmaceutical Research</i> , 2008 , 25, 551-62	4.5	105
72	Assessing the therapeutic potential of immunostimulatory nucleic acids. <i>Current Opinion in Immunology</i> , 2008 , 20, 389-95	7.8	84
71	RNA Interference in Scope of Immune System 2008 , 207-226		

70	Staphylococcus aureus protein A triggers T cell-independent B cell proliferation by sensitizing B cells for TLR2 ligands. <i>Journal of Immunology</i> , 2007 , 178, 2803-12	5.3	84
69	Immunostimulatory RNA oligonucleotides trigger an antigen-specific cytotoxic T-cell and IgG2a response. <i>Blood</i> , 2007 , 109, 2953-60	2.2	50
68	A mammalian microRNA expression atlas based on small RNA library sequencing. <i>Cell</i> , 2007 , 129, 1401-1462	36.2	3005
67	Immunotherapy with dendritic cells and CpG oligonucleotides can be combined with chemotherapy without loss of efficacy in a mouse model of colon cancer. <i>International Journal of Cancer</i> , 2006 , 118, 2790-5	7.5	35
66	Analysis of plasmacytoid and myeloid dendritic cells in nasal epithelium. <i>Vaccine Journal</i> , 2006 , 13, 1278-86		42
65	Immunostimulatory properties of CpG-oligonucleotides are enhanced by the use of protamine nanoparticles. <i>Oligonucleotides</i> , 2006 , 16, 313-22		35
64	T cell-independent, TLR-induced IL-12p70 production in primary human monocytes. <i>Journal of Immunology</i> , 2006 , 176, 7438-46	5.3	98
63	siRNA and isRNA: two edges of one sword. <i>Molecular Therapy</i> , 2006 , 14, 463-70	11.7	181
62	5RTriphosphate RNA is the ligand for RIG-I. <i>Science</i> , 2006 , 314, 994-7	33.3	1826
61	Inhibition of toll-like receptor 7- and 9-mediated alpha/beta interferon production in human plasmacytoid dendritic cells by respiratory syncytial virus and measles virus. <i>Journal of Virology</i> , 2005 , 79, 5507-15	6.6	196
60	CpG oligonucleotides induce strong humoral but only weak CD4+ T cell responses to protein antigens in rhesus macaques in vivo. <i>Vaccine</i> , 2005 , 23, 3310-7	4.1	20
59	Sequence-specific potent induction of IFN-alpha by short interfering RNA in plasmacytoid dendritic cells through TLR7. <i>Nature Medicine</i> , 2005 , 11, 263-70	50.5	1026
58	Preferential expression and function of Toll-like receptor 3 in human astrocytes. <i>Journal of Neuroimmunology</i> , 2005 , 159, 12-9	3.5	212
57	CpG ODN enhance antigen-specific NKT cell activation via plasmacytoid dendritic cells. <i>European Journal of Immunology</i> , 2005 , 35, 2347-57	6.1	64
56	No indication for a defect in toll-like receptor signaling in patients with hyper-IgE syndrome. <i>Journal of Clinical Immunology</i> , 2005 , 25, 321-8	5.7	14
55	Spontaneous formation of nucleic acid-based nanoparticles is responsible for high interferon-alpha induction by CpG-A in plasmacytoid dendritic cells. <i>Journal of Biological Chemistry</i> , 2005 , 280, 8086-93	5.4	146
54	B-cell lymphomas differ in their responsiveness to CpG oligodeoxynucleotides. <i>Clinical Cancer Research</i> , 2005 , 11, 1490-9	12.9	110
53	Plasmacytoid dendritic cells control TLR7 sensitivity of naive B cells via type I IFN. <i>Journal of Immunology</i> , 2005 , 174, 4043-50	5.3	281

52	Replication-dependent potent IFN-alpha induction in human plasmacytoid dendritic cells by a single-stranded RNA virus. <i>Journal of Immunology</i> , 2004 , 173, 5935-43	5-3	175
51	IL-12p70-dependent Th1 induction by human B cells requires combined activation with CD40 ligand and CpG DNA. <i>Journal of Immunology</i> , 2004 , 172, 954-63	5-3	137
50	CpG-A and CpG-B oligonucleotides differentially enhance human peptide-specific primary and memory CD8+ T-cell responses in vitro. <i>Blood</i> , 2004 , 103, 2162-9	2-2	79
49	CpG oligonucleotides elicit antitumor responses in a human melanoma NOD/SCID xenotransplantation model. <i>Journal of Investigative Dermatology</i> , 2004 , 122, 387-91	4-3	16
48	Structural studies of oligonucleotides containing G-quadruplex motifs using AFM. <i>Biochemical and Biophysical Research Communications</i> , 2004 , 313, 1065-72	3-4	39
47	Role of adenosine receptors in regulating chemotaxis and cytokine production of plasmacytoid dendritic cells. <i>Blood</i> , 2004 , 103, 1391-7	2-2	143
46	Plasmacytoid dendritic cells, antigen, and CpG-C license human B cells for plasma cell differentiation and immunoglobulin production in the absence of T-cell help. <i>Blood</i> , 2004 , 103, 3058-64	2-2	235
45	Technology evaluation: BAY-50-4798, Bayer. <i>Current Opinion in Molecular Therapeutics</i> , 2004 , 6, 221-7		
44	Activation of Dendritic Cells and Induction of T Cell Responses by Hpv 16 L1/E7 Chimeric Virus-Like Particles are Enhanced by Cpg ODN or Sorbitol. <i>Antiviral Therapy</i> , 2004 , 9, 479-489	1-6	13
43	Activation with CpG-A and CpG-B oligonucleotides reveals two distinct regulatory pathways of type I IFN synthesis in human plasmacytoid dendritic cells. <i>Journal of Immunology</i> , 2003 , 170, 4465-74	5-3	282
42	CpG-A oligonucleotides induce a monocyte-derived dendritic cell-like phenotype that preferentially activates CD8 T cells. <i>Journal of Immunology</i> , 2003 , 170, 3468-77	5-3	63
41	CpG: unraveling the key to B-cell function. <i>Blood</i> , 2003 , 101, 4230-4231	2-2	1
40	Rational design of new CpG oligonucleotides that combine B cell activation with high IFN-alpha induction in plasmacytoid dendritic cells. <i>European Journal of Immunology</i> , 2003 , 33, 1633-41	6-1	257
39	Identification and functional analysis of tumor-infiltrating plasmacytoid dendritic cells in head and neck cancer. <i>Cancer Research</i> , 2003 , 63, 6478-87	10-1	230
38	Combined dendritic cell- and CpG oligonucleotide-based immune therapy cures large murine tumors that resist chemotherapy. <i>European Journal of Immunology</i> , 2002 , 32, 3235-45	6-1	100
37	Plasmacytoid dendritic cells: a new cutaneous dendritic cell subset with distinct role in inflammatory skin diseases. <i>Journal of Investigative Dermatology</i> , 2002 , 119, 1096-102	4-3	327
36	CpG oligonucleotides: novel regulators of osteoclast differentiation. <i>FASEB Journal</i> , 2002 , 16, 274-82	0-9	46
35	Peritumoral CpG DNA elicits a coordinated response of CD8 T cells and innate effectors to cure established tumors in a murine colon carcinoma model. <i>Journal of Immunology</i> , 2002 , 169, 3892-9	5-3	163

34	Plasmacytoid dendritic cells: the key to CpG. <i>Human Immunology</i> , 2002 , 63, 1111-9	2.3	122
33	Quantitative expression of toll-like receptor 1-10 mRNA in cellular subsets of human peripheral blood mononuclear cells and sensitivity to CpG oligodeoxynucleotides. <i>Journal of Immunology</i> , 2002 , 168, 4531-7	5.3	1574
32	Shock waves: a novel method for cytoplasmic delivery of antisense oligonucleotides. <i>Journal of Molecular Medicine</i> , 2001 , 79, 306-13	5.5	27
31	Toll-like receptor expression reveals CpG DNA as a unique microbial stimulus for plasmacytoid dendritic cells which synergizes with CD40 ligand to induce high amounts of IL-12. <i>European Journal of Immunology</i> , 2001 , 31, 3026-37	6.1	643
30	Identification of CpG oligonucleotide sequences with high induction of IFN-alpha/beta in plasmacytoid dendritic cells. <i>European Journal of Immunology</i> , 2001 , 31, 2154-63	6.1	733
29	Distinct CpG oligonucleotide sequences activate human gamma delta T cells via interferon-alpha/-beta. <i>European Journal of Immunology</i> , 2001 , 31, 3525-34	6.1	64
28	Antisense therapy in oncology: new hope for an old idea?. <i>Lancet, The</i> , 2001 , 358, 489-97	4.0	205
27	Neutralization of interleukin-18 reduces severity in murine colitis and intestinal IFN-gamma and TNF-alpha production. <i>American Journal of Physiology - Regulatory Integrative and Comparative Physiology</i> , 2001 , 281, R1264-73	3.2	219
26	Toll-like receptor expression reveals CpG DNA as a unique microbial stimulus for plasmacytoid dendritic cells which synergizes with CD40 ligand to induce high amounts of IL-12 2001 , 31, 3026		3
25	Identification of CpG oligonucleotide sequences with high induction of IFN- γ in plasmacytoid dendritic cells 2001 , 31, 2154		2
24	Delineation of a CpG phosphorothioate oligodeoxynucleotide for activating primate immune responses in vitro and in vivo. <i>Journal of Immunology</i> , 2000 , 164, 1617-24	5.3	512
23	Enhanced dendritic cell maturation by TNF-alpha or cytidine-phosphate-guanosine DNA drives T cell activation in vitro and therapeutic anti-tumor immune responses in vivo. <i>Journal of Immunology</i> , 2000 , 165, 6278-86	5.3	148
22	High altitude increases circulating interleukin-6, interleukin-1 receptor antagonist and C-reactive protein. <i>Cytokine</i> , 2000 , 12, 246-52	4	309
21	Mechanism and function of a newly identified CpG DNA motif in human primary B cells. <i>Journal of Immunology</i> , 2000 , 164, 944-53	5.3	530
20	CpG DNA: a potent signal for growth, activation, and maturation of human dendritic cells. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1999 , 96, 9305-10	11.5	525
19	CpG DNA and LPS induce distinct patterns of activation in human monocytes. <i>Gene Therapy</i> , 1999 , 6, 893-903	4	174
18	Mechanisms and therapeutic applications of immune stimulatory cpG DNA 1999 , 84, 113-20		58
17	How to Characterize and Improve Oligonucleotide Uptake into Leukocytes. <i>Perspectives in Antisense Science</i> , 1999 , 41-56		

16	The hairy cell leukemia cell line Eskol spontaneously synthesizes tumor necrosis factor-alpha and nitric oxide. <i>Leukemia Research</i> , 1998 , 22, 501-7	2.7	8
15	Adrenaline enhances LPS-induced IL-10 synthesis: evidence for protein kinase A-mediated pathway. <i>International Journal of Immunopharmacology</i> , 1998 , 20, 57-69		34
14	Raised leptin concentrations at high altitude associated with loss of appetite. <i>Lancet, The</i> , 1998 , 352, 1119-20	40	121
13	Oligonucleotide uptake in leukocytes is dependent on extracellular calcium: a hint for the involvement of adhesion molecules?. <i>Nucleosides & Nucleotides</i> , 1998 , 17, 1767-1773		1
12	In vivo synthesis of tumor necrosis factor-alpha in healthy humans after live yellow fever vaccination. <i>Journal of Infectious Diseases</i> , 1998 , 177, 774-8	7	52
11	Adrenomedullin: a player at high altitude?. <i>Chest</i> , 1998 , 113, 1428	5.3	13
10	Anti-inflammatory activities of cAMP-elevating agents: enhancement of IL-10 synthesis and concurrent suppression of TNF production. <i>Journal of Leukocyte Biology</i> , 1998 , 63, 101-7	6.5	143
9	Plasma adrenomedullin levels, body fluid status, and end-stage renal failure. <i>Nephron</i> , 1998 , 80, 232-3	3.3	7
8	Antisense Strategies for Inhibition of Tumor Necrosis Factor- α Synthesis. <i>Nucleosides & Nucleotides</i> , 1997 , 16, 629-634		
7	Suppression of tumor necrosis factor-alpha production by interleukin-10 is enhanced by cAMP-elevating agents. <i>European Journal of Pharmacology</i> , 1997 , 321, 231-9	5.3	26
6	Oligonucleotides: extrapolating from in vitro to in vivo. <i>Nature Medicine</i> , 1997 , 3, 702	50.5	2
5	Taming TNF: strategies to restrain this proinflammatory cytokine. <i>Trends in Immunology</i> , 1997 , 18, 487-92		237
4	Oligodeoxynucleotides Enhance Lipopolysaccharide-Stimulated Synthesis of Tumor Necrosis Factor: Dependence on Phosphorothioate Modification and Reversal by Heparin. <i>Molecular Medicine</i> , 1996 , 2, 429-438	6.2	76
3	Specific suppression of human tumor necrosis factor-alpha synthesis by antisense oligodeoxynucleotides. <i>Oligonucleotides</i> , 1996 , 6, 291-9		30
2	ADA2 is a lysosomal DNase regulating the type-I interferon response		3
1	RIG-I-induced innate antiviral immunity protects mice from lethal SARS-CoV-2 infection		2