

Joanna Dabrowska

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/7674759/publications.pdf>

Version: 2024-02-01

18
papers

918
citations

623734

14
h-index

839539

18
g-index

19
all docs

19
docs citations

19
times ranked

1051
citing authors

#	ARTICLE	IF	CITATIONS
1	Neuroanatomical evidence for reciprocal regulation of the corticotrophin-releasing factor and oxytocin systems in the hypothalamus and the bed nucleus of the stria terminalis of the rat: Implications for balancing stress and affect. <i>Psychoneuroendocrinology</i> , 2011, 36, 1312-1326.	2.7	210
2	Central CRF neurons are not created equal: phenotypic differences in CRF-containing neurons of the rat paraventricular hypothalamus and the bed nucleus of the stria terminalis. <i>Frontiers in Neuroscience</i> , 2013, 7, 156.	2.8	131
3	Oxytocin in the nucleus accumbens shell reverses CRFR2-evoked passive stress-coping after partner loss in monogamous male prairie voles. <i>Psychoneuroendocrinology</i> , 2016, 64, 66-78.	2.7	116
4	The response of neurons in the bed nucleus of the stria terminalis to serotonin: Implications for anxiety. <i>Progress in Neuro-Psychopharmacology and Biological Psychiatry</i> , 2009, 33, 1309-1320.	4.8	88
5	Striatal-Enriched Protein Tyrosine Phosphataseâ€”STEPS Toward Understanding Chronic Stress-Induced Activation of Corticotrophin Releasing Factor Neurons in the Rat Bed Nucleus of the Stria Terminalis. <i>Biological Psychiatry</i> , 2013, 74, 817-826.	1.3	47
6	Oxytocin facilitates adaptive fear and attenuates anxiety responses in animal models and human studiesâ€”potential interaction with the corticotropin-releasing factor (CRF) system in the bed nucleus of the stria terminalis (BNST). <i>Cell and Tissue Research</i> , 2019, 375, 143-172.	2.9	47
7	A transcriptomic analysis of type Iâ€”III neurons in the bed nucleus of the stria terminalis. <i>Molecular and Cellular Neurosciences</i> , 2011, 46, 699-709.	2.2	42
8	Oxytocin receptors in the dorsolateral bed nucleus of the stria terminalis (BNST) bias fear learning toward temporally predictable cued fear. <i>Translational Psychiatry</i> , 2019, 9, 140.	4.8	38
9	Neuronal diversity of the amygdala and the bed nucleus of the stria terminalis. <i>Handbook of Behavioral Neuroscience</i> , 2020, 26, 63-100.	0.7	34
10	Oxytocin receptor neurotransmission in the dorsolateral bed nucleus of the stria terminalis facilitates the acquisition of cued fear in the fear-potentiated startle paradigm in rats. <i>Neuropharmacology</i> , 2017, 121, 130-139.	4.1	33
11	Oxytocin Promotes Accurate Fear Discrimination and Adaptive Defensive Behaviors. <i>Frontiers in Neuroscience</i> , 2020, 14, 583878.	2.8	27
12	Corticotropin-Releasing Factor Receptors Modulate Oxytocin Release in the Dorsolateral Bed Nucleus of the Stria Terminalis (BNST) in Male Rats. <i>Frontiers in Neuroscience</i> , 2018, 12, 183.	2.8	22
13	Stereoselectivity of 8-OH-DPAT toward the serotonin 5-HT1A receptor: Biochemical and molecular modeling study. <i>Biochemical Pharmacology</i> , 2006, 72, 498-511.	4.4	18
14	Limbic Neuropeptidergic Modulators of Emotion and Their Therapeutic Potential for Anxiety and Post-Traumatic Stress Disorder. <i>Journal of Neuroscience</i> , 2021, 41, 901-910.	3.6	18
15	Desensitization of 5-HT1A autoreceptors induced by neonatal DSP-4 treatment. <i>European Neuropsychopharmacology</i> , 2007, 17, 129-137.	0.7	14
16	Repeated shock stress facilitates basolateral amygdala synaptic plasticity through decreased cAMP-specific phosphodiesterase type IV (PDE4) expression. <i>Brain Structure and Function</i> , 2018, 223, 1731-1745.	2.3	13
17	Reactivity of 5-HT1A receptor in adult rats after neonatal noradrenergic neurons' lesion â€” Implications for antidepressant-like action. <i>Brain Research</i> , 2008, 1239, 66-76.	2.2	10
18	Oxytocin excites BNST interneurons and inhibits BNST output neurons to the central amygdala. <i>Neuropharmacology</i> , 2021, 192, 108601.	4.1	7