

Emi Ushigome

List of Publications by Citations

Source: <https://exaly.com/author-pdf/7674979/emi-ushigome-publications-by-citations.pdf>

Version: 2024-04-19

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

72
papers

562
citations

12
h-index

20
g-index

90
ext. papers

902
ext. citations

4.5
avg, IF

3.81
L-index

#	Paper	IF	Citations
72	The coefficient variation of home blood pressure is a novel factor associated with macroalbuminuria in type 2 diabetes mellitus. <i>Hypertension Research</i> , 2011 , 34, 1271-5	4.7	68
71	Decreased the creatinine to cystatin C ratio is a surrogate marker of sarcopenia in patients with type 2 diabetes. <i>Diabetes Research and Clinical Practice</i> , 2018 , 139, 52-58	7.4	54
70	Late-night-dinner is associated with poor glycemic control in people with type 2 diabetes: The KAMOGAWA-DM cohort study. <i>Endocrine Journal</i> , 2018 , 65, 395-402	2.9	39
69	Uncontrolled home blood pressure in the morning is associated with nephropathy in Japanese type 2 diabetes. <i>Heart and Vessels</i> , 2011 , 26, 609-15	2.1	25
68	Usefulness and Safety of Remote Continuous Glucose Monitoring for a Severe COVID-19 Patient with Diabetes. <i>Diabetes Technology and Therapeutics</i> , 2021 , 23, 78-80	8.1	25
67	Sarcopenia is associated with blood pressure variability in older patients with type 2 diabetes: A cross-sectional study of the KAMOGAWA-DM cohort study. <i>Geriatrics and Gerontology International</i> , 2018 , 18, 1345-1349	2.9	23
66	Effect of coronavirus disease 2019 pandemic on the lifestyle and glycemic control in patients with type 2 diabetes: a cross-section and retrospective cohort study. <i>Endocrine Journal</i> , 2021 , 68, 201-210	2.9	18
65	Combined effect of body mass index and waist-height ratio on incident diabetes; a population based cohort study. <i>Journal of Clinical Biochemistry and Nutrition</i> , 2017 , 61, 118-122	3.1	15
64	Is home blood pressure reporting in patients with type 2 diabetes reliable?. <i>Hypertension Research</i> , 2014 , 37, 741-5	4.7	15
63	Prognostic significance of day-by-day variability of home blood pressure on progression to macroalbuminuria in patients with diabetes. <i>Journal of Hypertension</i> , 2018 , 36, 1068-1075	1.9	14
62	Maximum home systolic blood pressure is a useful indicator of arterial stiffness in patients with type 2 diabetes mellitus: post hoc analysis of a cross-sectional multicenter study. <i>Diabetes Research and Clinical Practice</i> , 2014 , 105, 344-51	7.4	14
61	Protein Intake, Especially Vegetable Protein Intake, Is Associated with Higher Skeletal Muscle Mass in Elderly Patients with Type 2 Diabetes. <i>Journal of Diabetes Research</i> , 2017 , 2017, 7985728	3.9	12
60	High brain natriuretic peptide is associated with sarcopenia in patients with type 2 diabetes: a cross-sectional study of KAMOGAWA-DM cohort study. <i>Endocrine Journal</i> , 2019 , 66, 369-377	2.9	10
59	Impact of masked hypertension on diabetic nephropathy in patients with type II diabetes: a KAMOGAWA-HBP study. <i>Journal of the American Society of Hypertension</i> , 2018 , 12, 364-371.e1		10
58	Optimal home SBP targets for preventing the progression of diabetic nephropathy in patients with type 2 diabetes mellitus. <i>Journal of Hypertension</i> , 2015 , 33, 1853-9; discussion 1859	1.9	10
57	Skipping breakfast is associated with glycemic variability in patients with type 2 diabetes. <i>Nutrition</i> , 2020 , 71, 110639	4.8	10
56	Utility of continuous glucose monitoring following gastrectomy. <i>Gastric Cancer</i> , 2020 , 23, 699-706	7.6	10

55	Threshold value of home pulse pressure predicting arterial stiffness in patients with type 2 diabetes: KAMOGAWA-HBP study. <i>Journal of Clinical Hypertension</i> , 2018 , 20, 472-477	2.3	9
54	Beneficial effect of calcium channel blockers on home blood pressure variability in the morning in patients with type 2 diabetes. <i>Journal of Diabetes Investigation</i> , 2013 , 4, 399-404	3.9	9
53	The Effects of Metformin on the Gut Microbiota of Patients with Type 2 Diabetes: A Two-Center, Quasi-Experimental Study. <i>Life</i> , 2020 , 10,	3	9
52	Group 3 Innate Lymphoid Cells Protect Steatohepatitis From High-Fat Diet Induced Toxicity. <i>Frontiers in Immunology</i> , 2021 , 12, 648754	8.4	9
51	Frequent Usage of Convenience Stores is Associated with Low Diet Quality. <i>Nutrients</i> , 2019 , 11,	6.7	8
50	Isolated high home systolic blood pressure in patients with type 2 diabetes is a prognostic factor for the development of diabetic nephropathy: KAMOGAWA-HBP study. <i>Diabetes Research and Clinical Practice</i> , 2019 , 158, 107920	7.4	7
49	Short energy intake is associated with muscle mass loss in older patients with type 2 diabetes: A prospective study of the KAMOGAWA-DM cohort. <i>Clinical Nutrition</i> , 2021 , 40, 1613-1620	5.9	7
48	The effect of COVID-19 pandemic on the lifestyle and glycemic control in patients with type 1 diabetes: a retrospective cohort study. <i>Diabetology International</i> , 2021 , 1-6	2.3	7
47	Reduced dietary omega-3 fatty acids intake is associated with sarcopenia in elderly patients with type 2 diabetes: a cross-sectional study of KAMOGAWA-DM cohort study. <i>Journal of Clinical Biochemistry and Nutrition</i> , 2020 , 66, 233-237	3.1	6
46	Morning pulse pressure is associated more strongly with elevated albuminuria than systolic blood pressure in patients with type 2 diabetes mellitus: post hoc analysis of a cross-sectional multicenter study. <i>Diabetes Research and Clinical Practice</i> , 2013 , 101, 270-7	7.4	6
45	Home Blood Pressure Variability From the Stored Memory Is Correlated With Albuminuria, but From the Logbook Is not. <i>American Journal of Hypertension</i> , 2017 , 30, 993-998	2.3	6
44	Habitual Miso (Fermented Soybean Paste) Consumption Is Associated with a Low Prevalence of Sarcopenia in Patients with Type 2 Diabetes: A Cross-Sectional Study. <i>Nutrients</i> , 2020 , 13,	6.7	6
43	Low circulating dihomo-gamma-linolenic acid is associated with diabetic retinopathy: a cross sectional study of KAMOGAWA-DM cohort study. <i>Endocrine Journal</i> , 2021 , 68, 421-428	2.9	6
42	Japanese radio calisthenics prevents the reduction of skeletal muscle mass volume in people with type 2 diabetes. <i>BMJ Open Diabetes Research and Care</i> , 2020 , 8,	4.5	5
41	Reduced Energy Intake Is Associated with Sarcopenia in Elderly Patients with Type 2 Diabetes: Kamogawa-DM Cohort Study. <i>Diabetes</i> , 2018 , 67, 774-P	0.9	5
40	Seasonal variation in home blood pressure and its relationship with room temperature in patients with type 2 diabetes. <i>Diabetes and Vascular Disease Research</i> , 2020 , 17, 1479164119883986	3.3	5
39	Eating Fast Is Associated with Nonalcoholic Fatty Liver Disease in Men But Not in Women with Type 2 Diabetes: A Cross-Sectional Study. <i>Nutrients</i> , 2020 , 12,	6.7	5
38	Trans Fatty Acid Intake Induces Intestinal Inflammation and Impaired Glucose Tolerance. <i>Frontiers in Immunology</i> , 2021 , 12, 669672	8.4	5

37	Erythritol Ameliorates Small Intestinal Inflammation Induced by High-Fat Diets and Improves Glucose Tolerance. <i>International Journal of Molecular Sciences</i> , 2021 , 22,	6.3	5
36	Predictive power of home blood pressure measurement for cardiovascular outcomes in patients with type 2 diabetes: KAMOGAWA-HBP study. <i>Hypertension Research</i> , 2021 , 44, 348-354	4.7	5
35	Home blood pressure is associated with cognitive impairment among elderly patients with type 2 diabetes: KAMOGAWA-HBP study. <i>Diabetes and Vascular Disease Research</i> , 2019 , 16, 506-512	3.3	4
34	Prognostic significance of home pulse pressure for progression of diabetic nephropathy: KAMOGAWA-HBP study. <i>Hypertension Research</i> , 2018 , 41, 363-371	4.7	4
33	Understanding of antidiabetic medication is associated with blood glucose in patients with type 2 diabetes: At baseline date of the KAMOGAWA-DM cohort study. <i>Journal of Diabetes Investigation</i> , 2019 , 10, 458-465	3.9	4
32	ILC2s Improve Glucose Metabolism Through the Control of Saturated Fatty Acid Absorption Within Visceral Fat. <i>Frontiers in Immunology</i> , 2021 , 12, 669629	8.4	4
31	Olmесartan with azelnidipine versus with trichlormethiazide on home blood pressure variability in patients with type II diabetes mellitus. <i>Journal of the American Society of Hypertension</i> , 2017 , 11, 140-147		3
30	Maximum morning home systolic blood pressure is an indicator of the development of diabetic nephropathy: The KAMOGAWA-HBP study. <i>Journal of Diabetes Investigation</i> , 2019 , 10, 1543-1549	3.9	3
29	Living alone is associated with visit-to-visit HbA1c variability in men but not in women in people with type 2 diabetes: KAMOGAWA-DM cohort study. <i>Endocrine Journal</i> , 2020 , 67, 419-426	2.9	3
28	Caffeine intake enhances the benefits of sodium glucose transporter 2 inhibitor. <i>Diabetes/Metabolism Research and Reviews</i> , 2016 , 32, 694-699	7.5	3
27	Serum N-terminal Pro-brain Natriuretic Peptide Level is Associated with the Development of Chronic Kidney Diseases in Patients with Type 2 Diabetes. <i>Endocrine, Metabolic and Immune Disorders - Drug Targets</i> , 2018 , 18, 590-595	2.2	3
26	Habitual Dietary Intake Affects the Altered Pattern of Gut Microbiome by Acarbose in Patients with Type 2 Diabetes. <i>Nutrients</i> , 2021 , 13,	6.7	3
25	Critical discrepancy in blood glucose control levels evaluated by glycated albumin and estimated hemoglobin A1c levels determined from a flash continuous glucose monitoring system in patients with type 2 diabetes on hemodialysis. <i>Journal of Diabetes Investigation</i> , 2020 , 11, 1570-1574	3.9	2
24	Effects of dietary salt restriction on home blood pressure in diabetic patients with excessive salt intake: a pilot study. <i>Journal of Clinical Biochemistry and Nutrition</i> , 2019 , 65, 252-257	3.1	2
23	Home-measured heart rate is associated with albuminuria in patients with type 2 diabetes mellitus: a post-hoc analysis of a cross-sectional multicenter study. <i>Hypertension Research</i> , 2014 , 37, 533-7	4.7	2
22	Extracellular lipidome change by an SGLT2 inhibitor, luseogliflozin, contributes to prevent skeletal muscle atrophy in db/db mice. <i>Journal of Cachexia, Sarcopenia and Muscle</i> , 2021 ,	10.3	2
21	Impact of Isolated High Home Systolic Blood Pressure and Diabetic Nephropathy in Patients with Type 2 Diabetes Mellitus: A 5-Year Prospective Cohort Study. <i>Journal of Clinical Medicine</i> , 2021 , 10,	5.1	2
20	Vitamin Intake and Loss of Muscle Mass in Older People with Type 2 Diabetes: A Prospective Study of the KAMOGAWA-DM Cohort. <i>Nutrients</i> , 2021 , 13,	6.7	2

19	Usefulness of Exercise for Home Blood Pressure Control in People with Diabetes: A Study Protocol for a Crossover Randomized Controlled Trial. <i>Diabetes, Metabolic Syndrome and Obesity: Targets and Therapy</i> , 2020 , 13, 4747-4753	3.4	1
18	Maximum home blood pressure is a useful indicator of diabetic nephropathy in patients with type 2 diabetes mellitus: KAMOGAWA-HBP study. <i>Diabetes and Vascular Disease Research</i> , 2017 , 14, 477-484	3.3	1
17	Dynamics of glucose levels after Billroth I versus Roux-en-Y reconstruction in patients who undergo distal gastrectomy. <i>Surgery Today</i> , 2021 , 1	3	1
16	Association between Geriatric Nutrition Risk Index and The Presence of Sarcopenia in People with Type 2 Diabetes Mellitus: A Cross-Sectional Study. <i>Nutrients</i> , 2021 , 13,	6.7	1
15	Impact of extracellular-to-intracellular fluid volume ratio on albuminuria in patients with type 2 diabetes: A cross-sectional and longitudinal cohort study. <i>Journal of Diabetes Investigation</i> , 2021 , 12, 1202-1211	3.9	1
14	Habitual Miso (Fermented Soybean Paste) Consumption Is Associated with Glycemic Variability in Patients with Type 2 Diabetes: A Cross-Sectional Study. <i>Nutrients</i> , 2021 , 13,	6.7	1
13	Diabetic nephropathy ameliorated in patients with normal home blood pressure compared to those with isolated high home systolic blood pressure: A 5-year prospective cohort study among patients with type 2 diabetes mellitus.. <i>Diabetes and Vascular Disease Research</i> , 2022 , 19, 14791641221098193	3.3	1
12	Impact of untreated diabetes and COVID-19-related diabetes on severe COVID-19.. <i>Heliyon</i> , 2022 , e088036	3.6	0
11	Asymptomatic postprandial hypotension in patients with diabetes: The KAMOGAWA-HBP study. <i>Journal of Diabetes Investigation</i> , 2021 , 12, 837-844	3.9	0
10	Sarcopenia Is Associated With a Risk of Mortality in People With Type 2 Diabetes Mellitus. <i>Frontiers in Endocrinology</i> , 2021 , 12, 783363	5.7	0
9	Effect of probiotics, G9-1, on gastrointestinal symptoms in patients with type 2 diabetes mellitus: study protocol for open-label, single-arm, exploratory research trial (Big STAR study). <i>Journal of Clinical Biochemistry and Nutrition</i> , 2020 , 67, 223-227	3.1	0
8	Evaluation of the efficacy of simplified nutritional instructions from physicians on dietary salt restriction for patients with type 2 diabetes mellitus consuming excessive salt: protocol for a randomized controlled trial. <i>Trials</i> , 2019 , 20, 761	2.8	0
7	Status of online diet management program users in Japan during the 2020 Coronavirus disease 2019 pandemic. <i>Journal of Clinical Biochemistry and Nutrition</i> , 2021 , 69, 305-310	3.1	0
6	Multicenter, Open-Label, 2-Arm, Pilot Trial for Safe Reduction of Basal Insulin Dose Combined with SGLT2 Inhibitor in Type 1 Diabetes Mellitus: Study Protocol for a RISING-STAR Trial. <i>Clinical Medicine Insights: Endocrinology and Diabetes</i> , 2021 , 14, 11795514211040539	4.3	0
5	Low circulating arachidonic acid is associated with macroalbuminuria in diabetic patients: a cross-sectional examination of the KAMOGAWA-DM cohort study. <i>BMC Nephrology</i> , 2021 , 22, 68	2.7	0
4	Characterization of Peripheral Blood TCR in Patients with Type 1 Diabetes Mellitus by BD Rhapsody™ VDJ CDR3 Assay. <i>Cells</i> , 2022 , 11, 1623	7.9	0
3	Relationship between eosinophils counts and muscle mass decline in older people with type 2 diabetes: A prospective study of the KAMOGAWA-DM cohort.. <i>Experimental Gerontology</i> , 2022 , 159, 111671	4.5	1
2	Development of application to automatically calculate mean amplitude of glycaemic excursions using intermittently scanned continuous glucose monitoring data. <i>Diabetes, Obesity and Metabolism</i> , 2021 , 23, 2155-2160	6.7	

- 1 Association between the frequency of toothbrushing and lifestyle in people with type 2 diabetes mellitus: at the baseline date of the Kamogawa-DM cohort study. *Journal of Clinical Biochemistry and Nutrition*, **2021**, 69, 294-298

3.1