

Emi Ushigome

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/7674979/emi-ushigome-publications-by-year.pdf>

Version: 2024-04-19

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

72
papers

562
citations

12
h-index

20
g-index

90
ext. papers

902
ext. citations

4.5
avg, IF

3.81
L-index

#	Paper	IF	Citations
72	Impact of untreated diabetes and COVID-19-related diabetes on severe COVID-19.. <i>Heliyon</i> , 2022 , e088036	3.6	0
71	Relationship between eosinophils counts and muscle mass decline in older people with type 2 diabetes: A prospective study of the KAMOGAWA-DM cohort.. <i>Experimental Gerontology</i> , 2022 , 159, 111671	4.5	1
70	Diabetic nephropathy ameliorated in patients with normal home blood pressure compared to those with isolated high home systolic blood pressure: A 5-year prospective cohort study among patients with type 2 diabetes mellitus.. <i>Diabetes and Vascular Disease Research</i> , 2022 , 19, 14791641221098193	3.3	1
69	Characterization of Peripheral Blood TCR in Patients with Type 1 Diabetes Mellitus by BD Rhapsody™ VDJ CDR3 Assay. <i>Cells</i> , 2022 , 11, 1623	7.9	0
68	Asymptomatic postprandial hypotension in patients with diabetes: The KAMOGAWA-HBP study. <i>Journal of Diabetes Investigation</i> , 2021 , 12, 837-844	3.9	0
67	Sarcopenia Is Associated With a Risk of Mortality in People With Type 2 Diabetes Mellitus. <i>Frontiers in Endocrinology</i> , 2021 , 12, 783363	5.7	0
66	Dynamics of glucose levels after Billroth I versus Roux-en-Y reconstruction in patients who undergo distal gastrectomy. <i>Surgery Today</i> , 2021 , 1	3	1
65	Extracellular lipidome change by an SGLT2 inhibitor, luseogliflozin, contributes to prevent skeletal muscle atrophy in db/db mice. <i>Journal of Cachexia, Sarcopenia and Muscle</i> , 2021 ,	10.3	2
64	Association between Geriatric Nutrition Risk Index and The Presence of Sarcopenia in People with Type 2 Diabetes Mellitus: A Cross-Sectional Study. <i>Nutrients</i> , 2021 , 13,	6.7	1
63	Impact of extracellular-to-intracellular fluid volume ratio on albuminuria in patients with type 2 diabetes: A cross-sectional and longitudinal cohort study. <i>Journal of Diabetes Investigation</i> , 2021 , 12, 1202-1211	3.9	1
62	Group 3 Innate Lymphoid Cells Protect Steatohepatitis From High-Fat Diet Induced Toxicity. <i>Frontiers in Immunology</i> , 2021 , 12, 648754	8.4	9
61	Impact of Isolated High Home Systolic Blood Pressure and Diabetic Nephropathy in Patients with Type 2 Diabetes Mellitus: A 5-Year Prospective Cohort Study. <i>Journal of Clinical Medicine</i> , 2021 , 10,	5.1	2
60	Short energy intake is associated with muscle mass loss in older patients with type 2 diabetes: A prospective study of the KAMOGAWA-DM cohort. <i>Clinical Nutrition</i> , 2021 , 40, 1613-1620	5.9	7
59	Habitual Miso (Fermented Soybean Paste) Consumption Is Associated with Glycemic Variability in Patients with Type 2 Diabetes: A Cross-Sectional Study. <i>Nutrients</i> , 2021 , 13,	6.7	1
58	Trans Fatty Acid Intake Induces Intestinal Inflammation and Impaired Glucose Tolerance. <i>Frontiers in Immunology</i> , 2021 , 12, 669672	8.4	5
57	The effect of COVID-19 pandemic on the lifestyle and glycemic control in patients with type 1 diabetes: a retrospective cohort study. <i>Diabetology International</i> , 2021 , 1-6	2.3	7
56	Erythritol Ameliorates Small Intestinal Inflammation Induced by High-Fat Diets and Improves Glucose Tolerance. <i>International Journal of Molecular Sciences</i> , 2021 , 22,	6.3	5

55	Development of application to automatically calculate mean amplitude of glycaemic excursions using intermittently scanned continuous glucose monitoring data. <i>Diabetes, Obesity and Metabolism</i> , 2021 , 23, 2155-2160	6.7	
54	Habitual Dietary Intake Affects the Altered Pattern of Gut Microbiome by Acarbose in Patients with Type 2 Diabetes. <i>Nutrients</i> , 2021 , 13,	6.7	3
53	ILC2s Improve Glucose Metabolism Through the Control of Saturated Fatty Acid Absorption Within Visceral Fat. <i>Frontiers in Immunology</i> , 2021 , 12, 669629	8.4	4
52	Usefulness and Safety of Remote Continuous Glucose Monitoring for a Severe COVID-19 Patient with Diabetes. <i>Diabetes Technology and Therapeutics</i> , 2021 , 23, 78-80	8.1	25
51	Predictive power of home blood pressure measurement for cardiovascular outcomes in patients with type 2 diabetes: KAMOGAWA-HBP study. <i>Hypertension Research</i> , 2021 , 44, 348-354	4.7	5
50	Effect of coronavirus disease 2019 pandemic on the lifestyle and glycemic control in patients with type 2 diabetes: a cross-section and retrospective cohort study. <i>Endocrine Journal</i> , 2021 , 68, 201-210	2.9	18
49	Association between the frequency of toothbrushing and lifestyle in people with type 2 diabetes mellitus: at the baseline date of the Kamogawa-DM cohort study. <i>Journal of Clinical Biochemistry and Nutrition</i> , 2021 , 69, 294-298	3.1	
48	Status of online diet management program users in Japan during the 2020 Coronavirus disease 2019 pandemic. <i>Journal of Clinical Biochemistry and Nutrition</i> , 2021 , 69, 305-310	3.1	0
47	Multicenter, Open-Label, 2-Arm, Pilot Trial for Safe Reduction of Basal Insulin Dose Combined with SGLT2 Inhibitor in Type 1 Diabetes Mellitus: Study Protocol for a RISING-STAR Trial. <i>Clinical Medicine Insights: Endocrinology and Diabetes</i> , 2021 , 14, 11795514211040539	4.3	0
46	Low circulating arachidonic acid is associated with macroalbuminuria in diabetic patients: a cross-sectional examination of the KAMOGAWA-DM cohort study. <i>BMC Nephrology</i> , 2021 , 22, 68	2.7	0
45	Vitamin Intake and Loss of Muscle Mass in Older People with Type 2 Diabetes: A Prospective Study of the KAMOGAWA-DM Cohort. <i>Nutrients</i> , 2021 , 13,	6.7	2
44	Low circulating dihomo-gamma-linolenic acid is associated with diabetic retinopathy: a cross sectional study of KAMOGAWA-DM cohort study. <i>Endocrine Journal</i> , 2021 , 68, 421-428	2.9	6
43	Usefulness of Exercise for Home Blood Pressure Control in People with Diabetes: A Study Protocol for a Crossover Randomized Controlled Trial. <i>Diabetes, Metabolic Syndrome and Obesity: Targets and Therapy</i> , 2020 , 13, 4747-4753	3.4	1
42	Critical discrepancy in blood glucose control levels evaluated by glycated albumin and estimated hemoglobin A1c levels determined from a flash continuous glucose monitoring system in patients with type 2 diabetes on hemodialysis. <i>Journal of Diabetes Investigation</i> , 2020 , 11, 1570-1574	3.9	2
41	Japanese radio calisthenics prevents the reduction of skeletal muscle mass volume in people with type 2 diabetes. <i>BMJ Open Diabetes Research and Care</i> , 2020 , 8,	4.5	5
40	Reduced dietary omega-3 fatty acids intake is associated with sarcopenia in elderly patients with type 2 diabetes: a cross-sectional study of KAMOGAWA-DM cohort study. <i>Journal of Clinical Biochemistry and Nutrition</i> , 2020 , 66, 233-237	3.1	6
39	Living alone is associated with visit-to-visit HbA1c variability in men but not in women in people with type 2 diabetes: KAMOGAWA-DM cohort study. <i>Endocrine Journal</i> , 2020 , 67, 419-426	2.9	3
38	Habitual Miso (Fermented Soybean Paste) Consumption Is Associated with a Low Prevalence of Sarcopenia in Patients with Type 2 Diabetes: A Cross-Sectional Study. <i>Nutrients</i> , 2020 , 13,	6.7	6

37	Effect of probiotics, G9-1, on gastrointestinal symptoms in patients with type 2 diabetes mellitus: study protocol for open-label, single-arm, exploratory research trial (Big STAR study). <i>Journal of Clinical Biochemistry and Nutrition</i> , 2020 , 67, 223-227	3.1	0
36	Skipping breakfast is associated with glycemic variability in patients with type 2 diabetes. <i>Nutrition</i> , 2020 , 71, 110639	4.8	10
35	Utility of continuous glucose monitoring following gastrectomy. <i>Gastric Cancer</i> , 2020 , 23, 699-706	7.6	10
34	Seasonal variation in home blood pressure and its relationship with room temperature in patients with type 2 diabetes. <i>Diabetes and Vascular Disease Research</i> , 2020 , 17, 1479164119883986	3.3	5
33	Eating Fast Is Associated with Nonalcoholic Fatty Liver Disease in Men But Not in Women with Type 2 Diabetes: A Cross-Sectional Study. <i>Nutrients</i> , 2020 , 12,	6.7	5
32	The Effects of Metformin on the Gut Microbiota of Patients with Type 2 Diabetes: A Two-Center, Quasi-Experimental Study. <i>Life</i> , 2020 , 10,	3	9
31	Frequent Usage of Convenience Stores is Associated with Low Diet Quality. <i>Nutrients</i> , 2019 , 11,	6.7	8
30	High brain natriuretic peptide is associated with sarcopenia in patients with type 2 diabetes: a cross-sectional study of KAMOGAWA-DM cohort study. <i>Endocrine Journal</i> , 2019 , 66, 369-377	2.9	10
29	Home blood pressure is associated with cognitive impairment among elderly patients with type 2 diabetes: KAMOGAWA-HBP study. <i>Diabetes and Vascular Disease Research</i> , 2019 , 16, 506-512	3.3	4
28	Maximum morning home systolic blood pressure is an indicator of the development of diabetic nephropathy: The KAMOGAWA-HBP study. <i>Journal of Diabetes Investigation</i> , 2019 , 10, 1543-1549	3.9	3
27	Understanding of antidiabetic medication is associated with blood glucose in patients with type 2 diabetes: At baseline date of the KAMOGAWA-DM cohort study. <i>Journal of Diabetes Investigation</i> , 2019 , 10, 458-465	3.9	4
26	Effects of dietary salt restriction on home blood pressure in diabetic patients with excessive salt intake: a pilot study. <i>Journal of Clinical Biochemistry and Nutrition</i> , 2019 , 65, 252-257	3.1	2
25	Isolated high home systolic blood pressure in patients with type 2 diabetes is a prognostic factor for the development of diabetic nephropathy: KAMOGAWA-HBP study. <i>Diabetes Research and Clinical Practice</i> , 2019 , 158, 107920	7.4	7
24	Evaluation of the efficacy of simplified nutritional instructions from physicians on dietary salt restriction for patients with type 2 diabetes mellitus consuming excessive salt: protocol for a randomized controlled trial. <i>Trials</i> , 2019 , 20, 761	2.8	0
23	Threshold value of home pulse pressure predicting arterial stiffness in patients with type 2 diabetes: KAMOGAWA-HBP study. <i>Journal of Clinical Hypertension</i> , 2018 , 20, 472-477	2.3	9
22	Decreased the creatinine to cystatin C ratio is a surrogate marker of sarcopenia in patients with type 2 diabetes. <i>Diabetes Research and Clinical Practice</i> , 2018 , 139, 52-58	7.4	54
21	Prognostic significance of day-by-day variability of home blood pressure on progression to macroalbuminuria in patients with diabetes. <i>Journal of Hypertension</i> , 2018 , 36, 1068-1075	1.9	14
20	Prognostic significance of home pulse pressure for progression of diabetic nephropathy: KAMOGAWA-HBP study. <i>Hypertension Research</i> , 2018 , 41, 363-371	4.7	4

19	Impact of masked hypertension on diabetic nephropathy in patients with type II diabetes: a KAMOGAWA-HBP study. <i>Journal of the American Society of Hypertension</i> , 2018 , 12, 364-371.e1		10
18	Sarcopenia is associated with blood pressure variability in older patients with type 2 diabetes: A cross-sectional study of the KAMOGAWA-DM cohort study. <i>Geriatrics and Gerontology International</i> , 2018 , 18, 1345-1349	2.9	23
17	Serum N-terminal Pro-brain Natriuretic Peptide Level is Associated with the Development of Chronic Kidney Diseases in Patients with Type 2 Diabetes. <i>Endocrine, Metabolic and Immune Disorders - Drug Targets</i> , 2018 , 18, 590-595	2.2	3
16	Reduced Energy Intake Is Associated with Sarcopenia in Elderly Patients with Type 2 Diabetes KAMOGAWA-DM Cohort Study. <i>Diabetes</i> , 2018 , 67, 774-P	0.9	5
15	Late-night-dinner is associated with poor glycemic control in people with type 2 diabetes: The KAMOGAWA-DM cohort study. <i>Endocrine Journal</i> , 2018 , 65, 395-402	2.9	39
14	Olmesartan with azelnidipine versus with trichlormethiazide on home blood pressure variability in patients with type II diabetes mellitus. <i>Journal of the American Society of Hypertension</i> , 2017 , 11, 140-147		3
13	Combined effect of body mass index and waist-height ratio on incident diabetes; a population based cohort study. <i>Journal of Clinical Biochemistry and Nutrition</i> , 2017 , 61, 118-122	3.1	15
12	Maximum home blood pressure is a useful indicator of diabetic nephropathy in patients with type 2 diabetes mellitus: KAMOGAWA-HBP study. <i>Diabetes and Vascular Disease Research</i> , 2017 , 14, 477-484	3.3	1
11	Home Blood Pressure Variability From the Stored Memory Is Correlated With Albuminuria, but From the Logbook Is not. <i>American Journal of Hypertension</i> , 2017 , 30, 993-998	2.3	6
10	Protein Intake, Especially Vegetable Protein Intake, Is Associated with Higher Skeletal Muscle Mass in Elderly Patients with Type 2 Diabetes. <i>Journal of Diabetes Research</i> , 2017 , 2017, 7985728	3.9	12
9	Caffeine intake enhances the benefits of sodium glucose transporter 2 inhibitor. <i>Diabetes/Metabolism Research and Reviews</i> , 2016 , 32, 694-699	7.5	3
8	Optimal home SBP targets for preventing the progression of diabetic nephropathy in patients with type 2 diabetes mellitus. <i>Journal of Hypertension</i> , 2015 , 33, 1853-9; discussion 1859	1.9	10
7	Maximum home systolic blood pressure is a useful indicator of arterial stiffness in patients with type 2 diabetes mellitus: post hoc analysis of a cross-sectional multicenter study. <i>Diabetes Research and Clinical Practice</i> , 2014 , 105, 344-51	7.4	14
6	Is home blood pressure reporting in patients with type 2 diabetes reliable?. <i>Hypertension Research</i> , 2014 , 37, 741-5	4.7	15
5	Home-measured heart rate is associated with albuminuria in patients with type 2 diabetes mellitus: a post-hoc analysis of a cross-sectional multicenter study. <i>Hypertension Research</i> , 2014 , 37, 533-7	4.7	2
4	Morning pulse pressure is associated more strongly with elevated albuminuria than systolic blood pressure in patients with type 2 diabetes mellitus: post hoc analysis of a cross-sectional multicenter study. <i>Diabetes Research and Clinical Practice</i> , 2013 , 101, 270-7	7.4	6
3	Beneficial effect of calcium channel blockers on home blood pressure variability in the morning in patients with type 2 diabetes. <i>Journal of Diabetes Investigation</i> , 2013 , 4, 399-404	3.9	9
2	Uncontrolled home blood pressure in the morning is associated with nephropathy in Japanese type 2 diabetes. <i>Heart and Vessels</i> , 2011 , 26, 609-15	2.1	25

1 The coefficient variation of home blood pressure is a novel factor associated with macroalbuminuria in type 2 diabetes mellitus. *Hypertension Research*, **2011**, 34, 1271-5

4·7 68