

Hiroyoshi Taketani

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/7675875/publications.pdf>

Version: 2024-02-01

23
papers

522
citations

686830

13
h-index

676716

22
g-index

24
all docs

24
docs citations

24
times ranked

1074
citing authors

#	ARTICLE	IF	CITATIONS
1	Effect of sodium glucose cotransporter 2 inhibitor on liver function tests in Japanese patients with nonalcoholic fatty liver disease and type 2 diabetes mellitus. <i>Hepatology Research</i> , 2017, 47, 1072-1078.	1.8	72
2	Lower levels of insulin-like growth factor-1 standard deviation score are associated with histological severity of nonalcoholic fatty liver disease. <i>Hepatology Research</i> , 2015, 45, 771-781.	1.8	64
3	Effect of 12-week dulaglutide therapy in Japanese patients with biopsy-proven nonalcoholic fatty liver disease and type 2 diabetes mellitus. <i>Hepatology Research</i> , 2017, 47, 1206-1211.	1.8	64
4	Development of hepatocellular carcinoma in Japanese patients with biopsy-proven nonalcoholic fatty liver disease: Association between PNPLA3 genotype and hepatocarcinogenesis/fibrosis progression. <i>Hepatology Research</i> , 2017, 47, 1083-1092.	1.8	61
5	Serum alanine aminotransferase predicts the histological course of nonalcoholic steatohepatitis in Japanese patients. <i>Hepatology Research</i> , 2015, 45, E53-61.	1.8	46
6	Combination of PNPLA3 and TLL1 polymorphism can predict advanced fibrosis in Japanese patients with nonalcoholic fatty liver disease. <i>Journal of Gastroenterology</i> , 2018, 53, 438-448.	2.3	24
7	Insulin resistance increases the risk of incident type 2 diabetes mellitus in patients with non-alcoholic fatty liver disease. <i>Hepatology Research</i> , 2018, 48, E42-E51.	1.8	23
8	Hepatic nucleotide binding oligomerization domain-like receptors pyrin domain-containing 3 inflammasomes are associated with the histologic severity of nonalcoholic fatty liver disease. <i>Hepatology Research</i> , 2017, 47, 1459-1468.	1.8	20
9	Increase in the skeletal muscle mass to body fat mass ratio predicts the decline in transaminase in patients with nonalcoholic fatty liver disease. <i>Journal of Gastroenterology</i> , 2019, 54, 160-170.	2.3	20
10	Presence of varices in patients after hepatitis C virus eradication predicts deterioration in the FIB-4 index. <i>Hepatology Research</i> , 2019, 49, 473-478.	1.8	19
11	Association of coronary artery calcification with liver fibrosis in Japanese patients with nonalcoholic fatty liver disease. <i>Hepatology Research</i> , 2016, 46, 1107-1117.	1.8	17
12	Liver stiffness measurement to platelet ratio index predicts the stage of liver fibrosis in nonalcoholic fatty liver disease. <i>Hepatology Research</i> , 2017, 47, 721-730.	1.8	16
13	Clinical and pathological features of sarcopenia-related indices in patients with nonalcoholic fatty liver disease. <i>Hepatology Research</i> , 2019, 49, 627-636.	1.8	16
14	FIB-4 Index and Diabetes Mellitus Are Associated with Chronic Kidney Disease in Japanese Patients with Non-Alcoholic Fatty Liver Disease. <i>International Journal of Molecular Sciences</i> , 2020, 21, 171.	1.8	16
15	Honokiol Acts as a Potent Anti-Fibrotic Agent in the Liver through Inhibition of TGF- β 1/SMAD Signaling and Autophagy in Hepatic Stellate Cells. <i>International Journal of Molecular Sciences</i> , 2021, 22, 13354.	1.8	9
16	Effect of Sodium Glucose Cotransporter 2 Inhibitors on Renal Function in Patients with Nonalcoholic Fatty Liver Disease and Type 2 Diabetes in Japan. <i>Diagnostics</i> , 2020, 10, 86.	1.3	8
17	Honokiol Prevents Non-Alcoholic Steatohepatitis-Induced Liver Cancer via EGFR Degradation through the Glucocorticoid Receptor-MIG6 Axis. <i>Cancers</i> , 2021, 13, 1515.	1.7	7
18	Enhanced Antitumor Effect in Liver Cancer by Amino Acid Depletion-Induced Oxidative Stress. <i>Frontiers in Oncology</i> , 2021, 11, 758549.	1.3	6

#	ARTICLE	IF	CITATIONS
19	Drug-induced liver injury in a chronic hepatitis C patient treated by peginterferon, ribavirin and simeprevir. <i>Hepatology Research</i> , 2015, 45, E156-60.	1.8	5
20	<p>Aging-associated impairment in metabolic compensation by subcutaneous adipose tissue promotes diet-induced fatty liver disease in mice</p>. <i>Diabetes, Metabolic Syndrome and Obesity: Targets and Therapy</i> , 2019, Volume 12, 1473-1492.	1.1	4
21	Loss of KAP3 decreases intercellular adhesion and impairs intracellular transport of laminin in signet ring cell carcinoma of the stomach. <i>Scientific Reports</i> , 2022, 12, 5050.	1.6	3
22	SOX2 enhances cell survival and induces resistance to apoptosis under serum starvation conditions through the AKT/GSK β signaling pathway in esophageal squamous cell carcinoma. <i>Oncology Letters</i> , 2021, 21, 269.	0.8	2
23	The Association between the Platelet Count and Liver Volume in Compensated Cirrhosis Patients after the Eradication of Hepatitis C virus by Direct-acting Antivirals. <i>Internal Medicine</i> , 2020, 59, 1811-1817.	0.3	0