Eliana Janette Ojeda Lazo

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/768703/publications.pdf Version: 2024-02-01



#	Article	IF	CITATIONS
1	A large multinational study of vasomotor symptom prevalence, duration, and impact on quality of life in middle-aged women. Menopause, 2011, 18, 778-785.	0.8	135
2	A multinational study of sleep disorders during female mid-life. Maturitas, 2012, 72, 359-366.	1.0	125
3	Sexual dysfunction in middle-aged women. Menopause, 2009, 16, 1139-1148.	0.8	112
4	Impaired quality of life among middle aged women: A multicentre Latin American study. Maturitas, 2008, 61, 323-329.	1.0	109
5	Age at menopause in Latin America. Menopause, 2006, 13, 706-712.	0.8	105
6	The US National Cholesterol Education Programme Adult Treatment Panel III (NCEP ATP III): prevalence of the metabolic syndrome in postmenopausal Latin American women. Climacteric, 2007, 10, 164-170.	1.1	91
7	Sedentary lifestyle in middle-aged women is associated with severe menopausal symptoms and obesity. Menopause, 2016, 23, 488-493.	0.8	60
8	Menopausal symptoms appear before the menopause and persist 5 years beyond: a detailed analysis of a multinational study. Climacteric, 2012, 15, 542-551.	1.1	59
9	Type II diabetes mellitus and menopause: a multinational study. Climacteric, 2013, 16, 663-672.	1.1	57
10	Obesity and its relation to depressive symptoms and sedentary lifestyle in middle-aged women. Maturitas, 2015, 80, 100-105.	1.0	56
11	Menopause could be involved in the pathogenesis of muscle and joint aches in mid-aged women. Maturitas, 2013, 75, 94-100.	1.0	32
12	Association between anxiety and severe quality-of-life impairment in postmenopausal women: analysis of a multicenter Latin American cross-sectional study. Menopause, 2017, 24, 645-652.	0.8	29
13	Optimal waist circumference cutoff value for defining the metabolic syndrome in postmenopausal Latin American women. Menopause, 2012, 19, 433-437.	0.8	23
14	Metformin use is associated with a lower risk of osteoporosis in adult women independent of type 2 diabetes mellitus and obesity. REDLINC IX study. Gynecological Endocrinology, 2020, 36, 421-425.	0.7	23
15	A multicentric study regarding the use of hormone therapy during female mid-age (REDLINC VI). Climacteric, 2014, 17, 433-441.	1.1	22
16	Severe menopausal symptoms in mid-aged Latin American women can be related to their indigenous ethnic component. Climacteric, 2011, 14, 157-163.	1.1	21
17	Obesity is associated with a higher prevalence of musculoskeletal pain in middle-aged women. Gynecological Endocrinology, 2017, 33, 378-382.	0.7	14
18	Personal and professional use of menopausal hormone therapy among gynecologists: A multinational study (REDLINC VII). Maturitas, 2016, 87, 67-71.	1.0	13

#	Article	IF	CITATIONS
19	Evaluation of depressive symptoms in mid-aged women: report of a multicenter South American study. Menopause, 2017, 24, 1282-1288.	0.8	12
20	Climacteric symptoms in Quechua and Mestizo women from the Andean region of Cusco, Peru: Effects of altitude and ethnicity. Maturitas, 2014, 77, 356-360.	1.0	6
21	Does the menopausal status of female gynecologists affect their prescription of menopausal hormone therapy?. Climacteric, 2016, 19, 387-392.	1.1	5
22	Muscle health in Hispanic women. REDLINC VIII. Climacteric, 2020, 23, 184-191.	1.1	5
23	Association of bilateral oophorectomy and menopause hormone therapy with mild cognitive impairment: the REDLINC X study. Climacteric, 2022, 25, 195-202.	1.1	5