

Enzo Bonora

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/7695531/enzo-bonora-publications-by-year.pdf>

Version: 2023-12-10

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

299
papers

22,961
citations

74
h-index

146
g-index

305
ext. papers

25,558
ext. citations

7.7
avg, IF

6.56
L-index

#	Paper	IF	Citations
299	Role of monogenic diabetes genes on beta cell function in Italian patients with newly diagnosed type 2 diabetes. The Verona Newly Diagnosed Type 2 Diabetes Study (VNDS) 13.. <i>Diabetes and Metabolism</i> , 2022 , 101323	5.4	0
298	Growing up with type 1 diabetes mellitus: Data from the Verona Diabetes Transition Project. <i>Diabetic Medicine</i> , 2021 , e14719	3.5	1
297	Insulin resistance, diabetic kidney disease, and all-cause mortality in individuals with type 2 diabetes: a prospective cohort study. <i>BMC Medicine</i> , 2021 , 19, 66	11.4	9
296	Insulin-Mediated Substrate Use in Women With Different Phenotypes of PCOS: the Role of Androgens. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2021 , 106, e3414-e3425	5.6	2
295	Glomerular filtration rate decline in T2DM following diagnosis. The Verona newly diagnosed diabetes study-12. <i>Diabetes Research and Clinical Practice</i> , 2021 , 175, 108778	7.4	1
294	Transposition of cardiovascular outcome trial effects to the real-world population of patients with type 2 diabetes. <i>Cardiovascular Diabetology</i> , 2021 , 20, 103	8.7	1
293	Haplotypes of the genes (GCK and G6PC2) underlying the glucose/glucose-6-phosphate cycle are associated with pancreatic beta cell glucose sensitivity in patients with newly diagnosed type 2 diabetes from the VNDS study (VNDS 11). <i>Journal of Endocrinological Investigation</i> , 2021 , 44, 2567-2574	5.2	0
292	Effect of dulaglutide 3.0 and 4.5mg on weight in patients with type 2 diabetes: Exploratory analyses of AWARD-11. <i>Diabetes, Obesity and Metabolism</i> , 2021 , 23, 2242-2250	6.7	6
291	Incidence of diabetes mellitus in Italy in year 2018. A nationwide population-based study of the ARNO Diabetes Observatory. <i>Nutrition, Metabolism and Cardiovascular Diseases</i> , 2021 , 31, 2338-2344	4.5	2
290	Assessment of simple strategies for identifying undiagnosed diabetes and prediabetes in the general population. <i>Journal of Endocrinological Investigation</i> , 2021 , 44, 75-81	5.2	1
289	Independent association of atherogenic dyslipidaemia with all-cause mortality in individuals with type 2 diabetes and modifying effect of gender: a prospective cohort study. <i>Cardiovascular Diabetology</i> , 2021 , 20, 28	8.7	1
288	Common Variants Associated to Type 2 Diabetes in the Italian Population. <i>Open Journal of Endocrine and Metabolic Diseases</i> , 2021 , 11, 24-42	0.1	1
287	Efficacy and safety of dulaglutide 3.0 and 4.5 mg in patients aged younger than 65 and 65 years or older: Post hoc analysis of the AWARD-11 trial. <i>Diabetes, Obesity and Metabolism</i> , 2021 , 23, 2279-2288	6.7	2
286	SARS-CoV-2 and COVID-19 in diabetes mellitus. Population-based study on ascertained infections, hospital admissions and mortality in an Italian region with ~5 million inhabitants and ~250,000 diabetic people. <i>Nutrition, Metabolism and Cardiovascular Diseases</i> , 2021 , 31, 2612-2618	4.5	1
285	Initial treatment of diabetes in Italy. A nationwide population-based study from of the ARNO Diabetes Observatory. <i>Nutrition, Metabolism and Cardiovascular Diseases</i> , 2021 , 31, 2661-2668	4.5	0
284	Gastrointestinal Tolerability of Once-Weekly Dulaglutide 3.0 mg and 4.5 mg: A Post Hoc Analysis of the Incidence and Prevalence of Nausea, Vomiting, and Diarrhea in AWARD-11. <i>Diabetes Therapy</i> , 2021 , 12, 2783-2794	3.6	1
283	Glycaemic efficacy of an expanded dose range of dulaglutide according to baseline glycated haemoglobin (HbA1c) subgroup: Post hoc analysis of AWARD-11. <i>Diabetes, Obesity and Metabolism</i> , 2021 , 23, 2819-2824	6.7	

282	Time series of diabetes attributable mortality from 2008 to 2017. <i>Journal of Endocrinological Investigation</i> , 2021 , 1	5.2	2
281	Efficacy and Safety of Dulaglutide 3.0 mg and 4.5 mg Versus Dulaglutide 1.5 mg in Metformin-Treated Patients With Type 2 Diabetes in a Randomized Controlled Trial (AWARD-11). <i>Diabetes Care</i> , 2021 , 44, 765-773	14.6	36
280	Clinical Value Of Serum Levels Of 11-Oxygenated Metabolites Of Testosterone In Women With Polycystic Ovary Syndrome.. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2021 ,	5.6	2
279	Serum Androgens Are Independent Predictors of Insulin Clearance but Not of Insulin Secretion in Women With PCOS. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2020 , 105,	5.6	10
278	Association between On-Treatment Haemoglobin A and All-Cause Mortality in Individuals with Type 2 Diabetes: Importance of Personalized Goals and Type of Anti-Hyperglycaemic Treatment. <i>Journal of Clinical Medicine</i> , 2020 , 9,	5.1	1
277	Diabetes and NAFLD. <i>Endocrinology</i> , 2020 , 495-521	0.1	
276	Fifteen-Year Trends of Cardiogenic Shock and Mortality in Patients with Diabetes and Acute Coronary Syndromes. <i>American Journal of Medicine</i> , 2020 , 133, 331-339.e2	2.4	4
275	Screening for non-alcoholic fatty liver disease using liver stiffness measurement and its association with chronic kidney disease and cardiovascular complications in patients with type 2 diabetes. <i>Diabetes and Metabolism</i> , 2020 , 46, 296-303	5.4	21
274	A view on the quality of diabetes care in Italy and the role of Diabetes Clinics from the 2018 ARNO Diabetes Observatory. <i>Nutrition, Metabolism and Cardiovascular Diseases</i> , 2020 , 30, 1945-1953	4.5	10
273	Renal hyperfiltration is independently associated with increased all-cause mortality in individuals with type 2 diabetes: a prospective cohort study. <i>BMJ Open Diabetes Research and Care</i> , 2020 , 8,	4.5	5
272	Clinical burden of diabetes in Italy in 2018: a look at a systemic disease from the ARNO Diabetes Observatory. <i>BMJ Open Diabetes Research and Care</i> , 2020 , 8,	4.5	9
271	Chronic complications in patients with newly diagnosed type 2 diabetes: prevalence and related metabolic and clinical features: the Verona Newly Diagnosed Type 2 Diabetes Study (VNDS) 9. <i>BMJ Open Diabetes Research and Care</i> , 2020 , 8,	4.5	16
270	Depressive symptoms and glycaemic control in adults with type 1 diabetes: an exploratory study on the role of family functioning. <i>Acta Diabetologica</i> , 2020 , 57, 23-30	3.9	3
269	Continuous intraperitoneal insulin infusion: an alternative route for insulin delivery in type 1 diabetes. <i>Acta Diabetologica</i> , 2020 , 57, 101-104	3.9	1
268	Association Between Nonalcoholic Fatty Liver Disease and Reduced Bone Mineral Density in Children: A Meta-Analysis. <i>Hepatology</i> , 2019 , 70, 812-823	11.2	16
267	Cardiovascular Effects of Pioglitazone or Sulfonylureas According to Pretreatment Risk: Moving Toward Personalized Care. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2019 , 104, 3296-3302	5.6	8
266	Association between non-alcoholic fatty liver disease and decreased lung function in adults: A systematic review and meta-analysis. <i>Diabetes and Metabolism</i> , 2019 , 45, 536-544	5.4	17
265	Glucose-lowering therapy and cardiovascular outcomes in patients with type 2 diabetes mellitus and acute coronary syndrome. <i>Diabetes and Vascular Disease Research</i> , 2019 , 16, 399-414	3.3	13

264	Is resistant hypertension an independent predictor of all-cause mortality in individuals with type 2 diabetes? A prospective cohort study. <i>BMC Medicine</i> , 2019 , 17, 83	11.4	4
263	The use of real time continuous glucose monitoring or flash glucose monitoring in the management of diabetes: A consensus view of Italian diabetes experts using the Delphi method. <i>Nutrition, Metabolism and Cardiovascular Diseases</i> , 2019 , 29, 421-431	4.5	34
262	Loss of ZnT8 function protects against diabetes by enhanced insulin secretion. <i>Nature Genetics</i> , 2019 , 51, 1596-1606	36.3	45
261	Diabetes and NAFLD. <i>Endocrinology</i> , 2019 , 1-27	0.1	
260	Pasta Consumption and Connected Dietary Habits: Associations with Glucose Control, Adiposity Measures, and Cardiovascular Risk Factors in People with Type 2 Diabetes-TOSCA.IT Study. <i>Nutrients</i> , 2019 , 12,	6.7	8
259	Effects of Aerobic and Resistance Training on Circulating Micro-RNA Expression Profile in Subjects With Type 2 Diabetes. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2019 , 104, 1119-1130	5.6	13
258	Association between PNPLA3rs738409 polymorphism decreased kidney function in postmenopausal type 2 diabetic women with or without non-alcoholic fatty liver disease. <i>Diabetes and Metabolism</i> , 2019 , 45, 480-487	5.4	22
257	Ten years of experience with DPP-4 inhibitors for the treatment of type 2 diabetes mellitus. <i>Acta Diabetologica</i> , 2019 , 56, 605-617	3.9	29
256	Systematic review with meta-analysis: non-alcoholic fatty liver disease is associated with a history of osteoporotic fractures but not with low bone mineral density. <i>Alimentary Pharmacology and Therapeutics</i> , 2019 , 49, 375-388	6.1	22
255	Long-Acting GLP-1 Receptor Agonist Exenatide Influence on the Autonomic Cardiac Sympatho-Vagal Balance. <i>Journal of the Endocrine Society</i> , 2018 , 2, 53-62	0.4	4
254	Attending Diabetes Clinics is associated with a lower all-cause mortality. A meta-analysis of observational studies performed in Italy. <i>Nutrition, Metabolism and Cardiovascular Diseases</i> , 2018 , 28, 431-435	4.5	15
253	Haemoglobin A1c variability is a strong, independent predictor of all-cause mortality in patients with type 2 diabetes. <i>Diabetes, Obesity and Metabolism</i> , 2018 , 20, 1885-1893	6.7	27
252	Association of free-living physical activity measures with metabolic phenotypes in type 2 diabetes at the time of diagnosis. The Verona Newly Diagnosed Type 2 Diabetes Study (VNDS). <i>Nutrition, Metabolism and Cardiovascular Diseases</i> , 2018 , 28, 343-351	4.5	4
251	Mortality from infectious diseases in diabetes. <i>Nutrition, Metabolism and Cardiovascular Diseases</i> , 2018 , 28, 444-450	4.5	28
250	Nonalcoholic Fatty Liver Disease and Risk of Incident Type 2 Diabetes: A Meta-analysis. <i>Diabetes Care</i> , 2018 , 41, 372-382	14.6	262
249	Effects of aerobic or resistance exercise training on cardiovascular autonomic function of subjects with type 2 diabetes: A pilot study. <i>Nutrition, Metabolism and Cardiovascular Diseases</i> , 2018 , 28, 226-233	4.5	15
248	Sex differences in the association of psychological status with measures of physical activity and sedentary behaviour in adults with type 2 diabetes. <i>Acta Diabetologica</i> , 2018 , 55, 627-635	3.9	3
247	Diabetes and NAFLD. <i>Endocrinology</i> , 2018 , 1-27	0.1	

246	Defining the contribution of chronic kidney disease to all-cause mortality in patients with type 2 diabetes: the Renal Insufficiency And Cardiovascular Events (RIACE) Italian Multicenter Study. <i>Acta Diabetologica</i> , 2018 , 55, 603-612	3.9	23
245	Dietary intake and major food sources of polyphenols in people with type 2 diabetes: The TOSCA.IT Study. <i>European Journal of Nutrition</i> , 2018 , 57, 679-688	5.2	30
244	Hyperuricemia is associated with an increased prevalence of paroxysmal atrial fibrillation in patients with type 2 diabetes referred for clinically indicated 24-h Holter monitoring. <i>Journal of Endocrinological Investigation</i> , 2018 , 41, 223-231	5.2	14
243	Left ventricular chamber dilation and filling pressure may help to categorise patients with type 2 diabetes. <i>BMJ Open Diabetes Research and Care</i> , 2018 , 6, e000529	4.5	2
242	Non-albuminuric renal impairment is a strong predictor of mortality in individuals with type 2 diabetes: the Renal Insufficiency And Cardiovascular Events (RIACE) Italian multicentre study. <i>Diabetologia</i> , 2018 , 61, 2277-2289	10.3	52
241	Association Between Primary Hypothyroidism and Nonalcoholic Fatty Liver Disease: A Systematic Review and Meta-Analysis. <i>Thyroid</i> , 2018 , 28, 1270-1284	6.2	50
240	A renal genetic risk score (GRS) is associated with kidney dysfunction in people with type 2 diabetes. <i>Diabetes Research and Clinical Practice</i> , 2018 , 144, 137-143	7.4	
239	Impact of a Mediterranean Dietary Pattern and Its Components on Cardiovascular Risk Factors, Glucose Control, and Body Weight in People with Type 2 Diabetes: A Real-Life Study. <i>Nutrients</i> , 2018 , 10,	6.7	47
238	Diabetes and NAFLD. <i>Endocrinology</i> , 2018 , 495-521	0.1	
237	Nonalcoholic fatty liver disease increases risk of incident chronic kidney disease: A systematic review and meta-analysis. <i>Metabolism: Clinical and Experimental</i> , 2018 , 79, 64-76	12.7	171
236	The E/eR ratio difference between subjects with type 2 diabetes and controls. A meta-analysis of clinical studies. <i>PLoS ONE</i> , 2018 , 13, e0209794	3.7	4
235	Association between decreasing estimated glomerular filtration rate and risk of cardiac conduction defects in patients with type 2 diabetes. <i>Diabetes and Metabolism</i> , 2018 , 44, 473-481	5.4	1
234	Association between nonalcoholic fatty liver disease and colorectal tumours in asymptomatic adults undergoing screening colonoscopy: a systematic review and meta-analysis. <i>Metabolism: Clinical and Experimental</i> , 2018 , 87, 1-12	12.7	48
233	Psychological distress, self-efficacy and glycemic control in type 2 diabetes. <i>Nutrition, Metabolism and Cardiovascular Diseases</i> , 2017 , 27, 300-306	4.5	30
232	Nonalcoholic fatty liver disease is associated with an increased prevalence of distal symmetric polyneuropathy in adult patients with type 1 diabetes. <i>Journal of Diabetes and Its Complications</i> , 2017 , 31, 1021-1026	3.2	30
231	Association between subclinical left ventricular systolic dysfunction and glycemic control in asymptomatic type 2 diabetic patients with preserved left ventricular function. <i>Journal of Diabetes and Its Complications</i> , 2017 , 31, 1035-1040	3.2	6
230	Circulating MicroRNA-122 Is Associated With the Risk of New-Onset Metabolic Syndrome and Type 2 Diabetes. <i>Diabetes</i> , 2017 , 66, 347-357	0.9	141
229	Mechanisms linking empagliflozin to cardiovascular and renal protection. <i>International Journal of Cardiology</i> , 2017 , 241, 450-456	3.2	24

228	Association Between Vascular Cell Adhesion Molecule 1 and Atrial Fibrillation. <i>JAMA Cardiology</i> , 2017 , 2, 516-523	16.2	28
227	Influence of health locus of control and fear of hypoglycaemia on glycaemic control and treatment satisfaction in people with Type 1 diabetes on insulin pump therapy. <i>Diabetic Medicine</i> , 2017 , 34, 691-697	7.5	15
226	Relation of elevated serum uric acid levels to first-degree heart block and other cardiac conduction defects in hospitalized patients with type 2 diabetes. <i>Journal of Diabetes and Its Complications</i> , 2017 , 31, 1691-1697	3.2	5
225	Prognostic Impact of Diabetes on Long-term Survival Outcomes in Patients With Heart Failure: A Meta-analysis. <i>Diabetes Care</i> , 2017 , 40, 1597-1605	14.6	52
224	Insulin resistance in a large cohort of women with polycystic ovary syndrome: a comparison between euglycaemic-hyperinsulinaemic clamp and surrogate indexes. <i>Human Reproduction</i> , 2017 , 32, 2515-2521	5.7	43
223	Rationale and design of the DARWIN-T2D (Dapagliflozin Real World Evidence in Type 2 Diabetes): A multicenter retrospective nationwide Italian study and crowdsourcing opportunity. <i>Nutrition, Metabolism and Cardiovascular Diseases</i> , 2017 , 27, 1089-1097	4.5	19
222	Effects on the incidence of cardiovascular events of the addition of pioglitazone versus sulfonylureas in patients with type 2 diabetes inadequately controlled with metformin (TOSCA.IT): a randomised, multicentre trial. <i>Lancet Diabetes and Endocrinology</i> , 2017 , 5, 887-897	18.1	154
221	Mitral Regurgitation and Increased Risk of All-Cause and Cardiovascular Mortality in Patients with Type 2 Diabetes. <i>American Journal of Medicine</i> , 2017 , 130, 70-76.e1	2.4	15
220	Polyphenol intake and cardiovascular risk factors in a population with type 2 diabetes: The TOSCA.IT study. <i>Clinical Nutrition</i> , 2017 , 36, 1686-1692	5.9	42
219	Nonalcoholic fatty liver disease is associated with an increased risk of heart block in hospitalized patients with type 2 diabetes mellitus. <i>PLoS ONE</i> , 2017 , 12, e0185459	3.7	29
218	Influence of dietary fat and carbohydrates proportions on plasma lipids, glucose control and low-grade inflammation in patients with type 2 diabetes-The TOSCA.IT Study. <i>European Journal of Nutrition</i> , 2016 , 55, 1645-51	5.2	32
217	Hyperuricemia is associated with an increased prevalence of atrial fibrillation in hospitalized patients with type 2 diabetes. <i>Journal of Endocrinological Investigation</i> , 2016 , 39, 159-67	5.2	25
216	A Novel Insulin/Glucose Model after a Mixed-Meal Test in Patients with Type 1 Diabetes on Insulin Pump Therapy. <i>Scientific Reports</i> , 2016 , 6, 36029	4.9	8
215	Nonalcoholic fatty liver disease is independently associated with an increased incidence of cardiovascular disease in adult patients with type 1 diabetes. <i>International Journal of Cardiology</i> , 2016 , 225, 387-391	3.2	39
214	Inadequate hepcidin serum concentrations predict incident type 2 diabetes mellitus. <i>Diabetes/Metabolism Research and Reviews</i> , 2016 , 32, 187-92	7.5	15
213	Evidence of left atrial remodeling and left ventricular diastolic dysfunction in type 2 diabetes mellitus with preserved systolic function. <i>Nutrition, Metabolism and Cardiovascular Diseases</i> , 2016 , 26, 1026-1032	4.5	11
212	DPP-4 inhibitors and cardiovascular disease in type 2 diabetes mellitus. Expectations, observations and perspectives. <i>Nutrition, Metabolism and Cardiovascular Diseases</i> , 2016 , 26, 273-84	4.5	16
211	Implications of Androgen Assay Accuracy in the Phenotyping of Women With Polycystic Ovary Syndrome. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2016 , 101, 610-8	5.6	37

210	Interleukin-6 as a potential positive modulator of human beta-cell function: an exploratory analysis-the Verona Newly Diagnosed Type 2 Diabetes Study (VNDS) 6. <i>Acta Diabetologica</i> , 2016 , 53, 393-402	3.9	3
209	A post hoc analysis of saxagliptin efficacy and safety in patients with type 2 diabetes stratified by UKPDS 10-year cardiovascular risk score. <i>Nutrition, Metabolism and Cardiovascular Diseases</i> , 2016 , 26, 374-9	4.5	1
208	Nonalcoholic Fatty Liver Disease Is Associated With Ventricular Arrhythmias in Patients With Type 2 Diabetes Referred for Clinically Indicated 24-Hour Holter Monitoring. <i>Diabetes Care</i> , 2016 , 39, 1416-23	14.6	66
207	The relative burden of diabetes complications on healthcare costs: The population-based CINECA-SID ARNO Diabetes Observatory. <i>Nutrition, Metabolism and Cardiovascular Diseases</i> , 2016 , 26, 944-50	4.5	35
206	Is common genetic variation at IRS1, ENPP1 and TRIB3 loci associated with cardiometabolic phenotypes in type 2 diabetes? An exploratory analysis of the Verona Newly Diagnosed Type 2 Diabetes Study (VNDS) 5. <i>Nutrition, Metabolism and Cardiovascular Diseases</i> , 2016 , 26, 232-8	4.5	5
205	Sex differences in food choices, adherence to dietary recommendations and plasma lipid profile in type 2 diabetes - The TOSCA.IT study. <i>Nutrition, Metabolism and Cardiovascular Diseases</i> , 2016 , 26, 879-85	4.5	30
204	Incidence, prevalence, costs and quality of care of type 1 diabetes in Italy, age 0-29 years: The population-based CINECA-SID ARNO Observatory, 2002-2012. <i>Nutrition, Metabolism and Cardiovascular Diseases</i> , 2016 , 26, 1104-1111	4.5	11
203	Prevalence of neuropathy in type 2 diabetic patients and its association with other diabetes complications: The Verona Diabetic Foot Screening Program. <i>Journal of Diabetes and Its Complications</i> , 2015 , 29, 1066-70	3.2	53
202	Relationship between increased left atrial volume and microvascular complications in patients with type 2 diabetes. <i>Journal of Diabetes and Its Complications</i> , 2015 , 29, 822-8	3.2	9
201	Prevalence of Cardiovascular Autonomic Neuropathy in a Cohort of Patients With Newly Diagnosed Type 2 Diabetes: The Verona Newly Diagnosed Type 2 Diabetes Study (VNDS). <i>Diabetes Care</i> , 2015 , 38, 1487-93	14.6	42
200	Clinical factors that predict remission of diabetes after different bariatric surgical procedures: interdisciplinary group of bariatric surgery of Verona (G.I.C.O.V.). <i>Acta Diabetologica</i> , 2015 , 52, 937-42	3.9	8
199	Association of a 62 Variants Type 2 Diabetes Genetic Risk Score With Markers of Subclinical Atherosclerosis: A Transethnic, Multicenter Study. <i>Circulation: Cardiovascular Genetics</i> , 2015 , 8, 507-15		11
198	Serum testosterone predicts cardiorespiratory fitness impairment in normal-weight women with polycystic ovary syndrome. <i>Clinical Endocrinology</i> , 2015 , 83, 895-901	3.4	4
197	Nonalcoholic Fatty Liver Disease Is Independently Associated with Early Left Ventricular Diastolic Dysfunction in Patients with Type 2 Diabetes. <i>PLoS ONE</i> , 2015 , 10, e0135329	3.7	61
196	Heart valve calcification in patients with type 2 diabetes and nonalcoholic fatty liver disease. <i>Metabolism: Clinical and Experimental</i> , 2015 , 64, 879-87	12.7	55
195	Lower levels of 25-hydroxyvitamin D3 are associated with a higher prevalence of microvascular complications in patients with type 2 diabetes. <i>BMJ Open Diabetes Research and Care</i> , 2015 , 3, e000058	4.5	32
194	Total body fat and central fat mass independently predict insulin resistance but not hyperandrogenemia in women with polycystic ovary syndrome. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2015 , 100, 661-9	5.6	42
193	Usefulness of subclinical left ventricular midwall dysfunction to predict cardiovascular mortality in patients with type 2 diabetes mellitus. <i>American Journal of Cardiology</i> , 2014 , 113, 1409-14	3	20

192	Chronic kidney disease in type 2 diabetes: lessons from the Renal Insufficiency And Cardiovascular Events (RIACE) Italian Multicentre Study. <i>Nutrition, Metabolism and Cardiovascular Diseases</i> , 2014 , 24, 815-22	4.5	41
191	One size does not fit all glycemic targets for type 2 diabetes. <i>Journal of Diabetes Investigation</i> , 2014 , 5, 134-41	3.9	21
190	Nonalcoholic fatty liver disease is independently associated with an increased incidence of chronic kidney disease in patients with type 1 diabetes. <i>Diabetes Care</i> , 2014 , 37, 1729-36	14.6	98
189	Plasma levels of pentraxin-3, an inflammatory protein involved in fertility, are reduced in women with polycystic ovary syndrome. <i>European Journal of Endocrinology</i> , 2014 , 170, 401-9	6.5	15
188	Mortality from chronic liver diseases in diabetes. <i>American Journal of Gastroenterology</i> , 2014 , 109, 1020-5.7		83
187	Cardiovascular outcome trials of glucose-lowering strategies in type 2 diabetes. <i>Lancet, The</i> , 2014 , 384, 1096	4.0	2
186	Relationships between cardiorespiratory fitness, metabolic control, and fat distribution in type 2 diabetes subjects. <i>Acta Diabetologica</i> , 2014 , 51, 369-75	3.9	12
185	Under-treatment of migrants with diabetes in a universalistic health care system: the ARNO Observatory. <i>Nutrition, Metabolism and Cardiovascular Diseases</i> , 2014 , 24, 393-9	4.5	21
184	Association between plasma triglycerides and high-density lipoprotein cholesterol and microvascular kidney disease and retinopathy in type 2 diabetes mellitus: a global case-control study in 13 countries. <i>Circulation</i> , 2014 , 129, 999-1008	16.7	153
183	Hemostatic and fibrinolytic abnormalities in polycystic ovary syndrome. <i>Seminars in Thrombosis and Hemostasis</i> , 2014 , 40, 600-18	5.3	13
182	Antidromic sensory nerve conduction study of the digital branches of the medial plantar nerve: a novel method to detect early diabetic sensory axonal polyneuropathy. <i>Muscle and Nerve</i> , 2014 , 50, 193-9 ³⁻⁴		7
181	Haptoglobin 2-2 genotype is not associated with cardiovascular risk in subjects with elevated glycohemoglobin-results from the Bruneck Study. <i>Journal of the American Heart Association</i> , 2014 , 3, e000732	6	21
180	Distribution of cardiovascular disease and retinopathy in patients with type 2 diabetes according to different classification systems for chronic kidney disease: a cross-sectional analysis of the renal insufficiency and cardiovascular events (RIACE) Italian multicenter study. <i>Cardiovascular Diabetology</i> , 2014 , 13, 59	8.7	22
179	Association of nonalcoholic fatty liver disease with QTc interval in patients with type 2 diabetes. <i>Nutrition, Metabolism and Cardiovascular Diseases</i> , 2014 , 24, 663-9	4.5	55
178	HELP LDL apheresis reduces plasma pentraxin 3 in familial hypercholesterolemia. <i>PLoS ONE</i> , 2014 , 9, e101290	3.7	16
177	Six cases with severe insulin resistance (SIR) associated with mutations of insulin receptor: Is a Bartter-like syndrome a feature of congenital SIR?. <i>Acta Diabetologica</i> , 2013 , 50, 951-7	3.9	28
176	PPARG2 Pro12Ala and ADAMTS9 rs4607103 as "insulin resistance loci" and "insulin secretion loci" in Italian individuals. The GENFIEV study and the Verona Newly Diagnosed Type 2 Diabetes Study (VNDS) 4. <i>Acta Diabetologica</i> , 2013 , 50, 401-8	3.9	26
175	Both resistance training and aerobic training reduce hepatic fat content in type 2 diabetic subjects with nonalcoholic fatty liver disease (the RAED2 Randomized Trial). <i>Hepatology</i> , 2013 , 58, 1287-95	11.2	211

174	Type 2 diabetes mellitus: a disease of the governance of the glucose-insulin system: an experimental metabolic control analysis study. <i>Nutrition, Metabolism and Cardiovascular Diseases</i> , 2013 , 23, 23-30	4.5	5
173	HbA1c variability as an independent correlate of nephropathy, but not retinopathy, in patients with type 2 diabetes: the Renal Insufficiency And Cardiovascular Events (RIACE) Italian multicenter study. <i>Diabetes Care</i> , 2013 , 36, 2301-10	14.6	113
172	Gender differences in cardiovascular disease risk factors, treatments and complications in patients with type 2 diabetes: the RIACE Italian multicentre study. <i>Journal of Internal Medicine</i> , 2013 , 274, 176-91	10.8	90
171	Inappropriate left ventricular mass independently predicts cardiovascular mortality in patients with type 2 diabetes. <i>International Journal of Cardiology</i> , 2013 , 168, 4953-6	3.2	13
170	Divergences in insulin resistance between the different phenotypes of the polycystic ovary syndrome. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2013 , 98, E628-37	5.6	140
169	Blockade of receptor activator of nuclear factor- κ B (RANKL) signaling improves hepatic insulin resistance and prevents development of diabetes mellitus. <i>Nature Medicine</i> , 2013 , 19, 358-63	50.5	169
168	Metabolic inflexibility is a feature of women with polycystic ovary syndrome and is associated with both insulin resistance and hyperandrogenism. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2013 , 98, 2581-8	5.6	30
167	Non-alcoholic fatty liver disease is associated with an increased prevalence of atrial fibrillation in hospitalized patients with type 2 diabetes. <i>Clinical Science</i> , 2013 , 125, 301-9	6.5	80
166	Age, renal dysfunction, cardiovascular disease, and antihyperglycemic treatment in type 2 diabetes mellitus: findings from the Renal Insufficiency and Cardiovascular Events Italian Multicenter Study. <i>Journal of the American Geriatrics Society</i> , 2013 , 61, 1253-61	5.6	51
165	Glycated haemoglobin is inversely related to serum vitamin D levels in type 2 diabetic patients. <i>PLoS ONE</i> , 2013 , 8, e82733	3.7	36
164	Triglyceride-high-density lipoprotein cholesterol is associated with microvascular complications in type 2 diabetes mellitus. <i>Metabolism: Clinical and Experimental</i> , 2012 , 61, 22-9	12.7	45
163	Serum uric acid levels and incident chronic kidney disease in patients with type 2 diabetes and preserved kidney function. <i>Diabetes Care</i> , 2012 , 35, 99-104	14.6	167
162	Non-alcoholic fatty liver disease is independently associated with left ventricular hypertrophy in hypertensive Type 2 diabetic individuals. <i>Journal of Endocrinological Investigation</i> , 2012 , 35, 215-8	5.2	44
161	Increased risk of cardiovascular disease and chronic kidney disease in NAFLD. <i>Nature Reviews Gastroenterology and Hepatology</i> , 2012 , 9, 372-81	24.2	90
160	Aortic and mitral annular calcifications are predictive of all-cause and cardiovascular mortality in patients with type 2 diabetes. <i>Diabetes Care</i> , 2012 , 35, 1781-6	14.6	52
159	Addition of either pioglitazone or a sulfonylurea in type 2 diabetic patients inadequately controlled with metformin alone: impact on cardiovascular events. A randomized controlled trial. <i>Nutrition, Metabolism and Cardiovascular Diseases</i> , 2012 , 22, 997-1006	4.5	36
158	CACNA1E variants affect beta cell function in patients with newly diagnosed type 2 diabetes. the Verona newly diagnosed type 2 diabetes study (VNDS) 3. <i>PLoS ONE</i> , 2012 , 7, e32755	3.7	20
157	Differences in the acute effects of aerobic and resistance exercise in subjects with type 2 diabetes: results from the RAED2 Randomized Trial. <i>PLoS ONE</i> , 2012 , 7, e49937	3.7	28

156	Increased prevalence of chronic kidney disease in patients with Type 1 diabetes and non-alcoholic fatty liver. <i>Diabetic Medicine</i> , 2012 , 29, 220-6	3.5	48
155	The TOSCA.IT trial: a study designed to evaluate the effect of pioglitazone versus sulfonylureas on cardiovascular disease in type 2 diabetes. <i>Diabetes Care</i> , 2012 , 35, e82	14.6	23
154	Diverging association of reduced glomerular filtration rate and albuminuria with coronary and noncoronary events in patients with type 2 diabetes: the renal insufficiency and cardiovascular events (RIACE) Italian multicenter study. <i>Diabetes Care</i> , 2012 , 35, 143-9	14.6	91
153	Quality of diabetes care in Italy: information from a large population-based multiregional observatory (ARNO diabetes). <i>Diabetes Care</i> , 2012 , 35, e64	14.6	14
152	Predictors of estimated GFR decline in patients with type 2 diabetes and preserved kidney function. <i>Clinical Journal of the American Society of Nephrology: CJASN</i> , 2012 , 7, 401-8	6.9	138
151	Hyperinsulinemia amplifies GnRH agonist stimulated ovarian steroid secretion in women with polycystic ovary syndrome. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2012 , 97, 1712-9	5.6	56
150	Comparison of two creatinine-based estimating equations in predicting all-cause and cardiovascular mortality in patients with type 2 diabetes. <i>Diabetes Care</i> , 2012 , 35, 2347-53	14.6	20
149	Metabolic effects of aerobic training and resistance training in type 2 diabetic subjects: a randomized controlled trial (the RAED2 study). <i>Diabetes Care</i> , 2012 , 35, 676-82	14.6	123
148	Increased prevalence of cardiovascular disease in Type 1 diabetic patients with non-alcoholic fatty liver disease. <i>Journal of Endocrinological Investigation</i> , 2012 , 35, 535-40	5.2	19
147	Risk of chronic kidney disease in patients with non-alcoholic fatty liver disease: is there a link?. <i>Journal of Hepatology</i> , 2011 , 54, 1020-9	13.4	131
146	The Chronic Kidney Disease Epidemiology Collaboration (CKD-EPI) equation provides a better definition of cardiovascular burden associated with CKD than the Modification of Diet in Renal Disease (MDRD) Study formula in subjects with type 2 diabetes. <i>Atherosclerosis</i> , 2011 , 218, 194-9	3.1	43
145	Clinical significance of nonalbuminuric renal impairment in type 2 diabetes. <i>Journal of Hypertension</i> , 2011 , 29, 1802-9	1.9	147
144	The use of HbA(1c) for the diagnosis of diabetes. <i>Current Diabetes Reports</i> , 2011 , 11, 470-1	5.6	
143	Glomerular filtration rate, albuminuria and risk of cardiovascular and all-cause mortality in type 2 diabetic individuals. <i>Nutrition, Metabolism and Cardiovascular Diseases</i> , 2011 , 21, 294-301	4.5	23
142	The role of serum uric acid in cardiovascular disease in type 2 diabetic and non-diabetic subjects: a narrative review. <i>Journal of Endocrinological Investigation</i> , 2011 , 34, 881-6	5.2	22
141	Variants and haplotypes of TCF7L2 are associated with β cell function in patients with newly diagnosed type 2 diabetes: the Verona Newly Diagnosed Type 2 Diabetes Study (VNDS) 1. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2011 , 96, E389-93	5.6	32
140	Variants of GCKR affect both β cell and kidney function in patients with newly diagnosed type 2 diabetes: the Verona newly diagnosed type 2 diabetes study 2. <i>Diabetes Care</i> , 2011 , 34, 1205-10	14.6	27
139	Impact of reference category and number of traits in the cluster on risk of coronary heart disease in metabolic syndrome: prospective data from the Bruneck study. <i>Metabolic Syndrome and Related Disorders</i> , 2011 , 9, 313-8	2.6	1

138	Association of variation at the ABO locus with circulating levels of soluble intercellular adhesion molecule-1, soluble P-selectin, and soluble E-selectin: a meta-analysis. <i>Circulation: Cardiovascular Genetics</i> , 2011 , 4, 681-6		59
137	Insulin enhances ACTH-stimulated androgen and glucocorticoid metabolism in hyperandrogenic women. <i>European Journal of Endocrinology</i> , 2011 , 164, 197-203	6.5	31
136	High-normal HbA1c is a strong predictor of type 2 diabetes in the general population. <i>Diabetes Care</i> , 2011 , 34, 1038-40	14.6	39
135	The pros and cons of diagnosing diabetes with A1C. <i>Diabetes Care</i> , 2011 , 34 Suppl 2, S184-90	14.6	231
134	Supervised walking groups to increase physical activity in type 2 diabetic patients. <i>Diabetes Care</i> , 2010 , 33, 2333-5	14.6	34
133	Risk of cardiovascular disease in patients with nonalcoholic fatty liver disease. <i>New England Journal of Medicine</i> , 2010 , 363, 1341-50	59.2	1353
132	Terapia del diabete di tipo 2: obiettivi glicemici in relazione al tipo di paziente. <i>L Endocrinologo</i> , 2010 , 11, 254-259	0	
131	Prevalence of non-alcoholic fatty liver disease and its association with cardiovascular disease in patients with type 1 diabetes. <i>Journal of Hepatology</i> , 2010 , 53, 713-8	13.4	171
130	Anaemia, independent of chronic kidney disease, predicts all-cause and cardiovascular mortality in type 2 diabetic patients. <i>Atherosclerosis</i> , 2010 , 210, 575-80	3.1	26
129	Usefulness of the triglyceride to high-density lipoprotein cholesterol ratio for predicting mortality risk in type 2 diabetes: role of kidney dysfunction. <i>Atherosclerosis</i> , 2010 , 212, 287-91	3.1	14
128	Plasma microRNA profiling reveals loss of endothelial miR-126 and other microRNAs in type 2 diabetes. <i>Circulation Research</i> , 2010 , 107, 810-7	15.7	1086
127	Non-alcoholic fatty liver disease is independently associated with an increased prevalence of chronic kidney disease and retinopathy in type 1 diabetic patients. <i>Diabetologia</i> , 2010 , 53, 1341-8	10.3	115
126	Endogenous hyperinsulinaemia in insulinoma patients is not associated with changes in beta-cell area and turnover in the tumor-adjacent pancreas. <i>Regulatory Peptides</i> , 2010 , 165, 180-5		3
125	Elevated serum uric acid concentrations independently predict cardiovascular mortality in type 2 diabetic patients. <i>Diabetes Care</i> , 2009 , 32, 1716-20	14.6	94
124	Hemostatic and fibrinolytic abnormalities in endocrine diseases: a narrative review. <i>Seminars in Thrombosis and Hemostasis</i> , 2009 , 35, 605-12	5.3	10
123	Vascular cell adhesion molecule 1 as a predictor of severe osteoarthritis of the hip and knee joints. <i>Arthritis and Rheumatism</i> , 2009 , 60, 2381-9		62
122	Is fasting glucose variability a risk factor for retinopathy in people with type 2 diabetes?. <i>Nutrition, Metabolism and Cardiovascular Diseases</i> , 2009 , 19, 334-9	4.5	33
121	Hyperinsulinemia and insulin resistance are independently associated with plasma lipids, uric acid and blood pressure in non-diabetic subjects. The GISIR database. <i>Nutrition, Metabolism and Cardiovascular Diseases</i> , 2008 , 18, 624-31	4.5	59

120	Protection of pancreatic beta-cells: is it feasible?. <i>Nutrition, Metabolism and Cardiovascular Diseases</i> , 2008 , 18, 74-83	4.5	62
119	Insulin effect on serum potassium and auto-inhibition of insulin secretion is intact in a patient with leprechaunism despite severe impairment of substrates metabolism. <i>Diabetes/Metabolism Research and Reviews</i> , 2008 , 24, 205-10	7.5	0
118	Impact of lowering the criterion for impaired fasting glucose on identification of individuals with insulin resistance. The GISIR database. <i>Diabetes/Metabolism Research and Reviews</i> , 2008 , 24, 130-6	7.5	5
117	Variability of body weight, pulse pressure and glycaemia strongly predict total mortality in elderly type 2 diabetic patients. The Verona Diabetes Study. <i>Diabetes/Metabolism Research and Reviews</i> , 2008 , 24, 624-8	7.5	50
116	European guidelines on cardiovascular disease prevention in clinical practice: executive summary: Fourth Joint Task Force of the European Society of Cardiology and Other Societies on Cardiovascular Disease Prevention in Clinical Practice (Constituted by representatives of nine societies and by invited experts). <i>European Heart Journal</i> , 2007 , 28, 2375-414	9.5	1029
115	The metabolic syndrome: Is it still alive?. <i>International Congress Series</i> , 2007 , 1303, 3-9		
114	Antidiabetic medications in overweight/obese patients with type 2 diabetes: drawbacks of current drugs and potential advantages of incretin-based treatment on body weight. <i>International Journal of Clinical Practice</i> , 2007 , 61, 19-28	2.9	28
113	Disruptive mitochondrial DNA mutations in complex I subunits are markers of oncocytic phenotype in thyroid tumors. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2007 , 104, 9001-6	11.5	217
112	Insulin resistance as estimated by homeostasis model assessment predicts incident symptomatic cardiovascular disease in caucasian subjects from the general population: the Bruneck study. <i>Diabetes Care</i> , 2007 , 30, 318-24	14.6	230
111	Correspondence between the International Diabetes Federation criteria for metabolic syndrome and insulin resistance in a cohort of Italian nondiabetic Caucasians: the GISIR database. <i>Diabetes Care</i> , 2007 , 30, e33	14.6	3
110	Pulse pressure and mortality from cerebrovascular diseases in type 2 diabetic patients: the Verona Diabetes Study. <i>Cerebrovascular Diseases</i> , 2007 , 23, 20-6	3.2	7
109	Prevalence and correlates of post-prandial hyperglycaemia in a large sample of patients with type 2 diabetes mellitus. <i>Diabetologia</i> , 2006 , 49, 846-54	10.3	100
108	The metabolic syndrome and cardiovascular disease. <i>Annals of Medicine</i> , 2006 , 38, 64-80	1.5	139
107	Imatinib and regression of type 2 diabetes. <i>New England Journal of Medicine</i> , 2005 , 352, 1049-50	59.2	101
106	Analysis of IMGSAC autism susceptibility loci: evidence for sex limited and parent of origin specific effects. <i>Journal of Medical Genetics</i> , 2005 , 42, 132-7	5.8	94
105	Population-based incidence rates and risk factors for type 2 diabetes in white individuals: the Bruneck study. <i>Diabetes</i> , 2004 , 53, 1782-9	0.9	214
104	The Metabolic Syndrome is an independent predictor of cardiovascular disease in Type 2 diabetic subjects. Prospective data from the Verona Diabetes Complications Study. <i>Diabetic Medicine</i> , 2004 , 21, 52-8	3.5	213
103	Serum leptin level and the risk of nontraumatic fracture. <i>American Journal of Medicine</i> , 2004 , 117, 952-6	2.4	27

102	Mortality from site-specific malignancies in type 2 diabetic patients from Verona. <i>Diabetes Care</i> , 2003 , 26, 1047-51	14.6	118
101	Metabolic syndrome: epidemiology and more extensive phenotypic description. Cross-sectional data from the Bruneck Study. <i>International Journal of Obesity</i> , 2003 , 27, 1283-9	5.5	127
100	Body mass index and the risk of mortality in type II diabetic patients from Verona. <i>International Journal of Obesity</i> , 2003 , 27, 281-5	5.5	56
99	Analysis of reelin as a candidate gene for autism. <i>Molecular Psychiatry</i> , 2003 , 8, 885-92	15.1	108
98	Screening of nine candidate genes for autism on chromosome 2q reveals rare nonsynonymous variants in the cAMP-GEFII gene. <i>Molecular Psychiatry</i> , 2003 , 8, 916-24	15.1	94
97	Carotid atherosclerosis and coronary heart disease in the metabolic syndrome: prospective data from the Bruneck study. <i>Diabetes Care</i> , 2003 , 26, 1251-7	14.6	246
96	Altered homeostatic adaptation of first- and second-phase beta-cell secretion in the offspring of patients with type 2 diabetes: studies with a minimal model to assess beta-cell function. <i>Diabetes</i> , 2003 , 52, 470-80	0.9	64
95	Predictors of insulin sensitivity in Type 2 diabetes mellitus. <i>Diabetic Medicine</i> , 2002 , 19, 535-42	3.5	42
94	Insulin causes endothelial dysfunction in humans: sites and mechanisms. <i>Circulation</i> , 2002 , 105, 576-82	16.7	319
93	HOMA-estimated insulin resistance is an independent predictor of cardiovascular disease in type 2 diabetic subjects: prospective data from the Verona Diabetes Complications Study. <i>Diabetes Care</i> , 2002 , 25, 1135-41	14.6	390
92	Toll-like receptor 4 polymorphisms and atherogenesis. <i>New England Journal of Medicine</i> , 2002 , 347, 185-92	9.2	912
91	FOXP2 is not a major susceptibility gene for autism or specific language impairment. <i>American Journal of Human Genetics</i> , 2002 , 70, 1318-27	11	173
90	Postprandial peaks as a risk factor for cardiovascular disease: epidemiological perspectives. <i>International Journal of Clinical Practice, Supplement</i> , 2002 , 5-11		28
89	Assessment of beta-cell function during the oral glucose tolerance test by a minimal model of insulin secretion. <i>European Journal of Clinical Investigation</i> , 2001 , 31, 405-16	4.6	76
88	Postprandial blood glucose as a risk factor for cardiovascular disease in Type II diabetes: the epidemiological evidence. <i>Diabetologia</i> , 2001 , 44, 2107-14	10.3	339
87	ACE genotype and endothelium-dependent vasodilation of conduit arteries and forearm microcirculation in humans. <i>Arteriosclerosis, Thrombosis, and Vascular Biology</i> , 2001 , 21, 1313-9	9.4	17
86	Chronic infections and the risk of carotid atherosclerosis: prospective results from a large population study. <i>Circulation</i> , 2001 , 103, 1064-70	16.7	432
85	Glucose instability is associated with a high level of circulating p-selectin. <i>Diabetes Care</i> , 2001 , 24, 1685	14.6	9

84	Relation between soluble adhesion molecules and insulin sensitivity in type 2 diabetic individuals: role of adipose tissue. <i>Diabetes Care</i> , 2001 , 24, 1961-6	14.6	46
83	Intracellular partition of plasma glucose disposal in hypertensive and normotensive subjects with type 2 diabetes mellitus. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2001 , 86, 2073-9	5.6	8
82	Plasma glucose levels throughout the day and HbA(1c) interrelationships in type 2 diabetes: implications for treatment and monitoring of metabolic control. <i>Diabetes Care</i> , 2001 , 24, 2023-9	14.6	186
81	Nuovi criteri diagnostici e classificativi del diabete mellito. <i>L Endocrinologo</i> , 2001 , 2, 42-47	0	
80	Intracellular Partition of Plasma Glucose Disposal in Hypertensive and Normotensive Subjects with Type 2 Diabetes Mellitus. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2001 , 86, 2073-2079	5.6	7
79	Relationship between regional fat distribution and insulin resistance. <i>International Journal of Obesity</i> , 2000 , 24 Suppl 2, S32-5	5.5	64
78	Impaired glucose tolerance, Type II diabetes mellitus and carotid atherosclerosis: prospective results from the Bruneck Study. <i>Diabetologia</i> , 2000 , 43, 156-64	10.3	106
77	Homeostasis model assessment closely mirrors the glucose clamp technique in the assessment of insulin sensitivity: studies in subjects with various degrees of glucose tolerance and insulin sensitivity. <i>Diabetes Care</i> , 2000 , 23, 57-63	14.6	1906
76	Fasting plasma glucose variability predicts 10-year survival of type 2 diabetic patients: the Verona Diabetes Study. <i>Diabetes Care</i> , 2000 , 23, 45-50	14.6	205
75	No association between dehydroepiandrosterone sulfate and development of atherosclerosis in a prospective population study (Bruneck Study). <i>Arteriosclerosis, Thrombosis, and Vascular Biology</i> , 2000 , 20, 1094-100	9.4	56
74	Distinct risk profiles of early and advanced atherosclerosis: prospective results from the Bruneck Study. <i>Arteriosclerosis, Thrombosis, and Vascular Biology</i> , 2000 , 20, 529-37	9.4	123
73	Effects of glucosamine infusion on insulin secretion and insulin action in humans. <i>Diabetes</i> , 2000 , 49, 926-35	0.9	122
72	Effect of chronic treatment with lacidipine or lisinopril on intracellular partitioning of glucose metabolism in type 2 diabetes mellitus. <i>Journal of Clinical Endocrinology and Metabolism</i> , 1999 , 84, 1544-50	5.6	11
71	Plasma glucose within the normal range is not associated with carotid atherosclerosis: prospective results in subjects with normal glucose tolerance from the Bruneck Study. <i>Diabetes Care</i> , 1999 , 22, 1339-46	14.6	25
70	Attending the diabetes center is associated with reduced cardiovascular mortality in type 2 diabetic patients: the Verona Diabetes Study. <i>Diabetes/Metabolism Research and Reviews</i> , 1999 , 15, 170-4	7.5	13
69	Cause-specific mortality in type 2 diabetes. The Verona Diabetes Study. <i>Diabetes Care</i> , 1999 , 22, 756-61	14.6	303
68	Effect of Chronic Treatment with Lacidipine or Lisinopril on Intracellular Partitioning of Glucose Metabolism in Type 2 Diabetes Mellitus. <i>Journal of Clinical Endocrinology and Metabolism</i> , 1999 , 84, 1544-1550	5.6	9
67	Role of tissue-specific blood flow and tissue recruitment in insulin-mediated glucose uptake of human skeletal muscle. <i>Circulation</i> , 1998 , 98, 234-41	16.7	132

66	Prevalence of insulin resistance in metabolic disorders: the Bruneck Study. <i>Diabetes</i> , 1998 , 47, 1643-9	0.9	647
65	U-shaped and J-shaped relationships between serum insulin and coronary heart disease in the general population. The Bruneck Study. <i>Diabetes Care</i> , 1998 , 21, 221-30	14.6	44
64	Alcohol consumption and atherosclerosis: what is the relation? Prospective results from the Bruneck Study. <i>Stroke</i> , 1998 , 29, 900-7	6.7	142
63	Intimal-medial thickness of the carotid artery in nondiabetic and NIDDM patients. Relationship with insulin resistance. <i>Diabetes Care</i> , 1997 , 20, 627-31	14.6	128
62	Cigarette smoking and insulin resistance in patients with noninsulin-dependent diabetes mellitus. <i>Journal of Clinical Endocrinology and Metabolism</i> , 1997 , 82, 3619-24	5.6	134
61	The role of insulin in age-related sex differences of cardiovascular risk profile and morbidity. <i>Atherosclerosis</i> , 1997 , 130, 183-9	3.1	15
60	Obesity worsens cardiovascular risk profiles independently of hyperinsulinaemia. <i>Journal of Internal Medicine</i> , 1997 , 241, 463-70	10.8	4
59	Relationship between fasting insulin and cardiovascular risk factors is already present in young men: the Verona Young Men Atherosclerosis Risk Factors Study. <i>European Journal of Clinical Investigation</i> , 1997 , 27, 248-54	4.6	10
58	Long-term instability of fasting plasma glucose, a novel predictor of cardiovascular mortality in elderly patients with non-insulin-dependent diabetes mellitus: the Verona Diabetes Study. <i>Circulation</i> , 1997 , 96, 1750-4	16.7	99
57	Relationship between insulin and carotid atherosclerosis in the general population. The Bruneck Study. <i>Stroke</i> , 1997 , 28, 1147-52	6.7	23
56	Serum uric acid and related factors in 500 hospitalized subjects. <i>Metabolism: Clinical and Experimental</i> , 1996 , 45, 1557-61	12.7	62
55	Early changes in plasma glucagon and growth hormone response to oral glucose in experimental hyperthyroidism. <i>Metabolism: Clinical and Experimental</i> , 1996 , 45, 1029-33	12.7	15
54	Attending the diabetes center is associated with increased 5-year survival probability of diabetic patients: the Verona Diabetes Study. <i>Diabetes Care</i> , 1996 , 19, 211-3	14.6	74
53	Insulin sensitivity and regular alcohol consumption: large, prospective, cross sectional population study (Bruneck study). <i>BMJ: British Medical Journal</i> , 1996 , 313, 1040-4		240
52	Roles of glucose transport and glucose phosphorylation in muscle insulin resistance of NIDDM. <i>Diabetes</i> , 1996 , 45, 915-925	0.9	39
51	The Verona diabetes study: a population-based survey on known diabetes mellitus prevalence and 5-year all-cause mortality. <i>Diabetologia</i> , 1995 , 38, 318-25	10.3	123
50	Long-term instability of fasting plasma glucose predicts mortality in elderly NIDDM patients: the Verona Diabetes Study. <i>Diabetologia</i> , 1995 , 38, 672-9	10.3	56
49	Insulin sensitivity is not impaired in Mexican-American women without a family history of diabetes. <i>Diabetes Care</i> , 1995 , 18, 825-33	14.6	7

48	Is it possible to derive a reliable estimate of human visceral and subcutaneous abdominal adipose tissue from simple anthropometric measurements?. <i>Metabolism: Clinical and Experimental</i> , 1995 , 44, 1617-25	12.7	116
47	Mononuclear leukocytes from obese patients with type II diabetes have reduced activity of hexokinase, 6-phosphofructokinase and glucose-6-phosphate dehydrogenase. <i>Hormone and Metabolic Research</i> , 1993 , 25, 160-4	3.1	13
46	In vivo glucose metabolism in obese and type II diabetic subjects with or without hypertension. <i>Diabetes</i> , 1993 , 42, 764-72	0.9	33
45	Changes in insulin and lipid metabolism in males with asymptomatic hyperuricaemia. <i>Journal of Internal Medicine</i> , 1993 , 234, 25-30	10.8	94
44	Characterization of cellular defects of insulin action in type 2 (non-insulin-dependent) diabetes mellitus. <i>Journal of Clinical Investigation</i> , 1993 , 91, 484-94	15.9	128
43	Transmembrane glucose transport in skeletal muscle of patients with non-insulin-dependent diabetes. <i>Journal of Clinical Investigation</i> , 1993 , 92, 486-94	15.9	118
42	Detection of mutations in insulin receptor gene by denaturing gradient gel electrophoresis. <i>Diabetes</i> , 1992 , 41, 408-15	0.9	43
41	Total Body Fat Content and Fat Topography Are Associated Differently With In Vivo Glucose Metabolism in Nonobese and Obese Nondiabetic Women. <i>Diabetes</i> , 1992 , 41, 1151-1159	0.9	140
40	Influence of body fat and its regional localization on risk factors for atherosclerosis in young men. <i>American Journal of Epidemiology</i> , 1992 , 135, 1271-8	3.8	31
39	Studies on the mechanism of action of sulphonylureas in type II diabetic subjects: gliquidone. <i>Journal of Endocrinological Investigation</i> , 1992 , 15, 1-11	5.2	8
38	Total body fat content and fat topography are associated differently with in vivo glucose metabolism in nonobese and obese nondiabetic women. <i>Diabetes</i> , 1992 , 41, 1151-9	0.9	23
37	Total body fat content and fat topography are associated differently with in vivo glucose metabolism in nonobese and obese nondiabetic women. <i>Diabetes</i> , 1992 , 41, 1151-1159	0.9	33
36	Measurement of abdominal fat with T1-weighted MR images. <i>Journal of Magnetic Resonance Imaging</i> , 1991 , 1, 363-9	5.6	58
35	Plasma concentrations of glucagon during hyperglycemic clamp with or without somatostatin infusion in obese subjects. <i>Acta Diabetologica Latina</i> , 1990 , 27, 309-14		6
34	Enzymatic activities related to intermediary metabolism of glucose in circulating mononuclear cells from obese humans: relationship to enzyme activity in adipose tissue. <i>Enzyme</i> , 1990 , 43, 26-32		5
33	Normal inhibition by somatostatin of glucose-stimulated B cell secretion in the obese subjects. <i>Hormone and Metabolic Research</i> , 1990 , 22, 584-8	3.1	4
32	Glucose and insulin suppression of plasma free fatty acids in obese subjects with normal glucose tolerance or mild, newly diagnosed type 2 (non-insulin-dependent) diabetes. <i>Journal of Endocrinological Investigation</i> , 1990 , 13, 55-9	5.2	7
31	Plasma C-peptide response to oral glucose load in hyperthyroidism. <i>Journal of Endocrinological Investigation</i> , 1990 , 13, 555-8	5.2	4

30	Risk factors for coronary artery disease in healthy persons with hyperinsulinemia and normal glucose tolerance. <i>New England Journal of Medicine</i> , 1989 , 320, 702-6	59.2	663
29	Habitual leisure-time physical activity is associated with differences in various risk factors for coronary artery disease. <i>Journal of Internal Medicine</i> , 1989 , 226, 417-21	10.8	4
28	Estimates of in vivo insulin action in man: comparison of insulin tolerance tests with euglycemic and hyperglycemic glucose clamp studies. <i>Journal of Clinical Endocrinology and Metabolism</i> , 1989 , 68, 374-8	5.6	447
27	Insulin and C-peptide plasma levels in patients with severe chronic pancreatitis and fasting normoglycemia. <i>Digestive Diseases and Sciences</i> , 1988 , 33, 732-6	4	12
26	Differences in glucose metabolic enzyme activities in human adipose tissue from abdominal and gluteal regions. <i>Metabolism: Clinical and Experimental</i> , 1988 , 37, 820-3	12.7	4
25	Effect of glipizide on insulin secretion and insulin metabolism in obese type II diabetic patients. <i>Diabetes Care</i> , 1987 , 10, 687-91	14.6	8
24	Renal metabolism of C-peptide in man. <i>Journal of Clinical Endocrinology and Metabolism</i> , 1987 , 65, 494-8	5.6	53
23	Evidence that multiple risk factors for coronary artery disease exist in persons with abnormal glucose tolerance. <i>American Journal of Medicine</i> , 1987 , 83, 609-12	2.4	74
22	Influence of the menstrual cycle on glucose tolerance and insulin secretion. <i>American Journal of Obstetrics and Gynecology</i> , 1987 , 157, 140-1	6.4	24
21	Fractional hepatic extraction of insulin in man: is it constant?. <i>Journal of Endocrinological Investigation</i> , 1987 , 10, 45-9	5.2	
20	Relationship between blood pressure and plasma insulin in non-obese and obese non-diabetic subjects. <i>Diabetologia</i> , 1987 , 30, 719-23	10.3	114
19	Effect of age and environmental factors on glucose tolerance and insulin secretion in a worker population. <i>Journal of the American Geriatrics Society</i> , 1986 , 34, 271-5	5.6	46
18	Further evidence that insulin metabolism is a major determinant of peripheral insulin response to oral glucose in subjects with mild glucose intolerance. <i>Journal of Endocrinological Investigation</i> , 1986 , 9, 371-4	5.2	6
17	Effect of aging on growth hormone, ACTH and cortisol response to insulin-induced hypoglycemia in type I diabetes. <i>Acta Diabetologica Latina</i> , 1985 , 22, 159-68		2
16	Evidence for an independent relationship between plasma insulin and concentration of high density lipoprotein cholesterol and triglyceride. <i>Atherosclerosis</i> , 1985 , 55, 259-66	3.1	131
15	Glucose tolerance, insulin secretion and insulin removal in chronic persistent hepatitis. <i>Hormone and Metabolic Research</i> , 1984 , 16, 449-50	3.1	1
14	Peripheral hyperinsulinemia of simple obesity: pancreatic hypersecretion or impaired insulin metabolism?. <i>Journal of Clinical Endocrinology and Metabolism</i> , 1984 , 59, 1121-7	5.6	59
13	Hyperinsulinemia of chronic active hepatitis: impaired insulin removal rather than pancreatic hypersecretion. <i>Hormone and Metabolic Research</i> , 1984 , 16, 111-4	3.1	43

12	Lack of effect of intravenous metformin on plasma concentrations of glucose, insulin, C-peptide, glucagon and growth hormone in non-diabetic subjects. <i>Current Medical Research and Opinion</i> , 1984 , 9, 47-51	2.5	52
11	Residual B-cell function in type 1 (insulin-dependent) diabetes mellitus: its relation to clinical and metabolic features. <i>Acta Diabetologica Latina</i> , 1984 , 21, 375-83		5
10	Possible roles of insulin, glucagon, growth hormone and free fatty acids in the pathogenesis of insulin resistance of subjects with chronic liver diseases. <i>Acta Diabetologica Latina</i> , 1984 , 21, 241-50		24
9	Secretion and hepatic removal of insulin in the healthy offspring of type 2 (non-insulin-dependent) diabetic subjects. <i>Diabète & Métabolisme</i> , 1984 , 10, 250-4		3
8	Residual B cell function and insulin sensitivity in Type 1 (insulin-dependent) diabetes mellitus. <i>Diabetologia</i> , 1983 , 25, 298-298	10.3	1
7	Decreased hepatic insulin extraction in subjects with mild glucose intolerance. <i>Metabolism: Clinical and Experimental</i> , 1983 , 32, 438-46	12.7	132
6	Secretion and hepatic extraction of insulin in nondiabetic, obese, aged subjects. <i>Journal of the American Geriatrics Society</i> , 1983 , 31, 333-7	5.6	1
5	Evidence that basal beta cell activity may play a role in determining insulin sensitivity in healthy man. <i>Annals of Nutrition and Metabolism</i> , 1983 , 27, 501-4	4.5	1
4	Combined sulphonylurea-biguanide therapy for non-insulin dependent diabetics. Metabolic effects of glibenclamide and metformin or phenformin in newly diagnosed obese patients. <i>Current Medical Research and Opinion</i> , 1982 , 7, 677-683	2.5	11
3	Mortality and its Predictors in Type 2 Diabetes 103-117		
2	Michele Muggeo. <i>L Endocrinologo</i> , 1		0
1	Loss of ZnT8 function protects against diabetes by enhanced insulin secretion		3