## Rimke C Vos

## List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/7695590/publications.pdf

Version: 2024-02-01

69 1,377 papers citations

361045 20 h-index 395343 33 g-index

71 all docs 71 docs citations

71 times ranked 2332 citing authors

#	Article	IF	CITATIONS
1	Sex differences in the risk of vascular disease associated with diabetes. Biology of Sex Differences, 2020, 11, 1.	1.8	146
2	Dutch Translation and Psychometric Testing of the 9-Item Shared Decision Making Questionnaire (SDM-Q-9) and Shared Decision Making Questionnaire-Physician Version (SDM-Q-Doc) in Primary and Secondary Care. PLoS ONE, 2015, 10, e0132158.	1.1	109
3	Risk and management of pre-diabetes. European Journal of Preventive Cardiology, 2019, 26, 47-54.	0.8	87
4	Diabetes-Related Distress, Depression and Distress-Depression among Adults with Type 2 Diabetes Mellitus in Malaysia. PLoS ONE, 2016, 11, e0152095.	1.1	72
5	Psychological interventions for diabetes-related distress in adults with type 2 diabetes mellitus. The Cochrane Library, 2017, 2017, CD011469.	1.5	66
6	Developmental trajectories of receptive and expressive communication in children and young adults with cerebral palsy. Developmental Medicine and Child Neurology, 2014, 56, 951-959.	1.1	51
7	Physical fitness, functional ability and quality of life in children with severe haemophilia: a pilot study. Haemophilia, 2006, 12, 494-499.	1.0	49
8	Developmental Trajectories of Daily Activities in Children and Adolescents With Cerebral Palsy. Pediatrics, 2013, 132, e915-e923.	1.0	46
9	Diabetesâ€related distress over the course of illness: results from the Diacourse study. Diabetic Medicine, 2015, 32, 1617-1624.	1.2	43
10	Long-term effects of intensive multifactorial therapy in individuals with screen-detected type 2 diabetes in primary care: 10-year follow-up of the ADDITION-Europe cluster-randomised trial. Lancet Diabetes and Endocrinology,the, 2019, 7, 925-937.	5.5	39
11	Longitudinal development of gross motor function among <scp>D</scp> utch children and young adults with cerebral palsy: an investigation of motor growth curves. Developmental Medicine and Child Neurology, 2013, 55, 378-384.	1.1	35
12	Developmental trajectories of social participation in individuals with cerebral palsy: a multicentre longitudinal study. Developmental Medicine and Child Neurology, 2014, 56, 370-377.	1.1	32
13	Insulin monotherapy compared with the addition of oral glucose-lowering agents to insulin for people with type 2 diabetes already on insulin therapy and inadequate glycaemic control. The Cochrane Library, 2016, 9, CD006992.	1.5	32
14	Metabolomic biomarkers for personalised glucose lowering drugs treatment in type 2 diabetes. Metabolomics, 2016, 12, 27.	1.4	30
15	Long-term effect of lifestyle intervention on adiposity, metabolic parameters, inflammation and physical fitness in obese children: a randomized controlled trial. Nutrition and Diabetes, 2011, 1, e9-e9.	1.5	28
16	What determines treatment satisfaction of patients with type 2 diabetes on insulin therapy? An observational study in eight European countries. BMJ Open, 2017, 7, e016180.	0.8	28
17	Efficacy of three therapy approaches in preschool children with cerebral palsy: a randomized controlled trial. Developmental Medicine and Child Neurology, 2016, 58, 758-766.	1.1	25
18	Overtreatment of older patients with type 2 diabetes mellitus in primary care. Diabetes, Obesity and Metabolism, 2018, 20, 1066-1069.	2.2	25

#	Article	IF	CITATIONS
19	Sex differences in cardiovascular risk management for people with diabetes in primary care: a cross-sectional study. BJGP Open, 2019, 3, bjgpopen19X101645.	0.9	25
20	Validity and reliability of a Malay version of the brief illness perception questionnaire for patients with type 2 diabetes mellitus. BMC Medical Research Methodology, 2017, 17, 118.	1.4	23
21	Sex differences in the association of prediabetes and type 2 diabetes with microvascular complications and function: The Maastricht Study. Cardiovascular Diabetology, 2021, 20, 102.	2.7	23
22	The associations between diabetes distress and self-efficacy, medication adherence, self-care activities and disease control depend on the way diabetes distress is measured: Comparing the DDS-17, DDS-2 and the PAID-5. Diabetes Research and Clinical Practice, 2018, 142, 74-84.	1.1	22
23	The effectiveness of an emotionâ€focused educational programme in reducing diabetes distress in adults with Type 2 diabetes mellitus (VEMOFIT): a cluster randomized controlled trial. Diabetic Medicine, 2018, 35, 750-759.	1.2	21
24	Effectiveness of shared goal setting and decision making to achieve treatment targets in type 2 diabetes patients: A clusterâ€randomized trial (⟨scp⟩OPTIMAL⟨/scp⟩). Health Expectations, 2017, 20, 1172-1180.	1.1	19
25	Effectiveness of diabetes self-management education via a smartphone application in insulin treated type 2 diabetes patients – design of a randomised controlled trial (†TRIGGER study'). BMC Endocrine Disorders, 2018, 18, 74.	0.9	19
26	Shared decision making in type 2 diabetes with a support decision tool that takes into account clinical factors, the intensity of treatment and patient preferences: design of a cluster randomised (OPTIMAL) trial. BMC Family Practice, 2015, 16, 27.	2.9	18
27	Effectiveness of diabetes self-management education and support via a smartphone application in insulin-treated patients with type 2 diabetes: results of a randomized controlled trial (TRIGGER study). BMJ Open Diabetes Research and Care, 2019, 7, e000981.	1.2	18
28	The effect of family-based multidisciplinary cognitive behavioral treatment in children with obesity: study protocol for a randomized controlled trial. Trials, 2011, 12, 110.	0.7	17
29	Adverse differences in cardiometabolic risk factor levels between individuals with pre-diabetes and normal glucose metabolism are more pronounced in women than in men: the Maastricht Study. BMJ Open Diabetes Research and Care, 2019, 7, e000787.	1.2	17
30	Personalised treatment targets in type 2 diabetes patients: The Dutch approach. Primary Care Diabetes, 2017, 11, 71-77.	0.9	16
31	Repeat prescriptions of guideline-based secondary prevention medication in patients with type 2 diabetes and previous myocardial infarction in Dutch primary care. Family Practice, 2014, 31, 688-693.	0.8	15
32	Longitudinal Association Between Gross Motor Capacity and Neuromusculoskeletal Function in Children and Youth With Cerebral Palsy. Archives of Physical Medicine and Rehabilitation, 2016, 97, 1329-1337.	0.5	15
33	Comparison of perceptions of obesity among adults with central obesity with and without additional cardiometabolic risk factors and among those who were formally obese, 3 years after screening for central obesity. BMC Public Health, 2015, 15, 1214.	1.2	10
34	Theoryâ€based diabetes selfâ€management education with preâ€selection of participants: a randomized controlled trial with 2.5 years' followâ€up ( <scp>ELDES</scp> Study). Diabetic Medicine, 2019, 36, 827-835.	1.2	10
35	Effectiveness of tailored support for people with Type 2 diabetes after a first acute coronary event: a multicentre randomized controlled trial (the Diacourseâ€≺scp>ACE study). Diabetic Medicine, 2016, 33, 125-133.	1.2	9
36	Diabetes care providers' opinions and working methods after four years of experience with a diabetes patient web portal; a survey among health care providers in general practices and an outpatient clinic. BMC Family Practice, 2018, 19, 94.	2.9	9

#	Article	IF	CITATIONS
37	<p>De-Intensification Of Blood Glucose Lowering Medication In People Identified As Being Over-Treated: A Mixed Methods Study</p> . Patient Preference and Adherence, 2019, Volume 13, 1775-1783.	0.8	9
38	Large health disparities in cardiovascular death in men and women, by ethnicity and socioeconomic status in an urban based population cohort. EClinicalMedicine, 2021, 40, 101120.	3.2	9
39	The effectiveness of a value-based EMOtion-cognition-Focused educational programme to reduce diabetes-related distress in Malay adults with Type 2 diabetes (VEMOFIT): study protocol for a cluster randomised controlled trial. BMC Endocrine Disorders, 2017, 17, 22.	0.9	8
40	Psychological factors associated with changes in physical activity in Dutch people with type 2 diabetes under societal lockdown: A crossâ€sectional study. Endocrinology, Diabetes and Metabolism, 2021, 4, e00249.	1.0	8
41	Measuring positive health: Concurrent and factorial validity based on a representative Dutch sample. Health and Social Care in the Community, 2022, 30, .	0.7	8
42	Diabetes-Related Distress and Depressive Symptoms Are Not Merely Negative over a 3-Year Period in Malaysian Adults with Type 2 Diabetes Mellitus Receiving Regular Primary Diabetes Care. Frontiers in Psychology, 2017, 8, 1834.	1,1	7
43	Achieving diabetes treatment targets in people with registered mental illness is similar or improved compared with those without: Analyses of linked observational datasets. Diabetic Medicine, 2022, 39, e14835.	1.2	7
44	What follow-up care and self-management support do patients with type 2 diabetes want after their first acute coronary event? A qualitative study. Primary Care Diabetes, 2014, 8, 195-206.	0.9	6
45	Association of weight loss and weight loss maintenance following diabetes diagnosis by screening and incidence of cardiovascular disease and all ause mortality: An observational analysis of the ADDITIONâ€Europe trial. Diabetes, Obesity and Metabolism, 2021, 23, 730-741.	2.2	6
46	A simple to implement and low-cost supervised walking programme in highly motivated individuals with or at risk for type 2 diabetes: An observational study with a pre-post design. Preventive Medicine Reports, 2019, 13, 30-36.	0.8	5
47	The influence of an educational program on the HbA1c-level of adolescents with type 1 diabetes mellitus: a retrospective study. Journal of Pediatric Endocrinology and Metabolism, 2011, 24, 15-9.	0.4	4
48	The Effect of Multidisciplinary Lifestyle Intervention on the Pre- and Postprandial Plasma Gut Peptide Concentrations in Children with Obesity. Isrn Endocrinology, 2011, 2011, 1-7.	2.0	4
49	Differences in clinical characteristics between patients with and without type 2 diabetes hospitalized with a first myocardial infarction. Journal of Diabetes and Its Complications, 2016, 30, 830-833.	1.2	4
50	Population-based screen-detected type 2 diabetes mellitus is associated with less need for insulin therapy after 10 years. BMJ Open Diabetes Research and Care, 2020, 8, e000949.	1.2	4
51	Type 2 diabetes in South Asians compared to Europeans: Higher risk and earlier development of major cardiovascular events irrespective of the presence and degree of retinopathy. Results from The HinDu The Hague Diabetes Study. Endocrinology, Diabetes and Metabolism, 2021, 4, e00242.	1.0	4
52	Sex Disparities in Cardiovascular Risk Factor Assessment and Screening for Diabetes-Related Complications in Individuals With Diabetes: A Systematic Review. Frontiers in Endocrinology, 2021, 12, 617902.	1.5	4
53	Effect of six years intensified multifactorial treatment on levels of hs RP and adiponectin in patients with screen detected type 2 diabetes: The ADDITIONâ€Netherlands randomized trial. Diabetes/Metabolism Research and Reviews, 2015, 31, 758-766.	1.7	3
54	Diabetes self-management education after pre-selection of patients: design of a randomised controlled trial. Diabetology and Metabolic Syndrome, 2016, 8, 82.	1,2	3

#	Article	IF	Citations
55	The effectiveness of an emotion-focused educational programme in reducing diabetes distress in adults with type 2 diabetes mellitus at 12-month follow-up: a cluster randomized controlled trial. Therapeutic Advances in Endocrinology and Metabolism, 2019, 10, 204201881985376.	1.4	3
56	Individualised targets for insulin initiation in type 2 diabetes mellitusâ€"the influence of physician and practice: a cross-sectional study in eight European countries. BMJ Open, 2019, 9, e032040.	0.8	3
57	Internet-based Self-Management Support for Patients With Well-Controlled Type 2 Diabetes: A Real-Life Study. JMIR Research Protocols, 2017, 6, e47.	0.5	3
58	Oral Hypoglycemic Agents Added to Insulin Monotherapy for Type 2 Diabetes. JAMA - Journal of the American Medical Association, 2017, 318, 1489.	3.8	2
59	Insulin Therapy in Type 2 Diabetes Is Associated With Barriers to Activity and Worse Health Status: A Cross-Sectional Study in Primary Care. Frontiers in Endocrinology, 2021, 12, 573235.	1.5	2
60	Discussing overweight in children during a regular consultation in general practice: a qualitative study. BMC Family Practice, 2020, 21, 18.	2.9	2
61	Editorial: Achieving Effective Management and Treatment of Diabetes Mellitus in Future Primary Care. Frontiers in Endocrinology, 2022, 13, 854244.	1.5	2
62	Shared decision making in primary care: Process evaluation of the intervention in the OPTIMAL study, a cluster randomised trial. Primary Care Diabetes, 2022, 16, 375-380.	0.9	2
63	Cluster randomised trial on the effectiveness of a computerised prompt to refer (back) patients with type 2 diabetes. PLoS ONE, 2018, 13, e0207653.	1.1	1
64	Patientâ€reported outcomes after 10â€year followâ€up of intensive, multifactorial treatment in individuals with screenâ€detected type 2 diabetes: the ADDITIONâ€Europe trial. Diabetic Medicine, 2020, 37, 1509-1518.	1.2	1
65	Effectiveness of the Beyond Good Intentions Program on Improving Dietary Quality Among People With Type 2 Diabetes Mellitus: A Randomized Controlled Trial. Frontiers in Nutrition, 2021, 8, 583125.	1.6	1
66	Type 2 diabetes and COPD: treatment in the right healthcare setting? An observational study. BMC Family Practice, 2021, 22, 78.	2.9	1
67	Editorial: Achieving Efficient Diabetes Care Through Understanding the Risk Factors, Markers, and Patient Experiences. Frontiers in Endocrinology, 2022, 13, 854167.	1.5	1
68	Diabetes self-management education and support delivered by mobile health (m-health) interventions for adults with type 2 diabetes mellitus. The Cochrane Library, 0, , .	1.5	0
69	The impact of the new Dutch guideline on cardiovascular risk management in patients with COPD: a retrospective study. BJGP Open, 2021, 5, bjgpopen20X101139.	0.9	0