

Rainer Wolf

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/7698551/publications.pdf>

Version: 2024-02-01

8
papers

380
citations

1307594
7
h-index

1588992
8
g-index

8
all docs

8
docs citations

8
times ranked

932
citing authors

#	ARTICLE	IF	CITATIONS
1	The role of microglia in neuropsychiatric disorders and suicide. <i>European Archives of Psychiatry and Clinical Neuroscience</i> , 2022, 272, 929-945.	3.2	26
2	Effects of neonatal excitotoxic lesions in ventral thalamus on social interaction in the rat. <i>European Archives of Psychiatry and Clinical Neuroscience</i> , 2018, 268, 461-470.	3.2	4
3	First and second generation antipsychotics differentially affect structural and functional properties of rat hippocampal neuron synapses. <i>Neuroscience</i> , 2016, 337, 117-130.	2.3	16
4	Calretinin and parvalbumin in schizophrenia and affective disorders: a mini-review, a perspective on the evolutionary role of calretinin in schizophrenia, and a preliminary post-mortem study of calretinin in the septal nuclei. <i>Frontiers in Cellular Neuroscience</i> , 2015, 9, 393.	3.7	12
5	The Role of Dopamine in Schizophrenia from a Neurobiological and Evolutionary Perspective: Old Fashioned, but Still in Vogue. <i>Frontiers in Psychiatry</i> , 2014, 5, 47.	2.6	273
6	The role of microglia during life span in neuropsychiatric disease – an animal study. <i>Schizophrenia Research</i> , 2013, 143, 221-222.	2.0	26
7	Acute or subchronic clozapine treatment does not ameliorate prepulse inhibition (PPI) deficits in CPB-K mice with low levels of hippocampal NMDA receptor density. <i>Psychopharmacology</i> , 2007, 194, 93-102.	3.1	10
8	Prepulse inhibition is different in two inbred mouse strains (CPB-K and BALB/c) with different hippocampal NMDA receptor densities. <i>Behavioural Brain Research</i> , 2006, 166, 78-84.	2.2	13