

Giacomo Zoppini

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/7700806/publications.pdf>

Version: 2024-02-01

109
papers

7,835
citations

53660

45
h-index

51492

86
g-index

109
all docs

109
docs citations

109
times ranked

9598
citing authors

#	ARTICLE	IF	CITATIONS
1	Non-alcoholic fatty liver disease and risk of incident cardiovascular disease: A meta-analysis. <i>Journal of Hepatology</i> , 2016, 65, 589-600.	1.8	965
2	Relations Between Carotid Artery Wall Thickness and Liver Histology in Subjects With Nonalcoholic Fatty Liver Disease. <i>Diabetes Care</i> , 2006, 29, 1325-1330.	4.3	362
3	Both resistance training and aerobic training reduce hepatic fat content in type 2 diabetic subjects with nonalcoholic fatty liver disease (the RAED2 randomized trial). <i>Hepatology</i> , 2013, 58, 1287-1295.	3.6	275
4	Nonalcoholic fatty liver disease increases risk of incident chronic kidney disease: A systematic review and meta-analysis. <i>Metabolism: Clinical and Experimental</i> , 2018, 79, 64-76.	1.5	261
5	Non-alcoholic fatty liver disease in patients with chronic plaque psoriasis. <i>Journal of Hepatology</i> , 2009, 51, 758-764.	1.8	217
6	Serum Uric Acid Levels and Incident Chronic Kidney Disease in Patients With Type 2 Diabetes and Preserved Kidney Function. <i>Diabetes Care</i> , 2012, 35, 99-104.	4.3	207
7	Prevalence of non-alcoholic fatty liver disease and its association with cardiovascular disease in patients with type 1 diabetes. <i>Journal of Hepatology</i> , 2010, 53, 713-718.	1.8	202
8	Prevalence of Subclinical Hypothyroidism in Patients with Chronic Kidney Disease. <i>Clinical Journal of the American Society of Nephrology: CJASN</i> , 2008, 3, 1296-1300.	2.2	200
9	Relationship between Kidney Function and Liver Histology in Subjects with Nonalcoholic Steatohepatitis. <i>Clinical Journal of the American Society of Nephrology: CJASN</i> , 2010, 5, 2166-2171.	2.2	197
10	Increased Risk of CKD among Type 2 Diabetics with Nonalcoholic Fatty Liver Disease. <i>Journal of the American Society of Nephrology: JASN</i> , 2008, 19, 1564-1570.	3.0	187
11	NASH Predicts Plasma Inflammatory Biomarkers Independently of Visceral Fat in Men. <i>Obesity</i> , 2008, 16, 1394-1399.	1.5	180
12	Predictors of Estimated GFR Decline in Patients with Type 2 Diabetes and Preserved Kidney Function. <i>Clinical Journal of the American Society of Nephrology: CJASN</i> , 2012, 7, 401-408.	2.2	178
13	Metabolic Effects of Aerobic Training and Resistance Training in Type 2 Diabetic Subjects. <i>Diabetes Care</i> , 2012, 35, 676-682.	4.3	177
14	Relation of Nonalcoholic Hepatic Steatosis to Early Carotid Atherosclerosis in Healthy Men: Role of visceral fat accumulation. <i>Diabetes Care</i> , 2004, 27, 2498-2500.	4.3	173
15	Nonalcoholic Fatty Liver Disease Is Associated With Left Ventricular Diastolic Dysfunction in Patients With Type 2 Diabetes. <i>Diabetes Care</i> , 2012, 35, 389-395.	4.3	159
16	Associations between plasma adiponectin concentrations and liver histology in patients with nonalcoholic fatty liver disease. <i>Clinical Endocrinology</i> , 2006, 64, 679-683.	1.2	156
17	Non-Alcoholic Fatty Liver Disease Is Associated with an Increased Incidence of Atrial Fibrillation in Patients with Type 2 Diabetes. <i>PLoS ONE</i> , 2013, 8, e57183.	1.1	153
18	Risk of chronic kidney disease in patients with non-alcoholic fatty liver disease: Is there a link?. <i>Journal of Hepatology</i> , 2011, 54, 1020-1029.	1.8	152

#	ARTICLE	IF	CITATIONS
19	Relationship between red blood cell distribution width and kidney function tests in a large cohort of unselected outpatients. <i>Scandinavian Journal of Clinical and Laboratory Investigation</i> , 2008, 68, 745-748.	0.6	139
20	Effects of moderate-intensity exercise training on plasma biomarkers of inflammation and endothelial dysfunction in older patients with type 2 diabetes. <i>Nutrition, Metabolism and Cardiovascular Diseases</i> , 2006, 16, 543-549.	1.1	130
21	Nonalcoholic Fatty Liver Disease Is Independently Associated With an Increased Incidence of Chronic Kidney Disease in Patients With Type 1 Diabetes. <i>Diabetes Care</i> , 2014, 37, 1729-1736.	4.3	129
22	Nonalcoholic Fatty Liver Disease as a Contributor to Hypercoagulation and Thrombophilia in the Metabolic Syndrome. <i>Seminars in Thrombosis and Hemostasis</i> , 2009, 35, 277-287.	1.5	123
23	Mortality From Chronic Liver Diseases in Diabetes. <i>American Journal of Gastroenterology</i> , 2014, 109, 1020-1025.	0.2	121
24	Elevated Serum Uric Acid Concentrations Independently Predict Cardiovascular Mortality in Type 2 Diabetic Patients. <i>Diabetes Care</i> , 2009, 32, 1716-1720.	4.3	111
25	Non-alcoholic fatty liver disease is associated with an increased prevalence of atrial fibrillation in hospitalized patients with Type 2 diabetes. <i>Clinical Science</i> , 2013, 125, 301-310.	1.8	107
26	Nonalcoholic Fatty Liver Disease Is Associated With Ventricular Arrhythmias in Patients With Type 2 Diabetes Referred for Clinically Indicated 24-Hour Holter Monitoring. <i>Diabetes Care</i> , 2016, 39, 1416-1423.	4.3	95
27	Association Between Primary Hypothyroidism and Nonalcoholic Fatty Liver Disease: A Systematic Review and Meta-Analysis. <i>Thyroid</i> , 2018, 28, 1270-1284.	2.4	87
28	Associations between liver histology and cortisol secretion in subjects with nonalcoholic fatty liver disease. <i>Clinical Endocrinology</i> , 2006, 64, 337-341.	1.2	83
29	Heart valve calcification in patients with type 2 diabetes and nonalcoholic fatty liver disease. <i>Metabolism: Clinical and Experimental</i> , 2015, 64, 879-887.	1.5	82
30	Prognostic Impact of Diabetes on Long-term Survival Outcomes in Patients With Heart Failure: A Meta-analysis. <i>Diabetes Care</i> , 2017, 40, 1597-1605.	4.3	82
31	Nonalcoholic Fatty Liver Disease Is Independently Associated with Early Left Ventricular Diastolic Dysfunction in Patients with Type 2 Diabetes. <i>PLoS ONE</i> , 2015, 10, e0135329.	1.1	81
32	Association between nonalcoholic fatty liver disease and colorectal tumours in asymptomatic adults undergoing screening colonoscopy: a systematic review and meta-analysis. <i>Metabolism: Clinical and Experimental</i> , 2018, 87, 1-12.	1.5	80
33	Association of nonalcoholic fatty liver disease with QTc interval in patients with type 2 diabetes. <i>Nutrition, Metabolism and Cardiovascular Diseases</i> , 2014, 24, 663-669.	1.1	77
34	Association between nonalcoholic fatty liver disease and risk of atrial fibrillation in adult individuals: An updated meta-analysis. <i>Liver International</i> , 2019, 39, 758-769.	1.9	75
35	Prevalence of neuropathy in type 2 diabetic patients and its association with other diabetes complications: The Verona Diabetic Foot Screening Program. <i>Journal of Diabetes and Its Complications</i> , 2015, 29, 1066-1070.	1.2	69
36	Aortic and Mitral Annular Calcifications Are Predictive of All-Cause and Cardiovascular Mortality in Patients With Type 2 Diabetes. <i>Diabetes Care</i> , 2012, 35, 1781-1786.	4.3	62

#	ARTICLE	IF	CITATIONS
37	Triglycerideâ€“high-density lipoprotein cholesterol is associated with microvascular complications in type 2 diabetes mellitus. <i>Metabolism: Clinical and Experimental</i> , 2012, 61, 22-29.	1.5	62
38	Variability of body weight, pulse pressure and glycaemia strongly predict total mortality in elderly type 2 diabetic patients. <i>The Verona Diabetes Study. Diabetes/Metabolism Research and Reviews</i> , 2008, 24, 624-628.	1.7	61
39	Association between serum TSH, free T4 and serum liver enzyme activities in a large cohort of unselected outpatients. <i>Clinical Endocrinology</i> , 2008, 68, 481-484.	1.2	60
40	Relation of Elevated Serum Uric Acid Levels to Incidence of Atrial Fibrillation in Patients With Type 2 Diabetes Mellitus. <i>American Journal of Cardiology</i> , 2013, 112, 499-504.	0.7	58
41	Nonalcoholic fatty liver disease is independently associated with an increased incidence of cardiovascular disease in adult patients with type 1 diabetes. <i>International Journal of Cardiology</i> , 2016, 225, 387-391.	0.8	56
42	Prevalence of Cardiovascular Autonomic Neuropathy in a Cohort of Patients With Newly Diagnosed Type 2 Diabetes: The Verona Newly Diagnosed Type 2 Diabetes Study (VNDS). <i>Diabetes Care</i> , 2015, 38, 1487-1493.	4.3	55
43	Non-alcoholic fatty liver disease is independently associated with left ventricular hypertrophy in hypertensive Type 2 diabetic individuals. <i>Journal of Endocrinological Investigation</i> , 2012, 35, 215-218.	1.8	54
44	Effect of moderate aerobic exercise on sympatho-vagal balance in Type 2 diabetic patients. <i>Diabetic Medicine</i> , 2007, 24, 370-376.	1.2	50
45	High-Normal HbA1c Is a Strong Predictor of Type 2 Diabetes in the General Population. <i>Diabetes Care</i> , 2011, 34, 1038-1040.	4.3	47
46	The aspartate aminotransferase-to-alanine aminotransferase ratio predicts all-cause and cardiovascular mortality in patients with type 2 diabetes. <i>Medicine (United States)</i> , 2016, 95, e4821.	0.4	47
47	Nonalcoholic fatty liver disease is associated with an increased prevalence of distal symmetric polyneuropathy in adult patients with type 1 diabetes. <i>Journal of Diabetes and Its Complications</i> , 2017, 31, 1021-1026.	1.2	47
48	Glycated Haemoglobin Is Inversely Related to Serum Vitamin D Levels in Type 2 Diabetic Patients. <i>PLoS ONE</i> , 2013, 8, e82733.	1.1	47
49	Disorders of Coagulation and Hemostasis in Abdominal Obesity: Emerging Role of Fatty Liver. <i>Seminars in Thrombosis and Hemostasis</i> , 2010, 36, 041-048.	1.5	46
50	Lower levels of 25-hydroxyvitamin D ₃ are associated with a higher prevalence of microvascular complications in patients with type 2 diabetes. <i>BMJ Open Diabetes Research and Care</i> , 2015, 3, e000058.	1.2	45
51	Systematic review with metaâ€“analysis: nonâ€“alcoholic fatty liver disease is associated with a history of osteoporotic fractures but not with low bone mineral density. <i>Alimentary Pharmacology and Therapeutics</i> , 2019, 49, 375-388.	1.9	45
52	Mortality from infectious diseases in diabetes. <i>Nutrition, Metabolism and Cardiovascular Diseases</i> , 2018, 28, 444-450.	1.1	43
53	Nonalcoholic fatty liver disease is associated with an increased risk of heart block in hospitalized patients with type 2 diabetes mellitus. <i>PLoS ONE</i> , 2017, 12, e0185459.	1.1	42
54	Non-alcoholic fatty liver disease and increased risk of all-cause mortality in elderly patients admitted for acute heart failure. <i>International Journal of Cardiology</i> , 2018, 265, 162-168.	0.8	41

#	ARTICLE	IF	CITATIONS
55	Prevalence of thyroid autoimmunity and subclinical hypothyroidism in persons with chronic kidney disease not requiring chronic dialysis. <i>Clinical Chemistry and Laboratory Medicine</i> , 2009, 47, 1367-71.	1.4	39
56	Association between <i>Helicobacter pylori</i> infection and risk of nonalcoholic fatty liver disease: An updated meta-analysis. <i>Metabolism: Clinical and Experimental</i> , 2019, 96, 56-65.	1.5	38
57	Nonalcoholic fatty liver disease and increased risk of 1-year all-cause and cardiac hospital readmissions in elderly patients admitted for acute heart failure. <i>PLoS ONE</i> , 2017, 12, e0173398.	1.1	38
58	Multiple causes of death analysis of chronic diseases: the example of diabetes. <i>Population Health Metrics</i> , 2015, 13, 21.	1.3	35
59	Chronic complications in patients with newly diagnosed type 2 diabetes: prevalence and related metabolic and clinical features: the Verona Newly Diagnosed Type 2 Diabetes Study (VNDS) 9. <i>BMJ Open Diabetes Research and Care</i> , 2020, 8, e001549.	1.2	35
60	Anaemia, independent of chronic kidney disease, predicts all-cause and cardiovascular mortality in type 2 diabetic patients. <i>Atherosclerosis</i> , 2010, 210, 575-580.	0.4	32
61	Association Between Nonalcoholic Fatty Liver Disease and Reduced Bone Mineral Density in Children: A Meta-Analysis. <i>Hepatology</i> , 2019, 70, 812-823.	3.6	30
62	Hypertriglyceridemia Is Independently Associated with Renal, but Not Retinal Complications in Subjects with Type 2 Diabetes: A Cross-Sectional Analysis of the Renal Insufficiency And Cardiovascular Events (RIACE) Italian Multicenter Study. <i>PLoS ONE</i> , 2015, 10, e0125512.	1.1	30
63	Letter to the Editor. <i>Arteriosclerosis, Thrombosis, and Vascular Biology</i> , 2005, 25, 2687-2688.	1.1	29
64	Relationship of nonalcoholic hepatic steatosis to overnight low-dose dexamethasone suppression test in obese individuals. <i>Clinical Endocrinology</i> , 2004, 61, 711-715.	1.2	26
65	The role of serum uric acid in cardiovascular disease in Type 2 diabetic and non-diabetic subjects: A narrative review. <i>Journal of Endocrinological Investigation</i> , 2011, 34, 881-886.	1.8	26
66	Comparison of Two Creatinine-Based Estimating Equations in Predicting All-Cause and Cardiovascular Mortality in Patients With Type 2 Diabetes. <i>Diabetes Care</i> , 2012, 35, 2347-2353.	4.3	26
67	Usefulness of Subclinical Left Ventricular Midwall Dysfunction to Predict Cardiovascular Mortality in Patients With Type 2 Diabetes Mellitus. <i>American Journal of Cardiology</i> , 2014, 113, 1409-1414.	0.7	26
68	Hemostatic Disorders in Type 1 Diabetes Mellitus. <i>Seminars in Thrombosis and Hemostasis</i> , 2011, 37, 058-065.	1.5	24
69	Early impairment in left ventricular longitudinal systolic function is associated with an increased risk of incident atrial fibrillation in patients with type 2 diabetes. <i>Journal of Diabetes and Its Complications</i> , 2017, 31, 413-418.	1.2	24
70	Relationship between serum bilirubin and kidney function in non-diabetic and diabetic individuals. <i>Kidney International</i> , 2009, 75, 863.	2.6	21
71	Usefulness of the triglyceride to high-density lipoprotein cholesterol ratio for predicting mortality risk in type 2 diabetes: Role of kidney dysfunction. <i>Atherosclerosis</i> , 2010, 212, 287-291.	0.4	19
72	Relationship of Serum γ -Glutamyltransferase to Atherogenic Dyslipidemia and Glycemic Control in Type 2 Diabetes. <i>Obesity</i> , 2009, 17, 370-374.	1.5	18

#	ARTICLE	IF	CITATIONS
73	Hemostatic and Fibrinolytic Abnormalities in Polycystic Ovary Syndrome. <i>Seminars in Thrombosis and Hemostasis</i> , 2014, 40, 600-618.	1.5	18
74	Diabetes and cancer mortality: A multifaceted association. <i>Diabetes Research and Clinical Practice</i> , 2014, 106, e86-e89.	1.1	18
75	Mitral Regurgitation and Increased Risk of All-Cause and Cardiovascular Mortality in Patients with Type 2 Diabetes. <i>American Journal of Medicine</i> , 2017, 130, 70-76.e1.	0.6	18
76	Nonalcoholic Fatty Liver Disease Is Associated With Higher 1-year All-Cause Rehospitalization Rates in Patients Admitted for Acute Heart Failure. <i>Medicine (United States)</i> , 2016, 95, e2760.	0.4	17
77	Evidence of left atrial remodeling and left ventricular diastolic dysfunction in type 2 diabetes mellitus with preserved systolic function. <i>Nutrition, Metabolism and Cardiovascular Diseases</i> , 2016, 26, 1026-1032.	1.1	16
78	Inappropriate left ventricular mass independently predicts cardiovascular mortality in patients with type 2 diabetes. <i>International Journal of Cardiology</i> , 2013, 168, 4953-4956.	0.8	15
79	Prevalence of diabetes across different immigrant groups in North-eastern Italy. <i>Nutrition, Metabolism and Cardiovascular Diseases</i> , 2015, 25, 924-930.	1.1	15
80	Risk of all-cause and cardiovascular mortality in patients with chronic liver disease. <i>Gut</i> , 2011, 60, 1602-1603.	6.1	13
81	Relationship between increased left atrial volume and microvascular complications in patients with type 2 diabetes. <i>Journal of Diabetes and Its Complications</i> , 2015, 29, 822-828.	1.2	12
82	Hemostatic and Fibrinolytic Abnormalities in Endocrine Diseases: A Narrative Review. <i>Seminars in Thrombosis and Hemostasis</i> , 2009, 35, 605-612.	1.5	11
83	Association between subclinical left ventricular systolic dysfunction and glycemic control in asymptomatic type 2 diabetic patients with preserved left ventricular function. <i>Journal of Diabetes and Its Complications</i> , 2017, 31, 1035-1040.	1.2	11
84	Relation of elevated serum uric acid levels to first-degree heart block and other cardiac conduction defects in hospitalized patients with type 2 diabetes. <i>Journal of Diabetes and Its Complications</i> , 2017, 31, 1691-1697.	1.2	10
85	The E/e [™] ratio difference between subjects with type 2 diabetes and controls. A meta-analysis of clinical studies. <i>PLoS ONE</i> , 2018, 13, e0209794.	1.1	10
86	SARS-CoV-2 and COVID-19 in diabetes mellitus. Population-based study on ascertained infections, hospital admissions and mortality in an Italian region with ≈ 4.5 million inhabitants and $\approx 250,000$ diabetic people. <i>Nutrition, Metabolism and Cardiovascular Diseases</i> , 2021, 31, 2612-2618.	1.1	10
87	Soluble CD40L in Young Type 1 Diabetic Individuals Without Clinical Microvascular and Macrovascular Complications. <i>Diabetes Care</i> , 2004, 27, 1236-1237.	4.3	9
88	Pulse Pressure and Mortality from Cerebrovascular Diseases in Type 2 Diabetic Patients: The Verona Diabetes Study. <i>Cerebrovascular Diseases</i> , 2007, 23, 20-26.	0.8	8
89	Independent correlates of urinary albumin excretion within the normoalbuminuric range in patients with type 2 diabetes: The Renal Insufficiency And Cardiovascular Events (RIACE) Italian Multicentre Study. <i>Acta Diabetologica</i> , 2015, 52, 971-981.	1.2	8
90	Severe hypoglycemia in patients with known diabetes requiring emergency department care: A report from an Italian multicenter study. <i>Journal of Clinical and Translational Endocrinology</i> , 2016, 5, 46-52.	1.0	8

#	ARTICLE	IF	CITATIONS
91	Long-Acting GLP-1 Receptor Agonist Exenatide Influence on the Autonomic Cardiac Sympatho-Vagal Balance. <i>Journal of the Endocrine Society</i> , 2018, 2, 53-62.	0.1	8
92	Effect of Serum Gamma-Glutamyltransferase and Obesity on the Risk of Dyslipidemia and Poor Glycemic Control in Type 2 Diabetic Patients: Cross-Sectional Findings from the Verona Diabetes Study. <i>Clinical Chemistry</i> , 2007, 53, 1867-1869.	1.5	6
93	A renal genetic risk score (GRS) is associated with kidney dysfunction in people with type 2 diabetes. <i>Diabetes Research and Clinical Practice</i> , 2018, 144, 137-143.	1.1	5
94	Increased aortic stiffness index in patients with type 1 diabetes without cardiovascular disease compared to controls. <i>Journal of Endocrinological Investigation</i> , 2019, 42, 1109-1115.	1.8	5
95	Nonalcoholic Fatty Liver Disease and Implications for Older Adults with Diabetes. <i>Clinics in Geriatric Medicine</i> , 2020, 36, 527-547.	1.0	5
96	Left ventricular chamber dilation and filling pressure may help to categorise patients with type 2 diabetes. <i>BMJ Open Diabetes Research and Care</i> , 2018, 6, e000529.	1.2	4
97	Time series of diabetes attributable mortality from 2008 to 2017. <i>Journal of Endocrinological Investigation</i> , 2022, 45, 275-278.	1.8	4
98	Thyroidectomies in Italy: A Population-Based National Analysis from 2001 to 2018. <i>Thyroid</i> , 2022, 32, 263-272.	2.4	4
99	Glomerular filtration rate decline in T2DM following diagnosis. The Verona newly diagnosed diabetes study-12. <i>Diabetes Research and Clinical Practice</i> , 2021, 175, 108778.	1.1	3
100	Estimating the real burden of cardiovascular mortality in diabetes. <i>European Review for Medical and Pharmacological Sciences</i> , 2019, 23, 6700-6706.	0.5	3
101	Relationship between soluble CD40 ligand and gamma-glutamyltransferase concentrations in non-drinking, young type 1 diabetic individuals.. <i>Diabetic Medicine</i> , 2008, 25, 1283-8.	1.2	2
102	Impact of Reference Category and Number of Traits in the Cluster on Risk of Coronary Heart Disease in Metabolic Syndrome: Prospective Data from the Bruneck Study. <i>Metabolic Syndrome and Related Disorders</i> , 2011, 9, 313-318.	0.5	2
103	Insulin effect on serum potassium and autoinhibition of insulin secretion is intact in a patient with leprechaunism despite severe impairment of substrates metabolism. <i>Diabetes/Metabolism Research and Reviews</i> , 2008, 24, 205-210.	1.7	1
104	Echocardiographic parameters according to insulin dose in young patients affected by type 1 diabetes. <i>PLoS ONE</i> , 2020, 15, e0244483.	1.1	0
105	Title is missing!. , 2020, 15, e0244483.		0
106	Title is missing!. , 2020, 15, e0244483.		0
107	Title is missing!. , 2020, 15, e0244483.		0
108	Title is missing!. , 2020, 15, e0244483.		0

#	ARTICLE	IF	CITATIONS
109	Estimated peak systolic pulmonary artery pressure in young non-complicated patients with type 1 diabetes. <i>European Review for Medical and Pharmacological Sciences</i> , 2020, 24, 5028-5035.	0.5	0