

Giacomo Zoppini

List of Publications by Citations

Source: <https://exaly.com/author-pdf/7700806/giacomo-zoppini-publications-by-citations.pdf>

Version: 2024-04-25

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

108
papers

5,882
citations

42
h-index

76
g-index

109
ext. papers

6,897
ext. citations

6.1
avg. IF

5.57
L-index

#	Paper	IF	Citations
108	Non-alcoholic fatty liver disease and risk of incident cardiovascular disease: A meta-analysis. <i>Journal of Hepatology</i> , 2016 , 65, 589-600	13.4	640
107	Relations between carotid artery wall thickness and liver histology in subjects with nonalcoholic fatty liver disease. <i>Diabetes Care</i> , 2006 , 29, 1325-30	14.6	312
106	Both resistance training and aerobic training reduce hepatic fat content in type 2 diabetic subjects with nonalcoholic fatty liver disease (the RAED2 Randomized Trial). <i>Hepatology</i> , 2013 , 58, 1287-95	11.2	211
105	Non-alcoholic fatty liver disease in patients with chronic plaque psoriasis. <i>Journal of Hepatology</i> , 2009 , 51, 758-64	13.4	179
104	Prevalence of non-alcoholic fatty liver disease and its association with cardiovascular disease in patients with type 1 diabetes. <i>Journal of Hepatology</i> , 2010 , 53, 713-8	13.4	171
103	Nonalcoholic fatty liver disease increases risk of incident chronic kidney disease: A systematic review and meta-analysis. <i>Metabolism: Clinical and Experimental</i> , 2018 , 79, 64-76	12.7	171
102	Serum uric acid levels and incident chronic kidney disease in patients with type 2 diabetes and preserved kidney function. <i>Diabetes Care</i> , 2012 , 35, 99-104	14.6	167
101	Relationship between kidney function and liver histology in subjects with nonalcoholic steatohepatitis. <i>Clinical Journal of the American Society of Nephrology: CJASN</i> , 2010 , 5, 2166-71	6.9	156
100	NASH predicts plasma inflammatory biomarkers independently of visceral fat in men. <i>Obesity</i> , 2008 , 16, 1394-9	8	152
99	Relation of nonalcoholic hepatic steatosis to early carotid atherosclerosis in healthy men: role of visceral fat accumulation. <i>Diabetes Care</i> , 2004 , 27, 2498-500	14.6	150
98	Increased risk of CKD among type 2 diabetics with nonalcoholic fatty liver disease. <i>Journal of the American Society of Nephrology: JASN</i> , 2008 , 19, 1564-70	12.7	143
97	Associations between plasma adiponectin concentrations and liver histology in patients with nonalcoholic fatty liver disease. <i>Clinical Endocrinology</i> , 2006 , 64, 679-83	3.4	141
96	Predictors of estimated GFR decline in patients with type 2 diabetes and preserved kidney function. <i>Clinical Journal of the American Society of Nephrology: CJASN</i> , 2012 , 7, 401-8	6.9	138
95	Prevalence of subclinical hypothyroidism in patients with chronic kidney disease. <i>Clinical Journal of the American Society of Nephrology: CJASN</i> , 2008 , 3, 1296-300	6.9	138
94	Nonalcoholic fatty liver disease is associated with left ventricular diastolic dysfunction in patients with type 2 diabetes. <i>Diabetes Care</i> , 2012 , 35, 389-95	14.6	134
93	Risk of chronic kidney disease in patients with non-alcoholic fatty liver disease: is there a link?. <i>Journal of Hepatology</i> , 2011 , 54, 1020-9	13.4	131
92	Metabolic effects of aerobic training and resistance training in type 2 diabetic subjects: a randomized controlled trial (the RAED2 study). <i>Diabetes Care</i> , 2012 , 35, 676-82	14.6	123

91	Non-alcoholic fatty liver disease is associated with an increased incidence of atrial fibrillation in patients with type 2 diabetes. <i>PLoS ONE</i> , 2013 , 8, e57183	3.7	119
90	Relationship between red blood cell distribution width and kidney function tests in a large cohort of unselected outpatients. <i>Scandinavian Journal of Clinical and Laboratory Investigation</i> , 2008 , 68, 745-8	2	112
89	Effects of moderate-intensity exercise training on plasma biomarkers of inflammation and endothelial dysfunction in older patients with type 2 diabetes. <i>Nutrition, Metabolism and Cardiovascular Diseases</i> , 2006 , 16, 543-9	4.5	112
88	Nonalcoholic fatty liver disease as a contributor to hypercoagulation and thrombophilia in the metabolic syndrome. <i>Seminars in Thrombosis and Hemostasis</i> , 2009 , 35, 277-87	5.3	106
87	Nonalcoholic fatty liver disease is independently associated with an increased incidence of chronic kidney disease in patients with type 1 diabetes. <i>Diabetes Care</i> , 2014 , 37, 1729-36	14.6	98
86	Elevated serum uric acid concentrations independently predict cardiovascular mortality in type 2 diabetic patients. <i>Diabetes Care</i> , 2009 , 32, 1716-20	14.6	94
85	Mortality from chronic liver diseases in diabetes. <i>American Journal of Gastroenterology</i> , 2014 , 109, 1020-5	7	83
84	Non-alcoholic fatty liver disease is associated with an increased prevalence of atrial fibrillation in hospitalized patients with type 2 diabetes. <i>Clinical Science</i> , 2013 , 125, 301-9	6.5	80
83	Associations between liver histology and cortisol secretion in subjects with nonalcoholic fatty liver disease. <i>Clinical Endocrinology</i> , 2006 , 64, 337-41	3.4	67
82	Nonalcoholic Fatty Liver Disease Is Associated With Ventricular Arrhythmias in Patients With Type 2 Diabetes Referred for Clinically Indicated 24-Hour Holter Monitoring. <i>Diabetes Care</i> , 2016 , 39, 1416-23	14.6	66
81	Nonalcoholic Fatty Liver Disease Is Independently Associated with Early Left Ventricular Diastolic Dysfunction in Patients with Type 2 Diabetes. <i>PLoS ONE</i> , 2015 , 10, e0135329	3.7	61
80	Heart valve calcification in patients with type 2 diabetes and nonalcoholic fatty liver disease. <i>Metabolism: Clinical and Experimental</i> , 2015 , 64, 879-87	12.7	55
79	Association of nonalcoholic fatty liver disease with QTc interval in patients with type 2 diabetes. <i>Nutrition, Metabolism and Cardiovascular Diseases</i> , 2014 , 24, 663-9	4.5	55
78	Prevalence of neuropathy in type 2 diabetic patients and its association with other diabetes complications: The Verona Diabetic Foot Screening Program. <i>Journal of Diabetes and Its Complications</i> , 2015 , 29, 1066-70	3.2	53
77	Relation of elevated serum uric acid levels to incidence of atrial fibrillation in patients with type 2 diabetes mellitus. <i>American Journal of Cardiology</i> , 2013 , 112, 499-504	3	53
76	Prognostic Impact of Diabetes on Long-term Survival Outcomes in Patients With Heart Failure: A Meta-analysis. <i>Diabetes Care</i> , 2017 , 40, 1597-1605	14.6	52
75	Aortic and mitral annular calcifications are predictive of all-cause and cardiovascular mortality in patients with type 2 diabetes. <i>Diabetes Care</i> , 2012 , 35, 1781-6	14.6	52
74	Association Between Primary Hypothyroidism and Nonalcoholic Fatty Liver Disease: A Systematic Review and Meta-Analysis. <i>Thyroid</i> , 2018 , 28, 1270-1284	6.2	50

73	Variability of body weight, pulse pressure and glycaemia strongly predict total mortality in elderly type 2 diabetic patients. The Verona Diabetes Study. <i>Diabetes/Metabolism Research and Reviews</i> , 2008 , 24, 624-8	7.5	50
72	Association between nonalcoholic fatty liver disease and colorectal tumours in asymptomatic adults undergoing screening colonoscopy: a systematic review and meta-analysis. <i>Metabolism: Clinical and Experimental</i> , 2018 , 87, 1-12	12.7	48
71	Triglyceride-high-density lipoprotein cholesterol is associated with microvascular complications in type 2 diabetes mellitus. <i>Metabolism: Clinical and Experimental</i> , 2012 , 61, 22-9	12.7	45
70	Non-alcoholic fatty liver disease is independently associated with left ventricular hypertrophy in hypertensive Type 2 diabetic individuals. <i>Journal of Endocrinological Investigation</i> , 2012 , 35, 215-8	5.2	44
69	Association between serum TSH, free T4 and serum liver enzyme activities in a large cohort of unselected outpatients. <i>Clinical Endocrinology</i> , 2008 , 68, 481-4	3.4	44
68	Association between non-alcoholic fatty liver disease and risk of atrial fibrillation in adult individuals: An updated meta-analysis. <i>Liver International</i> , 2019 , 39, 758-769	7.9	43
67	Prevalence of Cardiovascular Autonomic Neuropathy in a Cohort of Patients With Newly Diagnosed Type 2 Diabetes: The Verona Newly Diagnosed Type 2 Diabetes Study (VNDS). <i>Diabetes Care</i> , 2015 , 38, 1487-93	14.6	42
66	Effect of moderate aerobic exercise on sympatho-vagal balance in Type 2 diabetic patients. <i>Diabetic Medicine</i> , 2007 , 24, 370-6	3.5	42
65	Nonalcoholic fatty liver disease is independently associated with an increased incidence of cardiovascular disease in adult patients with type 1 diabetes. <i>International Journal of Cardiology</i> , 2016 , 225, 387-391	3.2	39
64	Disorders of coagulation and hemostasis in abdominal obesity: emerging role of fatty liver. <i>Seminars in Thrombosis and Hemostasis</i> , 2010 , 36, 41-8	5.3	39
63	High-normal HbA1c is a strong predictor of type 2 diabetes in the general population. <i>Diabetes Care</i> , 2011 , 34, 1038-40	14.6	39
62	Glycated haemoglobin is inversely related to serum vitamin D levels in type 2 diabetic patients. <i>PLoS ONE</i> , 2013 , 8, e82733	3.7	36
61	Lower levels of 25-hydroxyvitamin D3 are associated with a higher prevalence of microvascular complications in patients with type 2 diabetes. <i>BMJ Open Diabetes Research and Care</i> , 2015 , 3, e000058	4.5	32
60	Prevalence of thyroid autoimmunity and subclinical hypothyroidism in persons with chronic kidney disease not requiring chronic dialysis. <i>Clinical Chemistry and Laboratory Medicine</i> , 2009 , 47, 1367-71	5.9	31
59	Nonalcoholic fatty liver disease is associated with an increased prevalence of distal symmetric polyneuropathy in adult patients with type 1 diabetes. <i>Journal of Diabetes and Its Complications</i> , 2017 , 31, 1021-1026	3.2	30
58	Nonalcoholic fatty liver disease is associated with an increased risk of heart block in hospitalized patients with type 2 diabetes mellitus. <i>PLoS ONE</i> , 2017 , 12, e0185459	3.7	29
57	Mortality from infectious diseases in diabetes. <i>Nutrition, Metabolism and Cardiovascular Diseases</i> , 2018 , 28, 444-450	4.5	28
56	Anaemia, independent of chronic kidney disease, predicts all-cause and cardiovascular mortality in type 2 diabetic patients. <i>Atherosclerosis</i> , 2010 , 210, 575-80	3.1	26

55	Nonalcoholic fatty liver disease and increased risk of 1-year all-cause and cardiac hospital readmissions in elderly patients admitted for acute heart failure. <i>PLoS ONE</i> , 2017 , 12, e0173398	3.7	26
54	Associations between liver histology and carotid intima-media thickness in patients with nonalcoholic fatty liver disease. <i>Arteriosclerosis, Thrombosis, and Vascular Biology</i> , 2005 , 25, 2687-8	9.4	25
53	Association between Helicobacter pylori infection and risk of nonalcoholic fatty liver disease: An updated meta-analysis. <i>Metabolism: Clinical and Experimental</i> , 2019 , 96, 56-65	12.7	24
52	Hypertriglyceridemia Is Independently Associated with Renal, but Not Retinal Complications in Subjects with Type 2 Diabetes: A Cross-Sectional Analysis of the Renal Insufficiency And Cardiovascular Events (RIACE) Italian Multicenter Study. <i>PLoS ONE</i> , 2015 , 10, e0125512	3.7	24
51	The role of serum uric acid in cardiovascular disease in type 2 diabetic and non-diabetic subjects: a narrative review. <i>Journal of Endocrinological Investigation</i> , 2011 , 34, 881-6	5.2	22
50	Relationship of nonalcoholic hepatic steatosis to overnight low-dose dexamethasone suppression test in obese individuals. <i>Clinical Endocrinology</i> , 2004 , 61, 711-5	3.4	22
49	Systematic review with meta-analysis: non-alcoholic fatty liver disease is associated with a history of osteoporotic fractures but not with low bone mineral density. <i>Alimentary Pharmacology and Therapeutics</i> , 2019 , 49, 375-388	6.1	22
48	Non-alcoholic fatty liver disease and increased risk of all-cause mortality in elderly patients admitted for acute heart failure. <i>International Journal of Cardiology</i> , 2018 , 265, 162-168	3.2	20
47	Usefulness of subclinical left ventricular midwall dysfunction to predict cardiovascular mortality in patients with type 2 diabetes mellitus. <i>American Journal of Cardiology</i> , 2014 , 113, 1409-14	3	20
46	Multiple causes of death analysis of chronic diseases: the example of diabetes. <i>Population Health Metrics</i> , 2015 , 13, 21	3	20
45	Comparison of two creatinine-based estimating equations in predicting all-cause and cardiovascular mortality in patients with type 2 diabetes. <i>Diabetes Care</i> , 2012 , 35, 2347-53	14.6	20
44	Early impairment in left ventricular longitudinal systolic function is associated with an increased risk of incident atrial fibrillation in patients with type 2 diabetes. <i>Journal of Diabetes and Its Complications</i> , 2017 , 31, 413-418	3.2	19
43	The aspartate aminotransferase-to-alanine aminotransferase ratio predicts all-cause and cardiovascular mortality in patients with type 2 diabetes. <i>Medicine (United States)</i> , 2016 , 95, e4821	1.8	19
42	Hemostatic disorders in type 1 diabetes mellitus. <i>Seminars in Thrombosis and Hemostasis</i> , 2011 , 37, 58-65	3	18
41	Relationship between serum bilirubin and kidney function in non-diabetic and diabetic individuals. <i>Kidney International</i> , 2009 , 75, 863	9.9	18
40	Association Between Nonalcoholic Fatty Liver Disease and Reduced Bone Mineral Density in Children: A Meta-Analysis. <i>Hepatology</i> , 2019 , 70, 812-823	11.2	16
39	Diabetes and cancer mortality: a multifaceted association. <i>Diabetes Research and Clinical Practice</i> , 2014 , 106, e86-9	7.4	16
38	Chronic complications in patients with newly diagnosed type 2 diabetes: prevalence and related metabolic and clinical features: the Verona Newly Diagnosed Type 2 Diabetes Study (VNDS) 9. <i>BMJ Open Diabetes Research and Care</i> , 2020 , 8,	4.5	16

37	Nonalcoholic Fatty Liver Disease Is Associated With Higher 1-year All-Cause Rehospitalization Rates in Patients Admitted for Acute Heart Failure. <i>Medicine (United States)</i> , 2016 , 95, e2760	1.8	15
36	Mitral Regurgitation and Increased Risk of All-Cause and Cardiovascular Mortality in Patients with Type 2 Diabetes. <i>American Journal of Medicine</i> , 2017 , 130, 70-76.e1	2.4	15
35	Usefulness of the triglyceride to high-density lipoprotein cholesterol ratio for predicting mortality risk in type 2 diabetes: role of kidney dysfunction. <i>Atherosclerosis</i> , 2010 , 212, 287-91	3.1	14
34	Relationship of serum gamma-glutamyltransferase to atherogenic dyslipidemia and glycemic control in type 2 diabetes. <i>Obesity</i> , 2009 , 17, 370-4	8	14
33	Prevalence of diabetes across different immigrant groups in North-eastern Italy. <i>Nutrition, Metabolism and Cardiovascular Diseases</i> , 2015 , 25, 924-30	4.5	13
32	Inappropriate left ventricular mass independently predicts cardiovascular mortality in patients with type 2 diabetes. <i>International Journal of Cardiology</i> , 2013 , 168, 4953-6	3.2	13
31	Hemostatic and fibrinolytic abnormalities in polycystic ovary syndrome. <i>Seminars in Thrombosis and Hemostasis</i> , 2014 , 40, 600-18	5.3	13
30	Evidence of left atrial remodeling and left ventricular diastolic dysfunction in type 2 diabetes mellitus with preserved systolic function. <i>Nutrition, Metabolism and Cardiovascular Diseases</i> , 2016 , 26, 1026-1032	4.5	11
29	Hemostatic and fibrinolytic abnormalities in endocrine diseases: a narrative review. <i>Seminars in Thrombosis and Hemostasis</i> , 2009 , 35, 605-12	5.3	10
28	Risk of all-cause and cardiovascular mortality in patients with chronic liver disease. <i>Gut</i> , 2011 , 60, 1602-3; author reply 1603-4	19.2	10
27	Relationship between increased left atrial volume and microvascular complications in patients with type 2 diabetes. <i>Journal of Diabetes and Its Complications</i> , 2015 , 29, 822-8	3.2	9
26	Soluble CD40L in young type 1 diabetic individuals without clinical microvascular and macrovascular complications. <i>Diabetes Care</i> , 2004 , 27, 1236-7	14.6	8
25	Independent correlates of urinary albumin excretion within the normoalbuminuric range in patients with type 2 diabetes: The Renal Insufficiency And Cardiovascular Events (RIACE) Italian Multicentre Study. <i>Acta Diabetologica</i> , 2015 , 52, 971-81	3.9	7
24	Pulse pressure and mortality from cerebrovascular diseases in type 2 diabetic patients: the Verona Diabetes Study. <i>Cerebrovascular Diseases</i> , 2007 , 23, 20-6	3.2	7
23	Association between subclinical left ventricular systolic dysfunction and glycemic control in asymptomatic type 2 diabetic patients with preserved left ventricular function. <i>Journal of Diabetes and Its Complications</i> , 2017 , 31, 1035-1040	3.2	6
22	Relation of elevated serum uric acid levels to first-degree heart block and other cardiac conduction defects in hospitalized patients with type 2 diabetes. <i>Journal of Diabetes and Its Complications</i> , 2017 , 31, 1691-1697	3.2	5
21	Effect of serum gamma-glutamyltransferase and obesity on the risk of dyslipidemia and poor glycemic control in type 2 diabetic patients: cross-sectional findings from the Verona Diabetes Study. <i>Clinical Chemistry</i> , 2007 , 53, 1867-9; author reply 1869-70	5.5	5
20	Long-Acting GLP-1 Receptor Agonist Exenatide Influence on the Autonomic Cardiac Sympatho-Vagal Balance. <i>Journal of the Endocrine Society</i> , 2018 , 2, 53-62	0.4	4

19	Severe hypoglycemia in patients with known diabetes requiring emergency department care: A report from an Italian multicenter study. <i>Journal of Clinical and Translational Endocrinology</i> , 2016 , 5, 46-52 ⁴	4
18	The E/eSratio difference between subjects with type 2 diabetes and controls. A meta-analysis of clinical studies. <i>PLoS ONE</i> , 2018 , 13, e0209794	3.7 4
17	Increased aortic stiffness index in patients with type 1 diabetes without cardiovascular disease compared to controls. <i>Journal of Endocrinological Investigation</i> , 2019 , 42, 1109-1115	5.2 3
16	Nonalcoholic Fatty Liver Disease and Implications for Older Adults with Diabetes. <i>Clinics in Geriatric Medicine</i> , 2020 , 36, 527-547	3.8 3
15	Left ventricular chamber dilation and filling pressure may help to categorise patients with type 2 diabetes. <i>BMJ Open Diabetes Research and Care</i> , 2018 , 6, e000529	4.5 2
14	Time series of diabetes attributable mortality from 2008 to 2017. <i>Journal of Endocrinological Investigation</i> , 2021 , 1	5.2 2
13	Estimating the real burden of cardiovascular mortality in diabetes. <i>European Review for Medical and Pharmacological Sciences</i> , 2019 , 23, 6700-6706	2.9 2
12	Impact of reference category and number of traits in the cluster on risk of coronary heart disease in metabolic syndrome: prospective data from the Bruneck study. <i>Metabolic Syndrome and Related Disorders</i> , 2011 , 9, 313-8	2.6 1
11	Relationship between soluble CD40 ligand and gamma-glutamyltransferase concentrations in non-drinking, young Type 1 diabetic individuals. <i>Diabetic Medicine</i> , 2008 , 25, 1283-8	3.5 1
10	Glomerular filtration rate decline in T2DM following diagnosis. The Verona newly diagnosed diabetes study-12. <i>Diabetes Research and Clinical Practice</i> , 2021 , 175, 108778	7.4 1
9	SARS-CoV-2 and COVID-19 in diabetes mellitus. Population-based study on ascertained infections, hospital admissions and mortality in an Italian region with ~5 million inhabitants and ~250,000 diabetic people. <i>Nutrition, Metabolism and Cardiovascular Diseases</i> , 2021 , 31, 2612-2618	4.5 1
8	Insulin effect on serum potassium and auto-inhibition of insulin secretion is intact in a patient with leprechaunism despite severe impairment of substrates metabolism. <i>Diabetes/Metabolism Research and Reviews</i> , 2008 , 24, 205-10	7.5 0
7	A renal genetic risk score (GRS) is associated with kidney dysfunction in people with type 2 diabetes. <i>Diabetes Research and Clinical Practice</i> , 2018 , 144, 137-143	7.4
6	Echocardiographic parameters according to insulin dose in young patients affected by type 1 diabetes. <i>PLoS ONE</i> , 2020 , 15, e0244483	3.7
5	Echocardiographic parameters according to insulin dose in young patients affected by type 1 diabetes 2020 , 15, e0244483	
4	Echocardiographic parameters according to insulin dose in young patients affected by type 1 diabetes 2020 , 15, e0244483	
3	Echocardiographic parameters according to insulin dose in young patients affected by type 1 diabetes 2020 , 15, e0244483	
2	Echocardiographic parameters according to insulin dose in young patients affected by type 1 diabetes 2020 , 15, e0244483	

- 1 Estimated peak systolic pulmonary artery pressure in young non-complicated patients with type 1 diabetes. *European Review for Medical and Pharmacological Sciences*, **2020**, 24, 5028-5035 2.9