

Bettina Tonn

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/7701122/publications.pdf>

Version: 2024-02-01

27
papers

343
citations

759233

12
h-index

839539

18
g-index

27
all docs

27
docs citations

27
times ranked

405
citing authors

#	ARTICLE	IF	CITATIONS
1	Leaching of biomass from semi-natural grasslands – Effects on chemical composition and ash high-temperature behaviour. <i>Biomass and Bioenergy</i> , 2012, 36, 390-403.	5.7	44
2	Semi-natural grassland biomass for combustion: influence of botanical composition, harvest date and site conditions on fuel composition. <i>Grass and Forage Science</i> , 2010, 65, 383-397.	2.9	36
3	Soil pH and phosphorus drive species composition and richness in semi-natural heathlands and grasslands unaffected by twentieth-century agricultural intensification. <i>Plant Ecology and Diversity</i> , 2018, 11, 239-253.	2.4	24
4	How German dairy farmers perceive advantages and disadvantages of grazing and how it relates to their milk production systems. <i>Livestock Science</i> , 2018, 214, 112-119.	1.6	22
5	Grazing-induced patchiness, not grazing intensity, drives plant diversity in European low-input pastures. <i>Journal of Applied Ecology</i> , 2019, 56, 1624-1636.	4.0	21
6	Target-oriented habitat and wildlife management: estimating forage quantity and quality of semi-natural grasslands with Sentinel-1 and Sentinel-2 data. <i>Remote Sensing in Ecology and Conservation</i> , 2020, 6, 381-398.	4.3	21
7	Influence of leaching on the chemical composition of grassland biomass for combustion. <i>Grass and Forage Science</i> , 2011, 66, 464-473.	2.9	20
8	Mapping semi-natural grassland communities using multi-temporal RapidEye remote sensing data. <i>International Journal of Remote Sensing</i> , 2018, 39, 5638-5659.	2.9	18
9	Sward patterns created by patch grazing are stable over more than a decade. <i>Grass and Forage Science</i> , 2019, 74, 104-114.	2.9	16
10	Grazing by wild red deer maintains characteristic vegetation of semi-natural open habitats: Evidence from a three-year exclusion experiment. <i>Applied Vegetation Science</i> , 2020, 23, 522-538.	1.9	16
11	Tall wheatgrass (<i>Agropyron elongatum</i>) for biogas production: Crop management more important for biomass and methane yield than grass provenance. <i>Industrial Crops and Products</i> , 2017, 97, 653-663.	5.2	14
12	Grazing by wild red deer: Management options for the conservation of semi-natural open habitats. <i>Journal of Applied Ecology</i> , 2019, 56, 1311-1321.	4.0	13
13	White clover population effects on the productivity and yield stability of mixtures with perennial ryegrass and chicory. <i>Field Crops Research</i> , 2020, 252, 107802.	5.1	10
14	The Effect of Grazing Intensity and Sward Heterogeneity on the Movement Behavior of Suckler Cows on Semi-natural Grassland. <i>Frontiers in Veterinary Science</i> , 2021, 8, 639096.	2.2	10
15	Herbage biomass and uptake under low-input grazing as affected by cattle and sheep excrement patches. <i>Nutrient Cycling in Agroecosystems</i> , 2018, 112, 277-289.	2.2	9
16	Effect of grazing intensity and soil characteristics on soil organic carbon and nitrogen stocks in a temperate long-term grassland. <i>Archives of Agronomy and Soil Science</i> , 2017, 63, 1776-1783.	2.6	8
17	Primary productivity in patches of heterogeneous swards after 12 years of low-intensity cattle grazing. <i>Grass and Forage Science</i> , 2020, 75, 398-408.	2.9	8
18	Urine effects on grass and legume nitrogen isotopic composition: Pronounced short-term dynamics of $\delta^{15}N$. <i>PLoS ONE</i> , 2019, 14, e0210623.	2.5	7

#	ARTICLE	IF	CITATIONS
19	Using a Citizen Science Approach with German Horse Owners to Study the Locomotion Behaviour of Horses on Pasture. Sustainability, 2020, 12, 1835.	3.2	7
20	Impacts of cutting frequency and position to tree line on herbage accumulation in silvopastoral grassland reveal potential for grassland conservation based on land use and cover information. Annals of Applied Biology, 2021, 179, 75-84.	2.5	6
21	Multi-temporal RapidEye Tasselled Cap data for land cover classification. European Journal of Remote Sensing, 2019, 52, 653-666.	3.5	3
22	The Role of Socio-Economic Determinants of Horse Farms for Grassland Management, Vegetation Composition and Ecological Value. Sustainability, 2020, 12, 10641.	3.2	3
23	Seasonal plasticity is more important than population variability in effects on white clover architecture and productivity. Annals of Botany, 2021, 128, 73-82.	2.9	3
24	Grazing by wild red deer can mitigate nutrient enrichment in protected semi-natural open habitats. Oecologia, 2022, 199, 471-485.	2.0	2
25	Results from a biodiversity experiment fail to represent economic performance of semi-natural grasslands. Nature Communications, 2021, 12, 2125.	12.8	1
26	Including chicory and selecting white clover varieties as strategies to improve temporal stability of forage yield and quality in white-clover-based temporary grassland. European Journal of Agronomy, 2021, 130, 126362.	4.1	1
27	Effects of drought-stressed temperate forage legumes on the degradation and the rumen microbial community in vitro. Journal of Animal Physiology and Animal Nutrition, 2019, 103, 436-446.	2.2	0