Marius Henriksen

List of Publications by Citations

Source: https://exaly.com/author-pdf/7703564/marius-henriksen-publications-by-citations.pdf

Version: 2024-04-28

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

126 papers

2,818 citations

29 h-index

g-index

141 ext. papers

3,365 ext. citations

3.5 avg, IF

5.04 L-index

#	Paper	IF	Citations
126	Test-retest reliability of trunk accelerometric gait analysis. <i>Gait and Posture</i> , 2004 , 19, 288-97	2.6	201
125	Effect of whole body vibration exercise on muscle strength and proprioception in females with knee osteoarthritis. <i>Knee</i> , 2009 , 16, 256-61	2.6	122
124	Gait changes in patients with knee osteoarthritis are replicated by experimental knee pain. <i>Arthritis Care and Research</i> , 2010 , 62, 501-9	4.7	107
123	Effects of an intensive weight loss program on knee joint loading in obese adults with knee osteoarthritis. <i>Osteoarthritis and Cartilage</i> , 2011 , 19, 822-8	6.2	102
122	The role of muscle strengthening in exercise therapy for knee osteoarthritis: A systematic review and meta-regression analysis of randomized trials. <i>Seminars in Arthritis and Rheumatism</i> , 2017 , 47, 9-21	5.3	97
121	Experimental knee pain reduces muscle strength. <i>Journal of Pain</i> , 2011 , 12, 460-7	5.2	94
120	The concept of physical limitations in knee osteoarthritis I the view of patients and health professionals. <i>Trials</i> , 2015 , 16,	2.8	78
119	Increased joint loads during walkinga consequence of pain relief in knee osteoarthritis. <i>Knee</i> , 2006 , 13, 445-50	2.6	76
118	Weight loss is effective for symptomatic relief in obese subjects with knee osteoarthritis independently of joint damage severity assessed by high-field MRI and radiography. <i>Osteoarthritis and Cartilage</i> , 2012 , 20, 495-502	6.2	75
117	Experimental quadriceps muscle pain impairs knee joint control during walking. <i>Journal of Applied Physiology</i> , 2007 , 103, 132-9	3.7	74
116	Knee pain and inflammation in the infrapatellar fat pad estimated by conventional and dynamic contrast-enhanced magnetic resonance imaging in obese patients with osteoarthritis: a cross-sectional study. <i>Osteoarthritis and Cartilage</i> , 2014 , 22, 933-40	6.2	71
115	Weight loss for overweight and obese individuals with gout: a systematic review of longitudinal studies. <i>Annals of the Rheumatic Diseases</i> , 2017 , 76, 1870-1882	2.4	67
114	Association of exercise therapy and reduction of pain sensitivity in patients with knee osteoarthritis: a randomized controlled trial. <i>Arthritis Care and Research</i> , 2014 , 66, 1836-43	4.7	66
113	Effect of weight maintenance on symptoms of knee osteoarthritis in obese patients: a twelve-month randomized controlled trial. <i>Arthritis Care and Research</i> , 2015 , 67, 640-50	4.7	64
112	Walking on high heels changes muscle activity and the dynamics of human walking significantly. Journal of Applied Biomechanics, 2012 , 28, 20-8	1.2	60
111	Evaluation of the benefit of corticosteroid injection before exercise therapy in patients with osteoarthritis of the knee: a randomized clinical trial. <i>JAMA Internal Medicine</i> , 2015 , 175, 923-30	11.5	58
110	Changes in lower extremity muscle mass and muscle strength after weight loss in obese patients with knee osteoarthritis: a prospective cohort study. <i>Arthritis and Rheumatism</i> , 2012 , 64, 438-42		52

(2014-2009)

109	Experimentally reduced hip abductor function during walking: Implications for knee joint loads. <i>Journal of Biomechanics</i> , 2009 , 42, 1236-40	2.9	50	
108	Relationships between the fibromyalgia impact questionnaire, tender point count, and muscle strength in female patients with fibromyalgia: a cohort study. <i>Arthritis and Rheumatism</i> , 2009 , 61, 732-9		45	
107	The relationship between pain and dynamic knee joint loading in knee osteoarthritis varies with radiographic disease severity. A cross sectional study. <i>Knee</i> , 2012 , 19, 392-8	2.6	44	
106	Ultrasound colour Doppler measurements in a single joint as measure of disease activity in patients with rheumatoid arthritisassessment of concurrent validity. <i>Rheumatology</i> , 2009 , 48, 254-7	3.9	42	
105	Osteoarthritis year in review 2016: imaging. Osteoarthritis and Cartilage, 2017, 25, 216-226	6.2	36	
104	Enhancing the reporting and transparency of rheumatology research: a guide to reporting guidelines. <i>Arthritis Research and Therapy</i> , 2013 , 15, 109	5.7	34	
103	Synovitis assessed on static and dynamic contrast-enhanced magnetic resonance imaging and its association with pain in knee osteoarthritis: A cross-sectional study. <i>European Journal of Radiology</i> , 2016 , 85, 1099-108	4.7	34	
102	Nordic Walking does not reduce the loading of the knee joint. <i>Scandinavian Journal of Medicine and Science in Sports</i> , 2008 , 18, 436-41	4.6	33	
101	Biomechanical characteristics of the eccentric Achilles tendon exercise. <i>Journal of Biomechanics</i> , 2009 , 42, 2702-7	2.9	32	
100	Long-term weight-loss maintenance in obese patients with knee osteoarthritis: a randomized trial. <i>American Journal of Clinical Nutrition</i> , 2017 , 106, 755-763	7	31	
99	Different knee joint loading patterns in ACL deficient copers and non-copers during walking. <i>Knee Surgery, Sports Traumatology, Arthroscopy</i> , 2011 , 19, 615-21	5.5	31	
98	Is there a causal link between knee loading and knee osteoarthritis progression? A systematic review and meta-analysis of cohort studies and randomised trials. <i>BMJ Open</i> , 2014 , 4, e005368	3	30	
97	The risk associated with spinal manipulation: an overview of reviews. Systematic Reviews, 2017, 6, 64	3	29	
96	Association of different levels of depressive symptoms with symptomatology, overall disease severity, and quality of life in women with fibromyalgia. <i>Quality of Life Research</i> , 2015 , 24, 2951-7	3.7	28	
95	Comparison of three weight maintenance programs on cardiovascular risk, bone and vitamins in sedentary older adults. <i>Obesity</i> , 2013 , 21, 1982-90	8	28	
94	Gait variability and motor control in people with knee osteoarthritis. <i>Gait and Posture</i> , 2015 , 42, 479-84	2.6	27	
93	Experimental muscle pain during a forward lungethe effects on knee joint dynamics and electromyographic activity. <i>British Journal of Sports Medicine</i> , 2009 , 43, 503-7	10.3	27	
92	Effect of a 16 weeks weight loss program on osteoarthritis biomarkers in obese patients with knee osteoarthritis: a prospective cohort study. <i>Osteoarthritis and Cartilage</i> , 2014 , 22, 1817-25	6.2	25	

91	Association of Physical Fitness With Fibromyalgia Severity in Women: The al-Edalus Project. <i>Archives of Physical Medicine and Rehabilitation</i> , 2015 , 96, 1599-605	2.8	24
90	Altered visual and feet proprioceptive feedbacks during quiet standing increase postural sway in patients with severe knee osteoarthritis. <i>PLoS ONE</i> , 2013 , 8, e71253	3.7	24
89	Poor replicability of recommended exercise interventions for knee osteoarthritis: a descriptive analysis of evidence informing current guidelines and recommendations. <i>Osteoarthritis and Cartilage</i> , 2019 , 27, 3-22	6.2	24
88	The association between histological, macroscopic and magnetic resonance imaging assessed synovitis in end-stage knee osteoarthritis: a cross-sectional study. <i>Osteoarthritis and Cartilage</i> , 2017 , 25, 272-280	6.2	23
87	Muscle fatigue in fibromyalgia is in the brain, not in the muscles: a case-control study of perceived versus objective muscle fatigue. <i>Annals of the Rheumatic Diseases</i> , 2013 , 72, 963-6	2.4	23
86	Preparing for what the reporting checklists will not tell you: the PREPARE Trial guide for planning clinical research to avoid research waste. <i>British Journal of Sports Medicine</i> , 2017 , 51, 1494-1501	10.3	22
85	Effect of aerobic exercise training on asthma in adults: a systematic review and meta-analysis. <i>European Respiratory Journal</i> , 2020 , 56,	13.6	21
84	Structural changes in the knee during weight loss maintenance after a significant weight loss in obese patients with osteoarthritis: a report of secondary outcome analyses from a randomized controlled trial. Osteoarthritis and Cartilage, 2014, 22, 639-46	6.2	21
83	Impulse-forces during walking are not increased in patients with knee osteoarthritis. <i>Monthly Notices of the Royal Astronomical Society: Letters</i> , 2006 , 77, 650-6	4.3	21
82	Quantification of colour Doppler activity in the wrist in patients with rheumatoid arthritisthe reliability of different methods for image selection and evaluation. <i>Ultraschall in Der Medizin</i> , 2008 , 29, 393-8	3.8	20
81	Forward lunge as a functional performance test in ACL deficient subjects: test-retest reliability. <i>Knee</i> , 2009 , 16, 176-82	2.6	19
80	Motor responses to experimental Achilles tendon pain. <i>British Journal of Sports Medicine</i> , 2011 , 45, 393	- 8 10.3	19
79	Comparable effects of exercise and analgesics for pain secondary to knee osteoarthritis: a meta-analysis of trials included in Cochrane systematic reviews. <i>Journal of Comparative Effectiveness Research</i> , 2016 , 5, 417-31	2.1	19
78	Quadriceps-strengthening exercise and quadriceps and knee biomechanics during walking in knee osteoarthritis: A two-centre randomized controlled trial. <i>Clinical Biomechanics</i> , 2018 , 59, 199-206	2.2	18
77	Association of Physical Fitness with Depression in Women with Fibromyalgia. <i>Pain Medicine</i> , 2016 , 17, 1542-52	2.8	17
76	Is increased joint loading detrimental to obese patients with knee osteoarthritis? A secondary data analysis from a randomized trial. <i>Osteoarthritis and Cartilage</i> , 2013 , 21, 1865-75	6.2	17
75	Influence of recent exercise and skin temperature on ultrasound Doppler measurements in patients with rheumatoid arthritisan intervention study. <i>Rheumatology</i> , 2009 , 48, 1520-3	3.9	17
74	Intra-Articular Corticosteroids in Addition to Exercise for Reducing Pain Sensitivity in Knee Osteoarthritis: Exploratory Outcome from a Randomized Controlled Trial. <i>PLoS ONE</i> , 2016 , 11, e014916	58 ^{.7}	14

(2018-2016)

73	A Standardized "Rescue" Exercise Program for Symptomatic Flare-up of Knee Osteoarthritis: Description and Safety Considerations. <i>Journal of Orthopaedic and Sports Physical Therapy</i> , 2016 , 46, 942-946	4.2	14
72	Changes in bone marrow lesions in response to weight-loss in obese knee osteoarthritis patients: a prospective cohort study. <i>BMC Musculoskeletal Disorders</i> , 2013 , 14, 106	2.8	13
71	The effects of intra-articular glucocorticoids and exercise on pain and synovitis assessed on static and dynamic magnetic resonance imaging in knee osteoarthritis: exploratory outcomes from a randomized controlled trial. <i>Osteoarthritis and Cartilage</i> , 2017 , 25, 481-491	6.2	13
70	Associations between muscle perfusion and symptoms in knee osteoarthritis: a cross sectional study. <i>Osteoarthritis and Cartilage</i> , 2015 , 23, 1721-7	6.2	12
69	Exercise therapy after ultrasound-guided corticosteroid injections in patients with subacromial pain syndrome: a randomized controlled trial. <i>Arthritis Research and Therapy</i> , 2016 , 18, 129	5.7	12
68	Kinesthetic taping improves walking function in patients with stroke: a pilot cohort study. <i>Topics in Stroke Rehabilitation</i> , 2014 , 21, 495-501	2.6	11
67	Experimental knee pain evoke spreading hyperalgesia and facilitated temporal summation of pain. <i>Pain Medicine</i> , 2013 , 14, 874-83	2.8	11
66	Influence of pain and gender on impact loading during walking: a randomised trial. <i>Clinical Biomechanics</i> , 2008 , 23, 221-30	2.2	11
65	Physiotherapy for pain: a meta-epidemiological study of randomised trials. <i>British Journal of Sports Medicine</i> , 2016 , 50, 965-71	10.3	11
64	Hand exercise for women with rheumatoid arthritis and decreased hand function: an exploratory randomized controlled trial. <i>Arthritis Research and Therapy</i> , 2019 , 21, 158	5.7	10
63	The effect of foot progression angle on knee joint compression force during walking. <i>Journal of Applied Biomechanics</i> , 2013 , 29, 329-35	1.2	10
62	Is it possible to reduce the knee joint compression force during level walking with hiking poles?. <i>Scandinavian Journal of Medicine and Science in Sports</i> , 2011 , 21, e195-200	4.6	10
61	Can stimulating massage improve joint repositioning error in patients with knee osteoarthritis?. <i>Journal of Geriatric Physical Therapy</i> , 2009 , 32, 111-6	3.2	10
60	Changes in ultrasound assessed markers of inflammation following intra-articular steroid injection combined with exercise in knee osteoarthritis: exploratory outcome from a randomized trial. Osteoarthritis and Cartilage, 2016, 24, 814-21	6.2	10
59	Evidence-Based Research Series-Paper 2 : Using an Evidence-Based Research approach before a new study is conducted to ensure value. <i>Journal of Clinical Epidemiology</i> , 2021 , 129, 158-166	5.7	10
58	The effect of isometric exercise of the hand on the synovial blood flow in patients with rheumatoid arthritis measured by color Doppler ultrasound. <i>Rheumatology International</i> , 2013 , 33, 65-70	3.6	9
57	Liraglutide after diet-induced weight loss for pain and weight control in knee osteoarthritis: a randomized controlled trial. <i>American Journal of Clinical Nutrition</i> , 2021 , 113, 314-323	7	9
56	Progressive early passive and active exercise therapy after surgical rotator cuff repair - study protocol for a randomized controlled trial (the CUT-N-MOVE trial). <i>Trials</i> , 2018 , 19, 470	2.8	9

55	Reliability and Construct Validity of the SENS Motion [©] Activity Measurement System as a Tool to Detect Sedentary Behaviour in Patients with Knee Osteoarthritis. <i>Arthritis</i> , 2018 , 2018, 6596278		9
54	Relationship between weight loss in obese knee osteoarthritis patients and serum biomarkers of cartilage breakdown: secondary analyses of a randomised trial. <i>Osteoarthritis and Cartilage</i> , 2017 , 25, 1641-1646	6.2	8
53	Intra-Articular Analgesia and Steroid Reduce Pain Sensitivity in Knee OA Patients: An Interventional Cohort Study. <i>Pain Research and Treatment</i> , 2014 , 2014, 710490	1.9	8
52	Adaptations in the gait pattern with experimental hamstring pain. <i>Journal of Electromyography and Kinesiology</i> , 2011 , 21, 746-53	2.5	8
51	The association between submaximal quadriceps force steadiness and the knee adduction moment during walking in patients with knee osteoarthritis. <i>Journal of Orthopaedic and Sports Physical Therapy</i> , 2011 , 41, 592-9	4.2	8
50	Association Between Weight Loss and Spontaneous Changes in Physical Inactivity in Overweight/Obese Individuals With Knee Osteoarthritis: An Eight-Week Prospective Cohort Study. <i>Arthritis Care and Research</i> , 2020 , 72, 397-404	4.7	8
49	The effect of stimulating massage of thigh muscles on knee joint position sense. <i>Advances in Physiotherapy</i> , 2004 , 6, 29-36		7
48	No effects of functional exercise therapy on walking biomechanics in patients with knee osteoarthritis: exploratory outcome analyses from a randomised trial. <i>BMJ Open Sport and Exercise Medicine</i> , 2016 , 2, e000230	3.4	6
47	The concept of physical limitations in knee osteoarthritis: as viewed by patients and health professionals. <i>Quality of Life Research</i> , 2015 , 24, 2423-32	3.7	6
46	Experimental knee joint pain during strength training and muscle strength gain in healthy subjects: a randomized controlled trial. <i>Arthritis Care and Research</i> , 2012 , 64, 108-16	4.7	6
45	The dynamics of the pain system is intact in patients with knee osteoarthritis: An exploratory experimental study. <i>Scandinavian Journal of Pain</i> , 2015 , 6, 43-49	1.9	6
44	The Influence of Radiographic Severity on the Relationship between Muscle Strength and Joint Loading in Obese Knee Osteoarthritis Patients. <i>Arthritis</i> , 2011 , 2011, 571519		6
43	Improved gait in persons with knee related mobility limitations by a rosehip food supplement: A randomized, double-blind, placebo-controlled trial. <i>Gait and Posture</i> , 2015 , 42, 340-7	2.6	5
42	The effect of intra-articular glucocorticosteroids and exercise on symptoms and bone marrow lesions in knee osteoarthritis: a secondary analysis of results from a randomized controlled trial. <i>Osteoarthritis and Cartilage</i> , 2018 , 26, 895-902	6.2	5
41	Osteoarthritis: Time to put steroid injections behind us?. <i>Nature Reviews Rheumatology</i> , 2017 , 13, 519-5	280 1	5
40	The Associations between Pain Sensitivity and Knee Muscle Strength in Healthy Volunteers: A Cross-Sectional Study. <i>Pain Research and Treatment</i> , 2013 , 2013, 787054	1.9	5
39	Increased mortality in patients with severe COPD associated with high-intensity exercise: a preliminary cohort study. <i>International Journal of COPD</i> , 2016 , 11, 2329-2334	3	5
38	Evidence-Based Research Series-Paper 3: Using an Evidence-Based Research approach to place your results into context after the study is performed to ensure usefulness of the conclusion. <i>Journal of Clinical Epidemiology</i> , 2021 , 129, 167-171	5.7	5

(2019-2014)

37	Comparing different preparations and doses of rosehip powder in patients with osteoarthritis of the knee: an exploratory randomized active-controlled trial. <i>International Journal of Clinical Rheumatology</i> , 2014 , 9, 267-278	1.5	4	
36	Perfusion in bone marrow lesions assessed on DCE-MRI and its association with pain in knee osteoarthritis: a cross-sectional study. <i>Skeletal Radiology</i> , 2020 , 49, 757-764	2.7	4	
35	The effect of exercise therapy on inflammatory activity assessed by MRI in knee osteoarthritis: Secondary outcomes from a randomized controlled trial. <i>Knee</i> , 2021 , 28, 256-265	2.6	4	
34	Effectiveness of text messages for decreasing inactive behaviour in patients with knee osteoarthritis: a pilot randomised controlled study. <i>Pilot and Feasibility Studies</i> , 2019 , 5, 112	1.9	3	
33	Effect of liraglutide on body weight and pain in patients with overweight and knee osteoarthritis: protocol for a randomised, double-blind, placebo-controlled, parallel-group, single-centre trial. <i>BMJ Open</i> , 2019 , 9, e024065	3	3	
32	Dynamic weight-bearing assessment of pain in knee osteoarthritis: a reliability and agreement study. <i>Quality of Life Research</i> , 2015 , 24, 2985-92	3.7	3	
31	Validity and reliability of 3D US for the detection of erosions in patients with rheumatoid arthritis using MRI as the gold standard. <i>Ultraschall in Der Medizin</i> , 2014 , 35, 137-41	3.8	3	
30	Obesity and Walking: Implications for Knee Osteoarthritis and Plantar Heel Pain. <i>Current Obesity Reports</i> , 2012 , 1, 160-165	8.4	3	
29	Dynamic control of the lumbopelvic complex; lack of reliability of established test procedures. <i>European Spine Journal</i> , 2007 , 16, 733-40	2.7	3	
28	Placebo response and effect in randomized clinical trials: meta-research with focus on contextual effects. <i>Trials</i> , 2021 , 22, 493	2.8	3	
27	Local and Systemic Changes in Pain Sensitivity After 4 Weeks of Calf Muscle Stretching in a Nonpainful Population: A Randomized Trial. <i>Pain Practice</i> , 2016 , 16, 696-703	3	3	
26	Controversy and Debate on Meta-epidemiology. Paper 1: Treatment effect sizes vary in randomized trials depending on the type of outcome measure. <i>Journal of Clinical Epidemiology</i> , 2020 , 123, 27-38	5.7	2	
25	Exercise-induced pain changes associate with changes in muscle perfusion in knee osteoarthritis: exploratory outcome analyses of a randomised controlled trial. <i>BMC Musculoskeletal Disorders</i> , 2019 , 20, 491	2.8	2	
24	Visual assessment of dynamic knee joint alignment in patients with patellofemoral pain: an agreement study. <i>PeerJ</i> , 2021 , 9, e12203	3.1	2	
23	Changes in physical inactivity during supervised educational and exercise therapy in patients with knee osteoarthritis: A prospective cohort study. <i>Knee</i> , 2020 , 27, 1848-1856	2.6	2	
22	Hand-related physical function in rheumatic hand conditions: a protocol for developing a patient-reported outcome measurement instrument. <i>BMJ Open</i> , 2016 , 6, e011174	3	2	
21	Prevalence of self-reported knee symptoms and management strategies among elderly individuals from Frederiksberg municipality: protocol for a prospective and pragmatic Danish cohort study. <i>BMJ Open</i> , 2019 , 9, e028087	3	2	
20	Opioid-Induced Reductions in Gait Variability in Healthy Volunteers and Individuals with Knee Osteoarthritis. <i>Pain Medicine</i> , 2019 , 20, 2106-2114	2.8	2	

19	Associations between shoulder symptoms and concomitant pathology în patients with traumatic supraspinatus tears. <i>JSES International</i> , 2020 , 4, 85-90	1.2	2
18	Estimating the Prevalence of Knee Pain and the Association between Illness Perception Profiles and Self-Management Strategies in the Frederiksberg Cohort of Elderly Individuals with Knee Pain: A Cross-Sectional Study. <i>Journal of Clinical Medicine</i> , 2021 , 10,	5.1	2
17	Evaluation of a technology assisted physical activity intervention among hospitalised patients: A randomised study. <i>European Journal of Internal Medicine</i> , 2019 , 69, 50-56	3.9	1
16	The effect of graded activity and pain education (GAPE): an early post-surgical rehabilitation programme after lumbar spinal fusion-study protocol for a randomized controlled trial. <i>Trials</i> , 2020 , 21, 791	2.8	1
15	PEP-CoV protocol: a PEP flute-self-care randomised controlled trial to prevent respiratory deterioration and hospitalisation in early COVID-19. <i>BMJ Open</i> , 2021 , 11, e050582	3	1
14	Dynamic weight-bearing assessment of pain in knee osteoarthritis: construct validity, responsiveness, and interpretability in a research setting. <i>Health and Quality of Life Outcomes</i> , 2016 , 14, 91	3	1
13	Exercise therapy and patient education versus intra-articular saline injections in the treatment of knee osteoarthritis: an evidence-based protocol for an open-label randomised controlled trial (the DISCO trial). <i>Trials</i> , 2021 , 22, 18	2.8	1
12	Effects of 12 Weeks of Progressive Early Active Exercise Therapy After Surgical Rotator Cuff Repair: 12 Weeks and 1-Year Results From the CUT-N-MOVE Randomized Controlled Trial. <i>American Journal of Sports Medicine</i> , 2021 , 49, 321-331	6.8	1
11	Knee replacement outcome predicted by physiotherapists: a prospective cohort study. <i>PeerJ</i> , 2021 , 9, e10838	3.1	1
10	Effects of a 12-week supervised resistance training program, combined with home-based physical activity, on physical fitness and quality of life in female breast cancer survivors: the EFICAN randomized controlled trial <i>Journal of Cancer Survivorship</i> , 2022 ,	5.1	1
9	Triage Strategies Based on C-Reactive Protein Levels and SARS-CoV-2 Tests among Individuals Referred with Suspected COVID-19: A Prospective Cohort Study <i>Journal of Clinical Medicine</i> , 2021 , 11,	5.1	1
8	Effect of PEP flute self-care versus usual care in early covid-19: non-drug, open label, randomised controlled trial in a Danish community setting. <i>BMJ, The</i> , 2021 , 375, e066952	5.9	O
7	OP0172 EFFECT OF WEIGHT LOSS AND LIRAGLUTIDE ON SERUM URATE LEVELS AMONG OBESE KNEE OSTEOARTHRITIS PATIENTS: SECONDARY ANALYSIS OF A RANDOMISED CONTROLLED TRIAL. <i>Annals of the Rheumatic Diseases</i> , 2020 , 79, 107.1-108	2.4	О
6	THU0619-HPR PREVALENCE OF DISTAL INTERPHALANGEAL JOINT ULTRASONOGRAPHY FEATURES IN PSORIATIC ARTHRITIS, SKIN PSORIASIS, OSTEOARTHRITIS AND HEALTHY INDIVIDUALS: A CROSS-SECTIONAL STUDY. <i>Annals of the Rheumatic Diseases</i> , 2020 , 79, 552-553	2.4	
5	Test-retest reliability of cuff pressure pain algometry in patients with knee osteoarthritis. <i>Clinical and Experimental Rheumatology</i> , 2016 , 34, 158	2.2	
4	Intra-articular 2.5% polyacrylamide hydrogel for the treatment of knee osteoarthritis: an observational proof-of-concept cohort study. <i>Clinical and Experimental Rheumatology</i> , 2018 , 36, 1082-1	085	
3	PhysiotherapistsPprognosis of 1-year outcome after lumbar spinal fusion - A prospective cohort study <i>Physiotherapy Theory and Practice</i> , 2022 , 1-12	1.5	
2	Changes in physical activity during a one-year weight loss trial with liraglutide vs placebo in participants with knee osteoarthritis: Secondary analyses of a randomised controlled trial. Osteoarthritis and Cartilage Open, 2022, 4, 100255	1.5	

LIST OF PUBLICATIONS

Prognostic factors for work disability in patients with chronic widespread pain and fibromyalgia: protocol for a cohort study.. *BMJ Open*, **2021**, 11, e052919

3