

Haizhao Song

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/7703569/publications.pdf>

Version: 2024-02-01

18
papers

518
citations

759055

12
h-index

839398

18
g-index

18
all docs

18
docs citations

18
times ranked

616
citing authors

#	ARTICLE	IF	CITATIONS
1	Red pitaya betacyanins protects from diet-induced obesity, liver steatosis and insulin resistance in association with modulation of gut microbiota in mice. <i>Journal of Gastroenterology and Hepatology (Australia)</i> , 2016, 31, 1462-1469.	1.4	101
2	Purified Betacyanins from <i>Hylocereus undatus</i> Peel Ameliorate Obesity and Insulin Resistance in High-Fat-Diet-Fed Mice. <i>Journal of Agricultural and Food Chemistry</i> , 2016, 64, 236-244.	2.4	57
3	CaCO ₃ nanoparticles incorporated with KAE to enable amplified calcium overload cancer therapy. <i>Biomaterials</i> , 2021, 277, 121080.	5.7	53
4	Mulberry ethanol extract attenuates hepatic steatosis and insulin resistance in high-fat diet-fed mice. <i>Nutrition Research</i> , 2016, 36, 710-718.	1.3	44
5	White Pitaya (<i>Hylocereus undatus</i>) Juice Attenuates Insulin Resistance and Hepatic Steatosis in Diet-Induced Obese Mice. <i>PLoS ONE</i> , 2016, 11, e0149670.	1.1	40
6	Dietary anthocyanin-rich extract of açaí protects from diet-induced obesity, liver steatosis, and insulin resistance with modulation of gut microbiota in mice. <i>Nutrition</i> , 2021, 86, 111176.	1.1	37
7	Dietary sweet cherry anthocyanins attenuates diet-induced hepatic steatosis by improving hepatic lipid metabolism in mice. <i>Nutrition</i> , 2016, 32, 827-833.	1.1	31
8	Pomegranate fruit pulp polyphenols reduce diet-induced obesity with modulation of gut microbiota in mice. <i>Journal of the Science of Food and Agriculture</i> , 2022, 102, 1968-1977.	1.7	27
9	Black rice anthocyanins alleviate hyperlipidemia, liver steatosis and insulin resistance by regulating lipid metabolism and gut microbiota in obese mice. <i>Food and Function</i> , 2021, 12, 10160-10170.	2.1	23
10	Black Current Anthocyanins Improve Lipid Metabolism and Modulate Gut Microbiota in High-Fat Diet-Induced Obese Mice. <i>Molecular Nutrition and Food Research</i> , 2021, 65, e2001090.	1.5	23
11	A wheat germ-derived peptide YDWPGGRN facilitates skin wound-healing processes. <i>Biochemical and Biophysical Research Communications</i> , 2020, 524, 943-950.	1.0	20
12	<i>Vaccinium bracteatum</i> Thunb. fruit extract reduces high-fat diet-induced obesity with modulation of the gut microbiota in obese mice. <i>Journal of Food Biochemistry</i> , 2021, 45, e13808.	1.2	14
13	A review on processing methods and functions of wheat germ-derived bioactive peptides. <i>Critical Reviews in Food Science and Nutrition</i> , 2023, 63, 5577-5593.	5.4	13
14	Soybean-derived gma-miR159a alleviates colon tumorigenesis by suppressing TCF7/MYC in mice. <i>Journal of Nutritional Biochemistry</i> , 2021, 92, 108627.	1.9	12
15	Pomegranate peel anthocyanins prevent diet-induced obesity and insulin resistance in association with modulation of the gut microbiota in mice. <i>European Journal of Nutrition</i> , 2022, 61, 1837-1847.	1.8	12
16	The guideline for western blotting assay. <i>Food Frontiers</i> , 2022, 3, 347-349.	3.7	6
17	Coffee consumption is not associated with the risk of gastric cancer: An updated systematic review and meta-analysis of prospective cohort studies. <i>Nutrition Research</i> , 2022, 102, 35-44.	1.3	4
18	Ferric ammonium citrate (FAC)-induced inhibition of osteoblast proliferation/differentiation and its reversal by soybean-derived peptides (SDP). <i>Food and Chemical Toxicology</i> , 2021, 156, 112527.	1.8	1