

Yue-Ming Zhang

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/7712330/publications.pdf>

Version: 2024-02-01

10
papers

265
citations

1039880

9
h-index

1372474

10
g-index

10
all docs

10
docs citations

10
times ranked

327
citing authors

#	ARTICLE	IF	CITATIONS
1	Calycosin ameliorates doxorubicin-induced cardiotoxicity by suppressing oxidative stress and inflammation via the sirtuin 1 "NOD-like receptor protein 3 pathway. <i>Phytotherapy Research</i> , 2020, 34, 649-659.	2.8	59
2	Dysregulation of BSEP and MRP2 May Play an Important Role in Isoniazid-Induced Liver Injury & via the SIRT1/FXR Pathway in Rats and HepG2 Cells. <i>Biological and Pharmaceutical Bulletin</i> , 2018, 41, 1211-1218.	0.6	47
3	XingNaojing injection ameliorates cerebral ischaemia/reperfusion injury via SIRT1-mediated inflammatory response inhibition. <i>Pharmaceutical Biology</i> , 2020, 58, 16-24.	1.3	31
4	Xingnaojing Injection Protects against Cerebral Ischemia Reperfusion Injury via PI3K/Akt-Mediated eNOS Phosphorylation. <i>Evidence-based Complementary and Alternative Medicine</i> , 2018, 2018, 1-13.	0.5	25
5	Quercetin protected against isoniazide-induced HepG2 cell apoptosis by activating the SIRT1/ERK pathway. <i>Journal of Biochemical and Molecular Toxicology</i> , 2019, 33, e22369.	1.4	25
6	Quercetin attenuates NLRP3 inflammasome activation and apoptosis to protect INH-induced liver injury via regulating SIRT1 pathway. <i>International Immunopharmacology</i> , 2020, 85, 106634.	1.7	25
7	XingNaojing injections protect against cerebral ischemia/reperfusion injury and alleviate blood-brain barrier disruption in rats, through an underlying mechanism of NLRP3 inflammasomes suppression. <i>Chinese Journal of Natural Medicines</i> , 2019, 17, 498-505.	0.7	22
8	Ginsenoside Rg3 attenuates cisplatin-induced kidney injury through inhibition of apoptosis and autophagy-inhibited NLRP3. <i>Journal of Biochemical and Molecular Toxicology</i> , 2021, 35, e22896.	1.4	13
9	Salvianolic Acid B Attenuates Apoptosis of HUVEC Cells Treated with High Glucose or High Fat via Sirt1 Activation. <i>Evidence-based Complementary and Alternative Medicine</i> , 2019, 2019, 1-11.	0.5	12
10	Salvianolic acid inhibits the effects of high glucose on vascular endothelial dysfunction by modulating the Sirt1-eNOS pathway. <i>Journal of Biochemical and Molecular Toxicology</i> , 2019, 33, e22245.	1.4	6