Mark Myatt

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/7716692/publications.pdf Version: 2024-02-01



#	Article	IF	CITATIONS
1	Recommendations for surveillance of transmitted HIV drug resistance in countries scaling up antiretroviral treatment. Antiviral Therapy, 2008, 13, 25-36.	0.6	222
2	A Review of Methods to Detect Cases of Severely Malnourished Children in the Community for Their Admission into Community-Based Therapeutic Care Programs. Food and Nutrition Bulletin, 2006, 27, S7-S23.	0.5	215
3	Children who are both wasted and stunted are also underweight and have a high risk of death: a descriptive epidemiology of multiple anthropometric deficits using data from 51 countries. Archives of Public Health, 2018, 76, 28.	1.0	119
4	Boys are more likely to be undernourished than girls: a systematic review and meta-analysis of sex differences in undernutrition. BMJ Global Health, 2020, 5, e004030.	2.0	118
5	Cost-effectiveness of the community-based management of severe acute malnutrition by community health workers in southern Bangladesh. Health Policy and Planning, 2013, 28, 386-399.	1.0	87
6	A novel sequential sampling technique for the surveillance of transmitted HIV drug resistance by cross-sectional survey for use in low resource settings. Antiviral Therapy, 2008, 13, 37-48.	0.6	76
7	Concurrent wasting and stunting among underâ€five children in Niakhar, Senegal. Maternal and Child Nutrition, 2019, 15, e12736.	1.4	62
8	Low mid-upper arm circumference identifies children with a high risk of death who should be the priority target for treatment. BMC Nutrition, 2016, 2, .	0.6	56
9	The effect of body shape on weight-for-height and mid-upper arm circumference based case definitions of acute malnutrition in Ethiopian children. Annals of Human Biology, 2009, 36, 5-20.	0.4	54
10	Mothers Understand And Can do it (MUAC): a comparison of mothers and community health workers determining mid-upper arm circumference in 103 children aged from 6Âmonths to 5Âyears. Archives of Public Health, 2015, 73, 26.	1.0	52
11	Field trial of applicability of lot quality assurance sampling survey method for rapid assessment of prevalence of active trachoma. Bulletin of the World Health Organization, 2003, 81, 877-85.	1.5	41
12	A novel sequential sampling technique for the surveillance of transmitted HIV drug resistance by cross-sectional survey for use in low resource settings. Antiviral Therapy, 2008, 13 Suppl 2, 37-48.	0.6	35
13	Improving screening for malnourished children at high risk of death: a study of children aged 6–59 months in rural Senegal. Public Health Nutrition, 2019, 22, 862-871.	1.1	31
14	Safety and practicability of using mid-upper arm circumference as a discharge criterion in community based management of severe acute malnutrition in children aged 6 to 59 months programmes. Archives of Public Health, 2016, 74, 24.	1.0	30
15	Understanding Sex Differences in Childhood Undernutrition: A Narrative Review. Nutrients, 2022, 14, 948.	1.7	28
16	Factors associated with concurrent wasting and stunting among children 6–59 months in Karamoja, Uganda. Maternal and Child Nutrition, 2021, 17, e13074.	1.4	25
17	Using lot quality-assurance sampling and area sampling to identify priority areas for trachoma control: Viet Nam. Bulletin of the World Health Organization, 2005, 83, 756-63.	1.5	25
18	Improving Estimates of Numbers of Children With Severe Acute Malnutrition Using Cohort and Survey Data. American Journal of Epidemiology, 2016, 184, 861-869.	1.6	24

Mark Myatt

#	Article	IF	CITATIONS
19	A field trial of a survey method for estimating the coverage of selective feeding programmes. Bulletin of the World Health Organization, 2005, 83, 20-6.	1.5	23
20	Relationship between mid upper arm circumference and weight changes in children aged 6–59 months. Archives of Public Health, 2015, 73, 54.	1.0	18
21	The cost of preventing undernutrition: cost, cost-efficiency and cost-effectiveness of three cash-based interventions on nutrition outcomes in Dadu, Pakistan. Health Policy and Planning, 2018, 33, 743-754.	1.0	18
22	Improving estimates of the burden of severe acute malnutrition and predictions of caseload for programs treating severe acute malnutrition: experiences from Nigeria. Archives of Public Health, 2017, 75, 66.	1.0	17
23	Individual and household risk factors of severe acute malnutrition among under-five children in Mao, Chad: a matched case-control study. Archives of Public Health, 2018, 76, 35.	1.0	17
24	Concurrently wasted and stunted children 6â€59 months in Karamoja, Uganda: prevalence and case detection. Maternal and Child Nutrition, 2020, 16, e13000.	1.4	16
25	Concurrently wasted and stunted 6-59 months children admitted to the outpatient therapeutic feeding programme in Karamoja, Uganda: Prevalence, characteristics, treatment outcomes and response. PLoS ONE, 2020, 15, e0230480.	1.1	15
26	Mid-upper-arm circumference based case-detection, admission, and discharging of under five children in a large-scale community-based management of acute malnutrition program in Nigeria. Archives of Public Health, 2018, 76, 19.	1.0	14
27	Maternal and child nutrition. Lancet, The, 2013, 382, 1549.	6.3	12
28	Prevention of child wasting: Results of a Child Health & Nutrition Research Initiative (CHNRI) prioritisation exercise. PLoS ONE, 2020, 15, e0228151.	1.1	12
29	How Can Nutrition Research Better Reflect the Relationship Between Wasting and Stunting in Children? Learnings from the Wasting and Stunting Project. Journal of Nutrition, 2022, 152, 2645-2651.	1.3	8
30	Response to Malnutrition Treatment in Low Weight-for-Age Children: Secondary Analyses of Children 6–59 Months in the ComPAS Cluster Randomized Controlled Trial. Nutrients, 2021, 13, 1054.	1.7	7
31	Using cross-sectional surveys to estimate the number of severely malnourished children needing to be enrolled in specific treatment programmes. Public Health Nutrition, 2017, 20, 1362-1366.	1.1	6
32	Changing sex differences in undernutrition of African children: findings from Demographic and Health Surveys. Journal of Biosocial Science, 2022, 54, 847-857.	0.5	6
33	Concurrent wasting and stunting among children 6–59 months: an analysis using district-level survey data in Mozambique. BMC Nutrition, 2022, 8, 15.	0.6	4
34	Disability-adjusted life years for severe acute malnutrition: implications of alternative model specifications. Public Health Nutrition, 2019, 22, 2729-2737.	1.1	3
35	Effects on child growth of a reduction in the general food distribution ration and provision of small-quantity lipid-based nutrient supplements in refugee camps in eastern Chad. BMJ Nutrition, Prevention and Health, 2021, 4, 235-242.	1.9	3