

# Mario Gennaro Mazza

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/771990/publications.pdf>

Version: 2024-02-01

31  
papers

2,739  
citations

471371

17  
h-index

414303

32  
g-index

33  
all docs

33  
docs citations

33  
times ranked

3239  
citing authors

| #  | ARTICLE  | IF  | CITATIONS |
|----|--|-----|-----------|
| 1  | Cognitive remediation therapy for post-acute persistent cognitive deficits in COVID-19 survivors: A proof-of-concept study. <i>Neuropsychological Rehabilitation</i> , 2023, 33, 1207-1224.  | 1.0 | 8         |
| 2  | Long-term consequences of COVID-19 on cognitive functioning up to 6 months after discharge: role of depression and impact on quality of life. <i>European Archives of Psychiatry and Clinical Neuroscience</i> , 2022, 272, 773-782.   | 1.8 | 67        |
| 3  | Rapid response to selective serotonin reuptake inhibitors in post-COVID depression. <i>European Neuropsychopharmacology</i> , 2022, 54, 1-6.   | 0.3 | 37        |
| 4  | One-year mental health outcomes in a cohort of COVID-19 survivors. <i>Journal of Psychiatric Research</i> , 2022, 145, 118-124.  | 1.5 | 57        |
| 5  | Machine learning approaches for prediction of bipolar disorder based on biological, clinical and neuropsychological markers: A systematic review and meta-analysis. <i>Neuroscience and Biobehavioral Reviews</i> , 2022, 135, 104552. | 2.9 | 7         |
| 6  | Comment on: "Fluvoxamine for the Early Treatment of SARS-CoV-2 Infection: A Review of Current Evidence". <i>Drugs</i> , 2022, 82, 349.   | 4.9 | 1         |
| 7  | A Nomogram-Based Model to Predict Respiratory Dysfunction at 6 Months in Non-Critical COVID-19 Survivors. <i>Frontiers in Medicine</i> , 2022, 9, 781410.  | 1.2 | 3         |
| 8  | Antidepressant psychopharmacology: is inflammation a future target?. <i>International Clinical Psychopharmacology</i> , 2022, 37, 79-81.   | 0.9 | 17        |
| 9  | Mood-congruent negative thinking styles and cognitive vulnerability in depressed COVID-19 survivors: A comparison with major depressive disorder. <i>Journal of Affective Disorders</i> , 2022, 308, 554-561.                          | 2.0 | 6         |
| 10 | Post-COVID-19 Depressive Symptoms: Epidemiology, Pathophysiology, and Pharmacological Treatment. <i>CNS Drugs</i> , 2022, 36, 681-702.   | 2.7 | 83        |
| 11 | Lower levels of glutathione in the anterior cingulate cortex associate with depressive symptoms and white matter hyperintensities in COVID-19 survivors. <i>European Neuropsychopharmacology</i> , 2022, 61, 71-77.                    | 0.3 | 13        |
| 12 | A peripheral inflammatory signature discriminates bipolar from unipolar depression: A machine learning approach. <i>Progress in Neuro-Psychopharmacology and Biological Psychiatry</i> , 2021, 105, 110136.                            | 2.5 | 49        |
| 13 | Higher baseline interleukin-1 $\beta$ and TNF- $\alpha$ hamper antidepressant response in major depressive disorder. <i>European Neuropsychopharmacology</i> , 2021, 42, 35-44.  | 0.3 | 25        |
| 14 | Persistent psychopathology and neurocognitive impairment in COVID-19 survivors: Effect of inflammatory biomarkers at three-month follow-up. <i>Brain, Behavior, and Immunity</i> , 2021, 94, 138-147.                                  | 2.0 | 299       |
| 15 | Mental disorders and risk of COVID-19-related mortality, hospitalisation, and intensive care unit admission: a systematic review and meta-analysis. <i>Lancet Psychiatry</i> , 2021, 8, 797-812.                                       | 3.7 | 202       |
| 16 | Higher Interleukin 13 differentiates patients with a positive history of suicide attempts in major depressive disorder. <i>Journal of Affective Disorders Reports</i> , 2021, 6, 100254.   | 0.9 | 5         |
| 17 | Brain correlates of depression, post-traumatic distress, and inflammatory biomarkers in COVID-19 survivors: A multimodal magnetic resonance imaging study. <i>Brain, Behavior, &amp; Immunity - Health</i> , 2021, 18, 100387.         | 1.3 | 57        |
| 18 | Antipsychotics and COVID-19: the debate goes on " Authors' reply. <i>Lancet Psychiatry</i> , 2021, 8, 1030-1031.   | 3.7 | 2         |

| #  | ARTICLE  | IF  | CITATIONS |
|----|--|-----|-----------|
| 19 | Neutrophil-lymphocyte ratio, monocyte-lymphocyte ratio and platelet-lymphocyte ratio in non-affective psychosis: A meta-analysis and systematic review. <i>World Journal of Biological Psychiatry</i> , 2020, 21, 326-338.           | 1.3 | 95        |
| 20 | Prevalence of co-occurring psychiatric disorders in adults and adolescents with intellectual disability: A systematic review and meta-analysis. <i>Journal of Applied Research in Intellectual Disabilities</i> , 2020, 33, 126-138. | 1.3 | 86        |
| 21 | Anxiety and depression in COVID-19 survivors: Role of inflammatory and clinical predictors. <i>Brain, Behavior, and Immunity</i> , 2020, 89, 594-600.  | 2.0 | 1,118     |
| 22 | Proinflammatory Cytokines Predict Brain Metabolite Concentrations in the Anterior Cingulate Cortex of Patients With Bipolar Disorder. <i>Frontiers in Psychiatry</i> , 2020, 11, 590095.   | 1.3 | 16        |
| 23 | Monocyte count in schizophrenia and related disorders: a systematic review and meta-analysis. <i>Acta Neuropsychiatrica</i> , 2020, 32, 229-236.   | 1.0 | 28        |
| 24 | Residual clinical damage after COVID-19: A retrospective and prospective observational cohort study. <i>PLoS ONE</i> , 2020, 15, e0239570.   | 1.1 | 129       |
| 25 | Neutrophil-lymphocyte, monocyte-lymphocyte and platelet-lymphocyte ratio in schizoaffective disorder compared to schizophrenia. <i>General Hospital Psychiatry</i> , 2019, 61, 86-87.  | 1.2 | 6         |
| 26 | Cross-sectional study of neutrophil-lymphocyte, platelet-lymphocyte and monocyte-lymphocyte ratios in mood disorders. <i>General Hospital Psychiatry</i> , 2019, 58, 7-12.   | 1.2 | 46        |
| 27 | Neutrophil/lymphocyte ratio and platelet/lymphocyte ratio in mood disorders: A meta-analysis. <i>Progress in Neuro-Psychopharmacology and Biological Psychiatry</i> , 2018, 84, 229-236.   | 2.5 | 183       |
| 28 | White matter alterations associate with onset symptom dimension in obsessive-compulsive disorder. <i>Psychiatry and Clinical Neurosciences</i> , 2018, 72, 13-27.  | 1.0 | 10        |
| 29 | Vortioxetine overdose in a suicidal attempt. <i>Medicine (United States)</i> , 2018, 97, e10788.   | 0.4 | 8         |
| 30 | Uric acid levels in subjects with bipolar disorder: A comparative meta-analysis. <i>Journal of Psychiatric Research</i> , 2016, 81, 133-139.   | 1.5 | 68        |
| 31 | A single nucleotide polymorphism in SLC1A1 gene is associated with age of onset of obsessive-compulsive disorder. <i>European Psychiatry</i> , 2014, 29, 301-303.  | 0.1 | 7         |