

JosÃ© Luis Venero

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/7725688/publications.pdf>

Version: 2024-02-01

115
papers

6,801
citations

71004

43
h-index

75989

78
g-index

118
all docs

118
docs citations

118
times ranked

9379
citing authors

#	ARTICLE	IF	CITATIONS
1	Gal3 Plays a Deleterious Role in a Mouse Model of Endotoxemia. International Journal of Molecular Sciences, 2022, 23, 1170.	1.8	3
2	Arginine deprivation alters microglial polarity and synergizes with radiation to eradicate non-arginine-auxotrophic glioblastoma tumors. Journal of Clinical Investigation, 2022, 132, .	3.9	28
3	Inflammatory Animal Models of Parkinsonâ€™s Disease. Journal of Parkinson's Disease, 2022, 12, S165-S182.	1.5	9
4	Selective deletion of Caspase-3 gene in the dopaminergic system exhibits autistic-like behaviour. Progress in Neuro-Psychopharmacology and Biological Psychiatry, 2021, 104, 110030.	2.5	9
5	Galectin-3 Deletion Reduces LPS and Acute Colitis-Induced Pro-Inflammatory Microglial Activation in the Ventral Mesencephalon. Frontiers in Pharmacology, 2021, 12, 706439.	1.6	6
6	Hydroxytyrosol Decreases LPS- and Î±-Synuclein-Induced Microglial Activation In Vitro. Antioxidants, 2020, 9, 36.	2.2	28
7	Microglia: Agents of the CNS Pro-Inflammatory Response. Cells, 2020, 9, 1717.	1.8	174
8	Hyperinflammation and Fibrosis in Severe COVID-19 Patients: Galectin-3, a Target Molecule to Consider. Frontiers in Immunology, 2020, 11, 2069.	2.2	66
9	The Ubiquitin Proteasome System in Neuromuscular Disorders: Moving Beyond Movement. International Journal of Molecular Sciences, 2020, 21, 6429.	1.8	17
10	Microglial subtypes: diversity within the microglial community. EMBO Journal, 2019, 38, e101997.	3.5	345
11	Reformulating Pro-Oxidant Microglia in Neurodegeneration. Journal of Clinical Medicine, 2019, 8, 1719.	1.0	47
12	TET2 Regulates the Neuroinflammatory Response in Microglia. Cell Reports, 2019, 29, 697-713.e8.	2.9	74
13	Galectin-3, a novel endogenous TREM2 ligand, detrimentally regulates inflammatory response in Alzheimerâ€™s disease. Acta Neuropathologica, 2019, 138, 251-273.	3.9	187
14	Magnetofection as a new tool to study microglia biology. Neural Regeneration Research, 2019, 14, 767.	1.6	1
15	HERC1 Ubiquitin Ligase Is Required for Normal Axonal Myelination in the Peripheral Nervous System. Molecular Neurobiology, 2018, 55, 8856-8868.	1.9	14
16	Peripheral Inflammation Enhances Microglia Response and Nigral Dopaminergic Cell Death in an in vivo MPTP Model of Parkinsonâ€™s Disease. Frontiers in Cellular Neuroscience, 2018, 12, 398.	1.8	67
17	Divergent Effects of Metformin on an Inflammatory Model of Parkinsonâ€™s Disease. Frontiers in Cellular Neuroscience, 2018, 12, 440.	1.8	43
18	Caspases orchestrate microglia instrumental functions. Progress in Neurobiology, 2018, 171, 50-71.	2.8	27

#	ARTICLE	IF	CITATIONS
19	RIPK1 is a critical modulator of both tonic and TLR-responsive inflammatory and cell death pathways in human macrophage differentiation. <i>Cell Death and Disease</i> , 2018, 9, 973.	2.7	33
20	Galectin-3 released in response to traumatic brain injury acts as an alarmin orchestrating brain immune response and promoting neurodegeneration. <i>Scientific Reports</i> , 2017, 7, 41689.	1.6	120
21	Caspase-8 inhibition represses initial human monocyte activation in septic shock model. <i>Oncotarget</i> , 2016, 7, 37456-37470.	0.8	16
22	Chronic stress alters the expression levels of longevity-related genes in the rat hippocampus. <i>Neurochemistry International</i> , 2016, 97, 181-192.	1.9	26
23	Glioma-induced inhibition of caspase-3 in microglia promotes a tumor-supportive phenotype. <i>Nature Immunology</i> , 2016, 17, 1282-1290.	7.0	76
24	Spatio-temporal activation of caspase-8 in myeloid cells upon ischemic stroke. <i>Acta Neuropathologica Communications</i> , 2016, 4, 92.	2.4	17
25	Metformin, besides exhibiting strong in vivo anti-inflammatory properties, increases mptp-induced damage to the nigrostriatal dopaminergic system. <i>Toxicology and Applied Pharmacology</i> , 2016, 298, 19-30.	1.3	72
26	PGC-1 β controls mitochondrial biogenesis and dynamics in lead-induced neurotoxicity. <i>Aging</i> , 2015, 7, 629-643.	1.4	87
27	Relevance of chronic stress and the two faces of microglia in Parkinson's disease. <i>Frontiers in Cellular Neuroscience</i> , 2015, 9, 312.	1.8	36
28	Microglia-Secreted Galectin-3 Acts as a Toll-like Receptor 4 Ligand and Contributes to Microglial Activation. <i>Cell Reports</i> , 2015, 10, 1626-1638.	2.9	268
29	Synergistic Deleterious Effect of Chronic Stress and Sodium Azide in the Mouse Hippocampus. <i>Chemical Research in Toxicology</i> , 2015, 28, 651-661.	1.7	4
30	Neuromelanin activates proinflammatory microglia through a caspase-8-dependent mechanism. <i>Journal of Neuroinflammation</i> , 2015, 12, 5.	3.1	38
31	Evaluation of a method for murine monocyte isolation by bone marrow depletion. <i>Analytical Biochemistry</i> , 2015, 480, 42-48.	1.1	5
32	Deletion of caspase-8 in mouse myeloid cells blocks microglia pro-inflammatory activation and confers protection in MPTP neurodegeneration model. <i>Aging</i> , 2015, 7, 673-689.	1.4	28
33	Collateral Damage: Contribution of Peripheral Inflammation to Neurodegenerative Diseases. <i>Current Topics in Medicinal Chemistry</i> , 2015, 15, 2193-2210.	1.0	37
34	Regulation of caspase-3 processing by cIAP2 controls the switch between pro-inflammatory activation and cell death in microglia. <i>Cell Death and Disease</i> , 2014, 5, e1565-e1565.	2.7	65
35	Chronic stress as a risk factor for Alzheimer's disease. <i>Reviews in the Neurosciences</i> , 2014, 25, 785-804.	1.4	132
36	The role of Galectin-3 in α -synuclein-induced microglial activation. <i>Acta Neuropathologica Communications</i> , 2014, 2, 156.	2.4	63

#	ARTICLE	IF	CITATIONS
37	Role of dopamine in the recruitment of immune cells to the nigro-striatal dopaminergic structures. <i>NeuroToxicology</i> , 2014, 41, 89-101.	1.4	25
38	Chronic stress enhances microglia activation and exacerbates death of nigral dopaminergic neurons under conditions of inflammation. <i>Journal of Neuroinflammation</i> , 2014, 11, 34.	3.1	157
39	Caspases Playing in the Field of Neuroinflammation: Old and New Players. <i>Developmental Neuroscience</i> , 2013, 35, 88-101.	1.0	35
40	A Brief Overview of Multitalented Microglia. <i>Methods in Molecular Biology</i> , 2013, 1041, 3-8.	0.4	24
41	Peripheral inflammation increases the deleterious effect of CNS inflammation on the nigrostriatal dopaminergic system. <i>NeuroToxicology</i> , 2012, 33, 347-360.	1.4	87
42	The executioners sing a new song: killer caspases activate microglia. <i>Cell Death and Differentiation</i> , 2011, 18, 1679-1691.	5.0	47
43	Stress is critical for LPS-induced activation of microglia and damage in the rat hippocampus. <i>Neurobiology of Aging</i> , 2011, 32, 85-102.	1.5	128
44	Peripheral Inflammation Increases the Damage in Animal Models of Nigrostriatal Dopaminergic Neurodegeneration: Possible Implication in Parkinson's Disease Incidence. <i>Parkinson's Disease</i> , 2011, 2011, 1-10.	0.6	35
45	Caspase signalling controls microglia activation and neurotoxicity. <i>Nature</i> , 2011, 472, 319-324.	13.7	491
46	Apoptosis-inducing factor mediates dopaminergic cell death in response to LPS-induced inflammatory stimulus. <i>Neurobiology of Disease</i> , 2011, 41, 177-188.	2.1	64
47	Nanostructures for Drug Delivery to the Brain. <i>Current Medicinal Chemistry</i> , 2011, 18, 5303-5321.	1.2	43
48	Ulcerative colitis exacerbates lipopolysaccharide-induced damage to the nigral dopaminergic system: potential risk factor in Parkinson's disease. <i>Journal of Neurochemistry</i> , 2010, 114, 1687-1700.	2.1	169
49	Use of haptoglobin and transthyretin as potential biomarkers for the preclinical diagnosis of Parkinson's disease. <i>Neurochemistry International</i> , 2010, 57, 227-234.	1.9	37
50	Striatal ablation of GABAergic neurons prevents the in vivo neuroprotective effect of DCG-IV against the MPP+-induced neurotoxicity on dopaminergic nerve terminals. <i>Neurochemistry International</i> , 2010, 57, 979-984.	1.9	1
51	Regional-specific regulation of BDNF and <i>trkB</i> correlates with nigral dopaminergic cell sprouting following unilateral nigrostriatal axotomy. <i>Journal of Neuroscience Research</i> , 2008, 86, 2016-2027.	1.3	3
52	The intrastriatal injection of thrombin in rat induced a retrograde apoptotic degeneration of nigral dopaminergic neurons through synaptic elimination. <i>Journal of Neurochemistry</i> , 2008, 105, 750-762.	2.1	12
53	Intracerebral VEGF injection highly upregulates AQP4 mRNA and protein in the perivascular space and glia limitans externa. <i>Neurochemistry International</i> , 2008, 52, 897-903.	1.9	41
54	Proteomic identification of biomarkers in the cerebrospinal fluid in a rat model of nigrostriatal dopaminergic degeneration. <i>Journal of Neuroscience Research</i> , 2007, 85, 3607-3618.	1.3	25

#	ARTICLE	IF	CITATIONS
55	Blood-brain barrier disruption induces <i>in vivo</i> degeneration of nigral dopaminergic neurons. <i>Journal of Neurochemistry</i> , 2007, 101, 1567-1582.	2.1	125
56	Mitochondrial toxins and neurodegenerative diseases. <i>Frontiers in Bioscience - Landmark</i> , 2007, 12, 986.	3.0	53
57	Stress Increases Vulnerability to Inflammation in the Rat Prefrontal Cortex. <i>Journal of Neuroscience</i> , 2006, 26, 5709-5719.	1.7	187
58	Blood-brain barrier disruption highly induces aquaporin-4 mRNA and protein in perivascular and parenchymal astrocytes: Protective effect by estradiol treatment in ovariectomized animals. <i>Journal of Neuroscience Research</i> , 2005, 80, 235-246.	1.3	101
59	Kainate-induced zinc translocation from presynaptic terminals causes neuronal and astroglial cell death and mRNA loss of BDNF receptors in the hippocampal formation and amygdala. <i>Journal of Neuroscience Research</i> , 2005, 82, 184-195.	1.3	16
60	Inflammatory process as a determinant factor for the degeneration of substantia nigra dopaminergic neurons. <i>Journal of Neural Transmission</i> , 2005, 112, 111-119.	1.4	95
61	Divergent regulatory mechanisms governing BDNF mRNA expression in cerebral cortex and substantia nigra in response to striatal target ablation. <i>Experimental Neurology</i> , 2005, 192, 142-155.	2.0	9
62	In vivo expression of aquaporin-4 by reactive microglia. <i>Journal of Neurochemistry</i> , 2004, 91, 891-899.	2.1	62
63	Minocycline reduces the lipopolysaccharide-induced inflammatory reaction, peroxynitrite-mediated nitration of proteins, disruption of the blood-brain barrier, and damage in the nigral dopaminergic system. <i>Neurobiology of Disease</i> , 2004, 16, 190-201.	2.1	187
64	Importance of Aquaporins in the Physiopathology of Brain Edema. <i>Current Pharmaceutical Design</i> , 2004, 10, 2153-2161.	0.9	35
65	Evidence for dopamine-derived hydroxyl radical formation in the nigrostriatal system in response to axotomy. <i>Free Radical Biology and Medicine</i> , 2003, 34, 111-123.	1.3	8
66	Thrombin induces <i>in vivo</i> degeneration of nigral dopaminergic neurones along with the activation of microglia. <i>Journal of Neurochemistry</i> , 2003, 84, 1201-1214.	2.1	75
67	Expression of BDNF mRNA in substantia nigra is dependent on target integrity and independent of neuronal activation. <i>Journal of Neurochemistry</i> , 2003, 87, 709-721.	2.1	14
68	Semichronic Inhibition of Glutathione Reductase Promotes Oxidative Damage to Proteins and Induces both Transcription and Translation of Tyrosine Hydroxylase in the Nigrostriatal System. <i>Free Radical Research</i> , 2003, 37, 1003-1012.	1.5	6
69	Differential regulation of glutamic acid decarboxylase mRNA and tyrosine hydroxylase mRNA expression in the aged manganese-treated rats. <i>Molecular Brain Research</i> , 2002, 103, 116-129.	2.5	42
70	DCG-IV but not other group-II metabotropic receptor agonists induces microglial BDNF mRNA expression in the rat striatum. Correlation with neuronal injury. <i>Neuroscience</i> , 2002, 113, 857-869.	1.1	42
71	Melatonin induces tyrosine hydroxylase mRNA expression in the ventral mesencephalon but not in the hypothalamus. <i>Journal of Pineal Research</i> , 2002, 32, 6-14.	3.4	29
72	Histamine Infusion Induces a Selective Dopaminergic Neuronal Death Along with an Inflammatory Reaction in Rat Substantia Nigra. <i>Journal of Neurochemistry</i> , 2002, 75, 540-552.	2.1	68

#	ARTICLE	IF	CITATIONS
73	Potential Role of Endogenous Brain-Derived Neurotrophic Factor in Long-Term Neuronal Reorganization of the Superior Colliculus after Bilateral Visual Deprivation. <i>Neurobiology of Disease</i> , 2001, 8, 866-880.	2.1	34
74	Long-lasting induction of brain-derived neurotrophic factor is restricted to resistant cell populations in an animal model of status epilepticus. <i>Neuroscience</i> , 2001, 103, 955-969.	1.1	8
75	Aquaporins in the central nervous system. <i>Progress in Neurobiology</i> , 2001, 63, 321-336.	2.8	182
76	Group II metabotropic glutamate receptor activation protects striatal dopaminergic nerve terminals against MPP ⁺ -induced neurotoxicity along with brain-derived neurotrophic factor induction. <i>Journal of Neurochemistry</i> , 2001, 76, 351-360.	2.1	81
77	Upregulation of BDNF mRNA and trkB mRNA in the Nigrostriatal System and in the Lesion Site Following Unilateral Transection of the Medial Forebrain Bundle. <i>Experimental Neurology</i> , 2000, 161, 38-48.	2.0	22
78	The Single Intranigral Injection of LPS as a New Model for Studying the Selective Effects of Inflammatory Reactions on Dopaminergic System. <i>Neurobiology of Disease</i> , 2000, 7, 429-447.	2.1	373
79	Decreased messenger RNA expression of key markers of the nigrostriatal dopaminergic system following vitamin E deficiency in the rat. <i>Neuroscience</i> , 2000, 101, 1029-1036.	1.1	6
80	Low selenium diet induces tyrosine hydroxylase enzyme in nigrostriatal system of the rat. <i>Molecular Brain Research</i> , 2000, 84, 7-16.	2.5	16
81	Localization of Aquaporin-3 mRNA and protein along the gastrointestinal tract of Wistar rats. <i>Pflügers Archiv European Journal of Physiology</i> , 1999, 438, 94-100.	1.3	56
82	Serotonin hyperinnervation in the adult rat ventral mesencephalon following unilateral transection of the medial forebrain bundle. Correlation with reactive microglial and astroglial populations. <i>Neuroscience</i> , 1999, 91, 567-577.	1.1	20
83	Detailed localization of aquaporin-4 messenger RNA in the CNS: preferential expression in periventricular organs. <i>Neuroscience</i> , 1999, 94, 239-250.	1.1	121
84	Delayed apoptotic pyramidal cell death in CA4 and CA1 hippocampal subfields after a single intraseptal injection of kainate. <i>Neuroscience</i> , 1999, 94, 1071-1081.	1.1	26
85	Differential Upregulation of Aquaporin-4 mRNA Expression in Reactive Astrocytes after Brain Injury: Potential Role in Brain Edema. <i>Neurobiology of Disease</i> , 1999, 6, 245-258.	2.1	227
86	Regionally specific induction of BDNF and truncated trkB.T1 receptors in the hippocampal formation after intraseptal injection of kainic acid. <i>Brain Research</i> , 1998, 790, 270-277.	1.1	25
87	Developmental expression of 5-HT ₇ receptor mRNA in rat brain visual structures after neonatal enucleation. <i>NeuroReport</i> , 1997, 8, 1531-1535.	0.6	9
88	Deprenyl induces the tyrosine hydroxylase enzyme in the rat dopaminergic nigrostriatal system. <i>Molecular Brain Research</i> , 1997, 46, 31-38.	2.5	25
89	Expression of 5-HT ₇ receptor mRNA in rat brain during postnatal development. <i>Neuroscience Letters</i> , 1997, 227, 53-56.	1.0	47
90	Less induced 1-methyl-4-phenylpyridinium ion neurotoxicity on striatal slices from guinea-pigs fed with a vitamin C-deficient diet. <i>Neuroscience</i> , 1997, 77, 167-174.	1.1	9

#	ARTICLE	IF	CITATIONS
91	Protective effects of neurotrophin-4/5 and transforming growth factor- $\hat{\alpha}$ on striatal neuronal phenotypic degeneration after excitotoxic lesioning with quinolinic acid. <i>Neuroscience</i> , 1997, 78, 73-86.	1.1	48
92	Increased Activity and Expression of Tyrosine Hydroxylase in the Rat Substantia Nigra after Chronic Treatment with Nomifensine. <i>Molecular Pharmacology</i> , 1997, 52, 641-647.	1.0	4
93	Chronic Inhibition of the High-Affinity Dopamine Uptake System Increases Oxidative Damage to Proteins in the Aged Rat Substantia Nigra. <i>Free Radical Biology and Medicine</i> , 1997, 23, 1-7.	1.3	11
94	Deprenyl induces GFAP immunoreactivity in the intact and injured dopaminergic nigrostriatal system but fails to counteract axotomy-induced degenerative changes. <i>Glia</i> , 1997, 21, 204-216.	2.5	13
95	Time Course Changes in the Dopaminergic Nigrostriatal System Following Transection of the Medial Forebrain Bundle: Detection of Oxidatively Modified Proteins in Substantia Nigra. <i>Journal of Neurochemistry</i> , 1997, 68, 2458-2468.	2.1	41
96	MK-801 PARTIALLY PROTECTS AGAINST THE ACUTE MPP+ DEPLETING EFFECT ON DOPAMINE LEVELS IN RAT STRIATAL SLICES. <i>Neurochemistry International</i> , 1996, 29, 411-416.	1.9	8
97	IMT-4/5 protects against adrenalectomy-induced apoptosis of rat hippocampal granule cells. <i>NeuroReport</i> , 1996, 7, 682.	0.6	16
98	Oxidative inactivation of tyrosine hydroxylase in substantia nigra of aged rat. <i>Free Radical Biology and Medicine</i> , 1996, 20, 53-61.	1.3	66
99	Intrastratial quinolinic acid injections protect against 6-hydroxydopamine-induced lesions of the dopaminergic nigrostriatal system. <i>Brain Research</i> , 1995, 672, 153-158.	1.1	19
100	Retrograde transport of nerve growth factor from hippocampus and amygdala to trk A messenger RNA expressing neurons in paraventricular and reuniens nuclei of the thalamus. <i>Neuroscience</i> , 1995, 64, 855-860.	1.1	9
101	Intrastratial infusion of nerve growth factor after quinolinic acid prevents reduction of cellular expression of choline acetyltransferase messenger RNA and trkA messenger RNA, but not glutamate decarboxylase messenger RNA. <i>Neuroscience</i> , 1994, 61, 257-268.	1.1	52
102	Expression of neurotrophin and trk receptor genes in adult rats with fimbria transections: Effect of intraventricular nerve growth factor and brain-derived neurotrophic factor administration. <i>Neuroscience</i> , 1994, 59, 797-815.	1.1	90
103	6-Hydroxydopamine lesions reduce BDNF mRNA levels in adult rat brain substantia nigra. <i>NeuroReport</i> , 1994, 5, 429-432.	0.6	34
104	Effect of ageing on monoamine turnover in the prefrontal cortex of rats. <i>Mechanisms of Ageing and Development</i> , 1993, 72, 105-118.	2.2	13
105	Age-related changes on monoamine turnover in hippocampus of rats. <i>Brain Research</i> , 1993, 631, 89-96.	1.1	52
106	TrkA NGF receptor expression by non-cholinergic thalamic neurons. <i>NeuroReport</i> , 1993, 4, 959-962.	0.6	17
107	Changes in neurotransmitter levels associated with the deficiency of some essential amino acids in the diet. <i>British Journal of Nutrition</i> , 1992, 68, 409-420.	1.2	23
108	In vivo protection of striatum from MPP+ neurotoxicity by N-methyl-d-aspartate antagonists. <i>Brain Research</i> , 1992, 586, 203-207.	1.1	41

#	ARTICLE	IF	CITATIONS
109	Changes in the Turnover of Monoamines in Prefrontal Cortex of Rats Fed on Vitamin E-Deficient Diet. <i>Journal of Neurochemistry</i> , 1992, 58, 1889-1895.	2.1	6
110	Age effects on monoamine turnover of the rat substantia nigra. <i>Brain Research</i> , 1991, 557, 109-114.	1.1	20
111	Turnover of Dopamine and Serotonin and Their Metabolites in the Striatum of Aged Rats. <i>Journal of Neurochemistry</i> , 1991, 56, 1940-1948.	2.1	36
112	Effects of neonatal bilateral eye enucleation on postnatal development of the monoamines in posterior thalamus of the rat. <i>Journal of Neural Transmission</i> , 1991, 85, 231-242.	1.4	3
113	Determination of levels of biogenic amines and their metabolites and both forms of monoamine oxidase in prefrontal cortex of aged rats. <i>Mechanisms of Ageing and Development</i> , 1990, 56, 253-263.	2.2	9
114	Changes in monoamines and their metabolite levels in substantia nigra of aged rats. <i>Mechanisms of Ageing and Development</i> , 1989, 49, 227-233.	2.2	14
115	Determination of monoamines and both forms of monoamine oxidase in the rat's substantia nigra during postnatal development. <i>Life Sciences</i> , 1989, 45, 1277-1283.	2.0	17