
## Anna Kakehashi

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/7731062/publications.pdf Version: 2024-02-01



ΔΝΝΑ ΚΑΚΕΠΑςΗΙ

#	Article	IF	CITATIONS
1	FOXP3 and CXCR4-positive regulatory T cells in the tumor stroma as indicators of tumor immunity in the conjunctival squamous cell carcinoma microenvironment. PLoS ONE, 2022, 17, e0263895.	1.1	1
2	The carbonic anhydrase inhibitor acetazolamide inhibits urinary bladder cancers via suppression of βâ€catenin signaling. Cancer Science, 2022, 113, 2642-2653.	1.7	3
3	Cache Domain Containing 1 Is a Novel Marker of Non-Alcoholic Steatohepatitis-Associated Hepatocarcinogenesis. Cancers, 2021, 13, 1216.	1.7	5
4	Expression of thrombospondin-1 in conjunctival squamous cell carcinoma is correlated to the Ki67 index and associated with progression-free survival. Graefe's Archive for Clinical and Experimental Ophthalmology, 2021, 259, 3127-3136.	1.0	1
5	Canopy Homolog 2 as a Novel Molecular Target in Hepatocarcinogenesis. Cancers, 2021, 13, 3613.	1.7	4
6	Accumulation of 8-hydroxydeoxyguanosine, L-arginine and Glucose Metabolites by Liver Tumor Cells Are the Important Characteristic Features of Metabolic Syndrome and Non-Alcoholic Steatohepatitis-Associated Hepatocarcinogenesis. International Journal of Molecular Sciences, 2020, 21, 7746.	1.8	17
7	Expression, intracellular localization, and mutation of EGFR in conjunctival squamous cell carcinoma and the association with prognosis and treatment. PLoS ONE, 2020, 15, e0238120.	1.1	6
8	Title is missing!. , 2020, 15, e0238120.		0
9	Title is missing!. , 2020, 15, e0238120.		0
10	Title is missing!. , 2020, 15, e0238120.		0
11	Title is missing!. , 2020, 15, e0238120.		Ο
12	Title is missing!. , 2020, 15, e0238120.		0
13	Title is missing!. , 2020, 15, e0238120.		Ο
14	mTOR Activation in Liver Tumors Is Associated with Metabolic Syndrome and Non-Alcoholic Steatohepatitis in Both Mouse Models and Humans. Cancers, 2018, 10, 465.	1.7	12
15	Steroid sulfatase promotes invasion through epithelial‑mesenchymal transition and predicts the progression of bladder cancer. Experimental and Therapeutic Medicine, 2018, 16, 4463-4470.	0.8	1
16	Chronic dietary toxicity and carcinogenicity studies of dammar resin in F344 rats. Archives of Toxicology, 2018, 92, 3565-3583.	1.9	1
17	Carbonic anhydrase 2 is a novel invasionâ€associated factor in urinary bladder cancers. Cancer Science, 2017, 108, 331-337.	1.7	12
18	Enhanced Susceptibility of Ogg1 Mutant Mice to Multiorgan Carcinogenesis. International Journal of Molecular Sciences, 2017, 18, 1801.	1.8	16

Anna Kakehashi

#	Article	IF	CITATIONS
19	Proteome Characteristics of Non-Alcoholic Steatohepatitis Liver Tissue and Associated Hepatocellular Carcinomas. International Journal of Molecular Sciences, 2017, 18, 434.	1.8	20
20	Progression of Hepatic Adenoma to Carcinoma in <i>Ogg1</i> Mutant Mice Induced by Phenobarbital. Oxidative Medicine and Cellular Longevity, 2017, 2017, 1-16.	1.9	9
21	Diphenylarsinic acid exerts promotion effects on hepatobiliary carcinogenesis in a rat medium-term multiorgan carcinogenicity bioassay. Journal of Toxicologic Pathology, 2017, 30, 39-45.	0.3	3
22	Chemopreventive Action by Ethanol-extracted Brazilian Green Propolis on Post-initiation Phase of Inflammation-associated Rat Colon Tumorigenesis. In Vivo, 2017, 31, 187-198.	0.6	24
23	Pueraria mirifica Exerts Estrogenic Effects in the Mammary Gland and Uterus and Promotes Mammary Carcinogenesis in Donryu Rats. Toxins, 2016, 8, 275.	1.5	9
24	Examination of in vivo mutagenicity of sodium arsenite and dimethylarsinic acid in gpt delta rats. Journal of Environmental Sciences, 2016, 49, 125-130.	3.2	6
25	Ethanol-Extracted Brazilian Propolis Exerts Protective Effects on Tumorigenesis in Wistar Hannover Rats. PLoS ONE, 2016, 11, e0158654.	1.1	17
26	Induction of cell proliferation in the rat liver by the short-term administration of ethyl <i>tertiary</i> -butyl ether. Journal of Toxicologic Pathology, 2015, 28, 27-32.	0.3	8
27	Integrative analyses of miRNA and proteomics identify potential biological pathways associated with onset of pulmonary fibrosis in the bleomycin rat model. Toxicology and Applied Pharmacology, 2015, 286, 188-197.	1.3	14
28	Ethanol-extracted propolis enhances BBN-initiated urinary bladder carcinogenesis via non-mutagenic mechanisms in rats. Food and Chemical Toxicology, 2015, 83, 193-200.	1.8	7
29	Isoleucine, Leucine and Their Role in Experimental Models of Bladder Carcinogenesis. , 2015, , 253-260.		1
30	Valerian Inhibits Rat Hepatocarcinogenesis by Activating GABA(A) Receptor-Mediated Signaling. PLoS ONE, 2014, 9, e113610.	1.1	11
31	Comparative Proteomics Analysis of Gastric Cancer Stem Cells. PLoS ONE, 2014, 9, e110736.	1.1	39
32	l-Leucine and l-isoleucine enhance growth of BBN-induced urothelial tumors in the rat bladder by modulating expression of amino acid transporters and tumorigenesis-associated genes. Food and Chemical Toxicology, 2013, 59, 137-144.	1.8	19
33	Mode of action of ethyl tertiary-butyl ether hepatotumorigenicity in the rat: Evidence for a role of oxidative stress via activation of CAR, PXR and PPAR signaling pathways. Toxicology and Applied Pharmacology, 2013, 273, 390-400.	1.3	29
34	Oxidative Stress in the Carcinogenicity of Chemical Carcinogens. Cancers, 2013, 5, 1332-1354.	1.7	39
35	Evaluation of the Modifying Effect of Inhalation of Mainstream Cigarette Smoke on Mouse Bladder Carcinogenesis. Journal of Toxicologic Pathology, 2013, 26, 447-451.	0.3	2
36	2-Amino-3-Methylimidazo[4,5-f]Quinoline (IQ) Promotes Mouse Hepatocarcinogenesis by Activating Transforming Growth Factor-Â and Wnt/Â-Catenin Signaling Pathways. Toxicological Sciences, 2012, 125, 392-400.	1.4	4

#	Article	IF	CITATIONS
37	Lack of Hepatocarcinogenicity of Combinations of Low Doses of 2-amino-3, 8-dimethylimidazo[4,5- <i>f</i> ]quinoxaline and Diethylnitrosamine in Rats: Indication for the Existence of a Threshold for Genotoxic Carcinogens. Journal of Toxicologic Pathology, 2012, 25, 209-214.	0.3	3
38	Hormonally Active Doses of Isoflavone Aglycones Promote Mammary and Endometrial Carcinogenesis and Alter the Molecular Tumor Environment in Donryu Rats. Toxicological Sciences, 2012, 126, 39-51.	1.4	23
39	Long-term treatment with l-isoleucine or l-leucine in AIN-93G diet has promoting effects on rat bladder carcinogenesis. Food and Chemical Toxicology, 2012, 50, 3934-3940.	1.8	14
40	Dammar resin, a non-mutagen, inducts oxidative stress and metabolic enzymes in the liver of gpt delta transgenic mouse which is different from a mutagen, 2-amino-3-methylimidazo[4,5-f]quinoline. Mutation Research - Genetic Toxicology and Environmental Mutagenesis, 2012, 748, 29-35.	0.9	7
41	<scp>DDX</scp> 39 acts as a suppressor of invasion for bladder cancer. Cancer Science, 2012, 103, 1363-1369.	1.7	27
42	Methionine Sulfoxide Stimulates Hepatocarcinogenesis in Non-alcoholic Steatohepatitis (NASH) Mouse: Possible Role of Free Radical-mediated DNA Methylation. Genes and Environment, 2012, 34, 123-128.	0.9	3
43	Threshold for Genotoxic Carcinogens: The Central Concern in Carcinogenic Risk Assessment. Genes and Environment, 2012, 34, 153-156.	0.9	3
44	Non-genotoxic mode of action and possible threshold for hepatocarcinogenicity of Kojic acid in F344 rats. Food and Chemical Toxicology, 2011, 49, 471-476.	1.8	18
45	Mitochondrial Prohibitins and Septin 9 Are Implicated in the Onset of Rat Hepatocarcinogenesis. Toxicological Sciences, 2011, 119, 61-72.	1.4	44
46	Targeted Proteomics of Isolated Glomeruli from the Kidneys of Diabetic Rats: Sorbin and SH3 Domain Containing 2 Is a Novel Protein Associated with Diabetic Nephropathy. Experimental Diabetes Research, 2011, 2011, 1-11.	3.8	28
47	Enhanced Urinary Bladder, Liver and Colon Carcinogenesis in Zucker Diabetic Fatty Rats in a Multiorgan Carcinogenesis Bioassay: Evidence for Mechanisms Involving Activation of PI3K Signaling and Impairment of P53 on Urinary Bladder Carcinogenesis. Journal of Toxicologic Pathology, 2011, 24, 25-36.	0.3	12
48	Cytokeratin 8/18 as a new marker of mouse liver preneoplastic lesions. Toxicology and Applied Pharmacology, 2010, 242, 47-55.	1.3	29
49	Evaluation of the Subchronic Toxicity of Dietary Administered Equisetum arvense in F344 Rats. Journal of Toxicologic Pathology, 2010, 23, 245-251.	0.3	11
50	Rat Monoclonal Antibody Specific for Septin 9. Hybridoma, 2010, 29, 169-171.	0.5	3
51	Ethanol Does Not Promote MelQx-initiated Rat Colon Carcinogenesis Based on Evidence from Analysis of a Colon Cancer Surrogate Marker. Journal of Toxicologic Pathology, 2009, 22, 65-70.	0.3	1
52	Potassium Bromate Enhances N-Ethyl-N-Hydroxyethylnitrosamine–Induced Kidney Carcinogenesis Only at High Doses in Wistar Rats: Indication of the Existence of an Enhancement Threshold. Toxicologic Pathology, 2009, 37, 983-991.	0.9	14
53	Cytokeratin 8/18 overexpression and complex formation as an indicator of GST-P positive foci transformation into hepatocellular carcinomas. Toxicology and Applied Pharmacology, 2009, 238, 71-79.	1.3	32
54	Characteristic Upregulation of Glucose-Regulated Protein 78 in an Early Lesion Negative for Hitherto Established Cytochemical Markers in Rat Hepatocarcinogenesis. Journal of Toxicologic Pathology, 2009, 22, 281-288.	0.3	2

#	Article	lF	CITATIONS
55	Existence of a Threshold for the Genotoxic Carcinogens: Evidence from Mechanism-based Carcinogenicity Studies. Genes and Environment, 2009, 31, 33-36.	0.9	5
56	Chemopreventive effects of a serratane-type triterpenoid, 3α-methoxyserrat-14-en-21β-ol (PJ-1), against rat lung carcinogenesis. Food and Chemical Toxicology, 2008, 46, 1882-1888.	1.8	12
57	Are there thresholds for carcinogens carcinogenicity?. Mycotoxins, 2008, 58, 119-128.	0.2	0