

Daniela Patinha

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/7731487/publications.pdf>

Version: 2024-02-01

15
papers

349
citations

1040056

9
h-index

1199594

12
g-index

15
all docs

15
docs citations

15
times ranked

601
citing authors

#	ARTICLE	IF	CITATIONS
1	Acute SGLT inhibition normalizes O ₂ tension in the renal cortex but causes hypoxia in the renal medulla in anaesthetized control and diabetic rats. <i>American Journal of Physiology - Renal Physiology</i> , 2015, 309, F227-F234.	2.7	180
2	Role of H ₂ O ₂ in hypertension, renin-angiotensin system activation and renal medullary dysfunction caused by angiotensin II. <i>British Journal of Pharmacology</i> , 2012, 166, 2386-2401.	5.4	37
3	Microinjection of angiotensin II in the caudal ventrolateral medulla induces hyperalgesia. <i>Neuroscience</i> , 2009, 158, 1301-1310.	2.3	33
4	Angiotensin II contributes to glomerular hyperfiltration in diabetic rats independently of adenosine type I receptors. <i>American Journal of Physiology - Renal Physiology</i> , 2013, 304, F614-F622.	2.7	21
5	Activation of adenosine receptors improves renal antioxidant status in diabetic Wistar but not SHR rats. <i>Upsala Journal of Medical Sciences</i> , 2014, 119, 10-18.	0.9	16
6	Cooperative Oxygen Sensing by the Kidney and Carotid Body in Blood Pressure Control. <i>Frontiers in Physiology</i> , 2017, 8, 752.	2.8	14
7	Diabetes-induced increase of renal medullary hydrogen peroxide and urinary angiotensinogen is similar in normotensive and hypertensive rats. <i>Life Sciences</i> , 2014, 108, 71-79.	4.3	10
8	NADPH oxidase 1 is a novel pharmacological target for the development of an antiplatelet drug without bleeding side effects. <i>FASEB Journal</i> , 2020, 34, 13959-13977.	0.5	10
9	Angiotensin II-induced hypertension in rats is only transiently accompanied by lower renal oxygenation. <i>Scientific Reports</i> , 2018, 8, 16342.	3.3	9
10	Adenosine A2A and A3 Receptors as Targets for the Treatment of Hypertensive-Diabetic Nephropathy. <i>Biomedicines</i> , 2020, 8, 529.	3.2	9
11	Diabetes downregulates renal adenosine A2A receptors in an experimental model of hypertension. <i>PLoS ONE</i> , 2019, 14, e0217552.	2.5	7
12	Determinants of renal oxygen metabolism during low Na + diet: effect of angiotensin II AT 1 and aldosterone receptor blockade. <i>Journal of Physiology</i> , 2020, 598, 5573-5587.	2.9	3
13	Intrarenal Blockade of Angiotensin II AT 1 Receptor Abolishes Renal Cortical Hypoxia in Salt Restricted Animals. <i>FASEB Journal</i> , 2015, 29, 963.2.	0.5	0
14	Acute IP Furosemide Increases Medullary PO ₂ in The Diabetic Rat Kidney. <i>FASEB Journal</i> , 2015, 29, 963.9.	0.5	0
15	Acute IP Phlorizin Normalizes Cortical PO ₂ But Causes Medullary Hypoxia in The Diabetic Rat Kidney. <i>FASEB Journal</i> , 2015, 29, 959.2.	0.5	0