

# Katsuomi Iwakura

## List of Publications by Year in Descending Order

**Source:** <https://exaly.com/author-pdf/773855/katsuomi-iwakura-publications-by-year.pdf>

**Version:** 2024-04-20

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

35  
papers

1,526  
citations

15  
h-index

36  
g-index

36  
ext. papers

1,677  
ext. citations

3.5  
avg, IF

3.52  
L-index

#	Paper	IF	Citations
35	Wire Cutting Method Using Rotational Atherectomy for Stretched Spring Wire During Coronary Intervention.. <i>JACC: Case Reports</i> , <b>2021</b> , 3, 1842-1848	1.2	1
34	Usefulness of Post-Procedural Plasma Brain Natriuretic Peptide Levels to Predict Recurrence After Catheter Ablation of Atrial Fibrillation in Patients With Left Ventricular Systolic Dysfunction. <i>American Journal of Cardiology</i> , <b>2021</b> , 144, 67-76	3	0
33	A practical guide to the lung ultrasound for the assessment of congestive heart failure. <i>Journal of Echocardiography</i> , <b>2021</b> , 19, 195-204	1.6	0
32	Going back home digitally: gift for patients with end-stage heart failure. <i>European Heart Journal Digital Health</i> , <b>2021</b> , 2, 351-353	2.3	
31	Decrease in red cell distribution width as a useful predictor of success after catheter ablation for atrial fibrillation: a retrospective multi-center study. <i>Heart and Vessels</i> , <b>2021</b> , 1	2.1	
30	Burden and Long Firing of Premature Atrial Contraction Early After Catheter Ablation Predict Late Recurrence of Atrial Fibrillation. <i>Circulation Journal</i> , <b>2020</b> , 84, 894-901	2.9	6
29	Impact of baseline left atrial function on long-term outcome after catheter ablation for paroxysmal atrial fibrillation. <i>Journal of Cardiology</i> , <b>2020</b> , 75, 352-359	3	5
28	Early recurrence during the blanking period and left atrial reverse remodeling after catheter ablation for non-paroxysmal atrial fibrillation. <i>IJC Heart and Vasculature</i> , <b>2020</b> , 30, 100588	2.4	2
27	Heart failure in patients with type 2 diabetes mellitus: assessment with echocardiography and effects of antihyperglycemic treatments. <i>Journal of Echocardiography</i> , <b>2019</b> , 17, 177-186	1.6	9
26	Outcomes After Atrial Fibrillation Ablation in Patients With Premature Atrial Contractions Originating From Non-Pulmonary Veins. <i>JACC: Clinical Electrophysiology</i> , <b>2019</b> , 5, 1319-1327	4.6	8
25	Usefulness of Failed Electrical Cardioversion for Early Recurrence After Catheter Ablation for Atrial Fibrillation as a Predictor of Future Recurrence. <i>American Journal of Cardiology</i> , <b>2019</b> , 123, 794-800	3	1
24	Left Atrial Reverse Remodeling After Catheter Ablation of Nonparoxysmal Atrial Fibrillation in Patients With Heart Failure With Reduced Ejection Fraction. <i>American Journal of Cardiology</i> , <b>2018</b> , 122, 89-96	3	13
23	Automated Ablation Annotation Algorithm Reduces Re-conduction of Isolated Pulmonary Vein and Improves Outcome After Catheter Ablation for Atrial Fibrillation. <i>Circulation Journal</i> , <b>2017</b> , 81, 1596-1602 <sup>9</sup>	2.9	23
22	Stress hyperglycemia and microvascular obstruction after acute myocardial infarction. <i>Journal of Cardiology</i> , <b>2015</b> , 65, 270-1	3	1
21	Changes in left ventricular relaxation after azelnidipine treatment in hypertensive patients with diabetes: subanalysis of a prospective single-arm multicentre study. <i>BMJ Open</i> , <b>2014</b> , 4, e006136	3	1
20	Cause of very late recurrence of atrial fibrillation or flutter after catheter ablation for atrial fibrillation. <i>American Journal of Cardiology</i> , <b>2013</b> , 111, 552-6	3	38
19	Modulation of individual susceptibility to the no-reflow phenomenon after acute myocardial infarction. <i>Current Pharmaceutical Design</i> , <b>2013</b> , 19, 4519-28	3.3	6

18	Oral treatment with nicorandil at discharge is associated with reduced mortality after acute myocardial infarction. <i>Journal of Cardiology</i> , <b>2012</b> , 59, 14-21	3	22
17	Trigger-based mechanism of the persistence of atrial fibrillation and its impact on the efficacy of catheter ablation. <i>Circulation: Arrhythmia and Electrophysiology</i> , <b>2012</b> , 5, 295-301	6.4	65
16	Effect of elevated left ventricular diastolic filling pressure on the frequency of left atrial appendage thrombus in patients with nonvalvular atrial fibrillation. <i>American Journal of Cardiology</i> , <b>2011</b> , 107, 417-22	3	35
15	Visualization of myocardial hemorrhage with real-time three-dimensional myocardial contrast echocardiography in patients with acute myocardial infarction. <i>Journal of Echocardiography</i> , <b>2011</b> , 9, 154-5	1.6	1
14	Automated assessment of myocardial viability after acute myocardial infarction by global longitudinal peak strain on low-dose dobutamine stress echocardiography. <i>Circulation Journal</i> , <b>2010</b> , 74, 2158-65	2.9	8
13	Impact of azelnidipine treatment on left ventricular diastolic performance in patients with hypertension and mild diastolic dysfunction: multi-center study with echocardiography. <i>Hypertension Research</i> , <b>2009</b> , 32, 895-900	4.7	16
12	Nicorandil treatment in patients with acute myocardial infarction: a meta-analysis. <i>Circulation Journal</i> , <b>2009</b> , 73, 925-31	2.9	33
11	Normal values of echocardiographic parameters in relation to age in a healthy Japanese population: the JAMP study. <i>Circulation Journal</i> , <b>2008</b> , 72, 1859-66	2.9	146
10	Comparison of two- versus three-dimensional myocardial contrast echocardiography for assessing subendocardial perfusion abnormality after percutaneous coronary intervention in patients with acute myocardial infarction. <i>American Journal of Cardiology</i> , <b>2007</b> , 100, 1502-10	3	7
9	Usefulness of high-resolution real-time three-dimensional echocardiography to visualize the left ventricular endocardial surface in myocardial infarction. <i>American Journal of Cardiology</i> , <b>2006</b> , 97, 1578-81	3	6
8	Comparison of orifice area by transthoracic three-dimensional Doppler echocardiography versus proximal isovelocity surface area (PISA) method for assessment of mitral regurgitation. <i>American Journal of Cardiology</i> , <b>2006</b> , 97, 1630-7	3	82
7	Chronic pre-treatment of statins is associated with the reduction of the no-reflow phenomenon in the patients with reperfused acute myocardial infarction. <i>European Heart Journal</i> , <b>2006</b> , 27, 534-9	9.5	106
6	Intravenous nicorandil in conjunction with coronary reperfusion therapy is associated with better clinical and functional outcomes in patients with acute myocardial infarction. <i>Circulation Journal</i> , <b>2003</b> , 67, 295-300	2.9	58
5	Association between hyperglycemia and the no-reflow phenomenon in patients with acute myocardial infarction. <i>Journal of the American College of Cardiology</i> , <b>2003</b> , 41, 1-7	15.1	276
4	Predictive factors for development of the no-reflow phenomenon in patients with reperfused anterior wall acute myocardial infarction. <i>Journal of the American College of Cardiology</i> , <b>2001</b> , 38, 472-7	15.1	152
3	Use of echocardiography for predicting myocardial viability in patients with reperfused anterior wall myocardial infarction. <i>American Journal of Cardiology</i> , <b>2000</b> , 85, 744-8	3	15
2	Intravenous nicorandil can preserve microvascular integrity and myocardial viability in patients with reperfused anterior wall myocardial infarction. <i>Journal of the American College of Cardiology</i> , <b>1999</b> , 33, 654-60	15.1	313
1	Alpha 1-adrenoceptor activation increases ecto-5'-nucleotidase activity and adenosine release in rat cardiomyocytes by activating protein kinase C. <i>Circulation</i> , <b>1995</b> , 91, 2226-34	16.7	71

