

Chantal Michel

List of Publications by Citations

Source: <https://exaly.com/author-pdf/7738930/chantal-michel-publications-by-citations.pdf>

Version: 2024-04-26

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

61
papers

1,438
citations

21
h-index

37
g-index

99
ext. papers

1,828
ext. citations

4.8
avg. IF

5.05
L-index

#	Paper	IF	Citations
61	EPA guidance on the early detection of clinical high risk states of psychoses. <i>European Psychiatry</i> , 2015 , 30, 405-16	6	235
60	Age matters in the prevalence and clinical significance of ultra-high-risk for psychosis symptoms and criteria in the general population: Findings from the BEAR and BEARS-kid studies. <i>World Psychiatry</i> , 2015 , 14, 189-97	14.4	80
59	O10.5. CLINICAL HIGH RISK (CHR) CRITERIA IN CHILDREN AND ADOLESCENTS NOT SUSPECTED TO DEVELOP PSYCHOSIS: NO TRANSDIAGNOSTIC SEVERITY MARKER OF MENTAL HEALTH PROBLEMS. <i>Schizophrenia Bulletin</i> , 2020 , 46, S26-S26	1.3	78
58	O11.8. PREVALENCE AND PREDICTORS OF INTERVIEW-ASSESSED CLINICAL HIGH-RISK SYMPTOMS IN THE GENERAL POPULATION. <i>Schizophrenia Bulletin</i> , 2018 , 44, S109-S109	1.3	78
57	F26. THE NATURE OF CLINICAL HIGH-RISK SYMPTOMS: NEW INSIGHTS GAINED FROM AGE EFFECTS. <i>Schizophrenia Bulletin</i> , 2018 , 44, S228-S229	1.3	78
56	Prevalence and clinical significance of DSM-5-attenuated psychosis syndrome in adolescents and young adults in the general population: the Bern Epidemiological At-Risk (BEAR) study. <i>Schizophrenia Bulletin</i> , 2014 , 40, 1499-508	1.3	70
55	A stratified model for psychosis prediction in clinical practice. <i>Schizophrenia Bulletin</i> , 2014 , 40, 1533-42	1.3	53
54	⌈A rose is a rose is a rose⌋ but at-risk criteria differ. <i>Psychopathology</i> , 2013 , 46, 75-87	3.4	51
53	Psychosis-predictive value of self-reported schizotypy in a clinical high-risk sample. <i>Journal of Abnormal Psychology</i> , 2016 , 125, 923-932	7	46
52	Course of clinical high-risk states for psychosis beyond conversion. <i>European Archives of Psychiatry and Clinical Neuroscience</i> , 2018 , 268, 39-48	5.1	43
51	Duration of unspecific prodromal and clinical high risk states, and early help-seeking in first-admission psychosis patients. <i>Social Psychiatry and Psychiatric Epidemiology</i> , 2015 , 50, 1831-41	4.5	41
50	What percentage of people in the general population satisfies the current clinical at-risk criteria of psychosis?. <i>Schizophrenia Research</i> , 2011 , 125, 99-100	3.6	41
49	Revisiting the Basic Symptom Concept: Toward Translating Risk Symptoms for Psychosis into Neurobiological Targets. <i>Frontiers in Psychiatry</i> , 2016 , 7, 9	5	40
48	Personality disorders and accentuations in at-risk persons with and without conversion to first-episode psychosis. <i>Microbial Biotechnology</i> , 2012 , 6, 389-98	3.3	34
47	Differences in coping, self-efficacy, and external control beliefs between patients at-risk for psychosis and patients with first-episode psychosis. <i>Psychiatry Research</i> , 2014 , 219, 95-102	9.9	33
46	Prevalence and clinical relevance of interview-assessed psychosis-risk symptoms in the young adult community. <i>Psychological Medicine</i> , 2018 , 48, 1167-1178	6.9	32
45	Multimodal Machine Learning Workflows for Prediction of Psychosis in Patients With Clinical High-Risk Syndromes and Recent-Onset Depression. <i>JAMA Psychiatry</i> , 2021 , 78, 195-209	14.5	31

44	Abnormal involuntary movements are linked to psychosis-risk in children and adolescents: Results of a population-based study. <i>Schizophrenia Research</i> , 2016 , 174, 58-64	3.6	28
43	Coping as a mediator between locus of control, competence beliefs, and mental health: A systematic review and structural equation modelling meta-analysis. <i>Behaviour Research and Therapy</i> , 2019 , 121, 103442	5.2	26
42	Predictors for DSM-5 nonsuicidal self-injury in female adolescent inpatients: The role of childhood maltreatment, alexithymia, and dissociation. <i>Psychiatry Research</i> , 2016 , 239, 346-52	9.9	23
41	Age effect on prevalence of ultra-high risk for psychosis symptoms: replication in a clinical sample of an early detection of psychosis service. <i>European Child and Adolescent Psychiatry</i> , 2017 , 26, 1401-1405	5.5	21
40	The interrelationship between schizotypy, clinical high risk for psychosis and related symptoms: Cognitive disturbances matter. <i>Schizophrenia Research</i> , 2019 , 210, 188-196	3.6	19
39	Association of Structural Magnetic Resonance Imaging Measures With Psychosis Onset in Individuals at Clinical High Risk for Developing Psychosis: An ENIGMA Working Group Mega-analysis. <i>JAMA Psychiatry</i> , 2021 , 78, 753-766	14.5	17
38	The interplay of etiological knowledge and mental illness stigma on healthcare utilisation in the community: A structural equation model. <i>European Psychiatry</i> , 2018 , 51, 48-56	6	15
37	Reliability of telephone assessments of at-risk criteria of psychosis: a comparison to face-to-face interviews. <i>Schizophrenia Research</i> , 2014 , 153, 251-3	3.6	15
36	What Is the Incidence of Implant Malpositioning and Revision Surgery After Orbital Repair?. <i>Journal of Oral and Maxillofacial Surgery</i> , 2018 , 76, 146-153	1.8	13
35	Striatal cerebral blood flow, executive functioning, and fronto-striatal functional connectivity in clinical high risk for psychosis. <i>Schizophrenia Research</i> , 2018 , 201, 231-236	3.6	13
34	The trait-state distinction between schizotypy and clinical high risk: results from a one-year follow-up. <i>World Psychiatry</i> , 2019 , 18, 108-109	14.4	13
33	Clinical evaluation of the iterative metal artefact reduction algorithm for post-operative CT examination after maxillofacial surgery. <i>Dentomaxillofacial Radiology</i> , 2017 , 46, 20160355	3.9	11
32	Functioning mediates help-seeking for mental problems in the general population. <i>European Psychiatry</i> , 2018 , 54, 1-9	6	11
31	Is a black eye a useful sign of facial fractures in patients with minor head injuries? A retrospective analysis in a level I trauma centre over 10 years. <i>British Journal of Oral and Maxillofacial Surgery</i> , 2014 , 52, 518-22	1.4	11
30	Effects of age and sex on clinical high-risk for psychosis in the community. <i>World Journal of Psychiatry</i> , 2020 , 10, 101-124	3	11
29	What are the limitations of the non-patient-specific implant in titanium reconstruction of the orbit?. <i>British Journal of Oral and Maxillofacial Surgery</i> , 2020 , 58, e80-e85	1.4	11
28	Age effects on basic symptoms in the community: A route to gain new insight into the neurodevelopment of psychosis?. <i>European Archives of Psychiatry and Clinical Neuroscience</i> , 2020 , 270, 311-324	5.1	11
27	Development of stimulus material for research in alcohol use disorders. <i>International Journal of Methods in Psychiatric Research</i> , 2017 , 26,	4.3	10

26	Mediators Linking Childhood Adversities and Trauma to Suicidality in Individuals at Risk for Psychosis. <i>Frontiers in Psychiatry</i> , 2017 , 8, 242	5	10
25	Differentiating schizoaffective and bipolar I disorder in first-episode psychotic mania. <i>Schizophrenia Research</i> , 2012 , 140, 31-6	3.6	10
24	Convergent and concurrent validity of the Frankfurt Complaint Questionnaire as a screener for psychosis risk. <i>Journal of Risk Research</i> , 2017 , 20, 1480-1496	4.2	9
23	Functional and structural correlates of abnormal involuntary movements in psychosis risk and first episode psychosis. <i>Schizophrenia Research</i> , 2019 , 212, 196-203	3.6	9
22	Demographic and clinical characteristics of diagnosed and non-diagnosed psychotic disorders in the community. <i>Microbial Biotechnology</i> , 2018 , 12, 87-90	3.3	5
21	Associations of psychosis-risk symptoms with quality of life and self-rated health in the Community. <i>European Psychiatry</i> , 2019 , 62, 116-123	6	4
20	Twelve-month service use, suicidality and mental health problems of European adolescents after a school-based screening for current suicidality. <i>European Child and Adolescent Psychiatry</i> , 2020 , 1	5.5	4
19	Clinical high-risk of and conversion to psychosis in the community: A 3-year follow-up of a cohort study. <i>Schizophrenia Research</i> , 2021 , 228, 616-618	3.6	3
18	Prediction and Early Detection of First-Episode Psychosis 2011 , 207-267		2
17	Trapped in a Glass Bell Jar: Neural Correlates of Depersonalization and Derealization in Subjects at Clinical High-Risk of Psychosis and Depersonalization-Derealization Disorder. <i>Frontiers in Psychiatry</i> , 2020 , 11, 535652	5	2
16	S211. SCHIZOTYPY IN PATIENTS FROM A CLINICAL HIGH RISK SERVICE: TRAIT OR STATE?. <i>Schizophrenia Bulletin</i> , 2018 , 44, S407-S408	1.3	2
15	Predictors of help-seeking behaviour in people with mental health problems: a 3-year prospective community study. <i>BMC Psychiatry</i> , 2021 , 21, 432	4.2	2
14	Clinical High Risk Symptoms and Criteria in the Community: Prevalence, Clinical Significance and Risk Factors for Their Occurrence. <i>European Psychiatry</i> , 2017 , 41, S226-S226	6	1
13	Basic Symptoms in the Community and their Association with Age. <i>European Psychiatry</i> , 2017 , 41, S84-S85		1
12	Positive psychotic symptoms in childhood and adolescence.. <i>Current Opinion in Psychology</i> , 2021 , 45, 101287	6.2	1
11	Basic symptoms and gray matter volumes of patients at clinical high risk for psychosis. <i>Psychological Medicine</i> , 2021 , 51, 2666-2674	6.9	1
10	The Bern Early Recognition and Intervention Centre for mental crisis (FETZ Bern)-An 8-year evaluation. <i>Microbial Biotechnology</i> , 2021 ,	3.3	1
9	Psychoserisikosyndrome im Kindes- und Jugendalter. <i>Springer Reference Medizin</i> , 2021 , 1-12	0	1

8	Psychometric properties of the Kessler psychological scales in a Swiss young-adult community sample indicate poor suitability for community screening for mental disorders.. <i>Microbial Biotechnology</i> , 2022 ,	3.3	1
7	The Important Role of Stereotypes in the relation between Mental Health Literacy and Stigmatization of Depression and Psychosis in the Community. <i>Community Mental Health Journal</i> , 2021 , 1	2.1	0
6	Clinical high-risk criteria of psychosis in 8-17-year-old community subjects and inpatients not suspected of developing psychosis.. <i>World Journal of Psychiatry</i> , 2022 , 12, 425-449	3	0
5	Childhood Maltreatment and Its Association with Cognitive Ability in Young People Suspected to Be at Clinical High Risk of Psychosis. <i>Psychopathology</i> ,1-12	3.4	0
4	Subjective disturbances in emerging psychosis 2020 , 59-80		
3	The Impact of Age on the Prevalence and Clinical Relevance of Attenuated Psychotic Symptoms in Patients of an Early Detection Service. <i>European Psychiatry</i> , 2017 , 41, S84-S84	6	
2	Prevalence and Burden of at-Risk Criteria of Psychosis and Help-Seeking Behaviour - a Population Survey. <i>European Psychiatry</i> , 2011 , 26, 1128-1128	6	
1	Clinical High Risk for Psychosis Syndromes Among Swiss and German Youth and Young Adults: Early Identification and Intervention 2019 , 115-142		