

# Yoichi Kase

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/7749389/publications.pdf>

Version: 2024-02-01

10  
papers

204  
citations

1307594

7  
h-index

1372567

10  
g-index

12  
all docs

12  
docs citations

12  
times ranked

362  
citing authors

#	ARTICLE	IF	CITATIONS
1	Phase 2 single-arm study on the efficacy and safety of niraparib in Japanese patients with heavily pretreated, homologous recombination-deficient ovarian cancer. <i>Journal of Gynecologic Oncology</i> , 2021, 32, e16.	2.2	8
2	Comparison of the analgesic effects continuous epidural anesthesia and continuous rectus sheath block in patients undergoing gynecological cancer surgery: a non-inferiority randomized control trial. <i>Journal of Anesthesia</i> , 2021, 35, 663-670.	1.7	5
3	Phase 2 single-arm study on the safety of maintenance niraparib in Japanese patients with platinum-sensitive relapsed ovarian cancer. <i>Journal of Gynecologic Oncology</i> , 2021, 32, e21.	2.2	7
4	Reply to the letter titled "Difficulty in evaluating the effect of CEA or CRSB" <i>Journal of Anesthesia</i> , 2021, , 1.	1.7	0
5	Phase 1 study of ixazomib alone or combined with lenalidomide-dexamethasone in Japanese patients with relapsed/refractory multiple myeloma. <i>International Journal of Hematology</i> , 2017, 105, 445-452.	1.6	13
6	Effect of a selective neutrophil elastase inhibitor on mortality and ventilator-free days in patients with increased extravascular lung water: a post hoc analysis of the PiCCO Pulmonary Edema Study. <i>Journal of Intensive Care</i> , 2014, 2, 67.	2.9	78
7	Early-phase changes of extravascular lung water index as a prognostic indicator in acute respiratory distress syndrome patients. <i>Annals of Intensive Care</i> , 2014, 4, 27.	4.6	38
8	Global end-diastolic volume is an important contributor to increased extravascular lung water in patients with acute lung injury and acute respiratory distress syndrome: a multicenter observational study. <i>Journal of Intensive Care</i> , 2014, 2, 25.	2.9	9
9	Limitations of global end-diastolic volume index as a parameter of cardiac preload in the early phase of severe sepsis: a subgroup analysis of a multicenter, prospective observational study. <i>Journal of Intensive Care</i> , 2013, 1, 11.	2.9	19
10	Removal of 2- <sup>15</sup> Arachidonylglycerol by Direct Hemoperfusion Therapy With Polymyxin B Immobilized Fibers Benefits Patients With Septic Shock. <i>Therapeutic Apheresis and Dialysis</i> , 2008, 12, 374-380.	0.9	26