

# Karin Palmblad

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/7754223/publications.pdf>

Version: 2024-02-01

10  
papers

161  
citations

1306789

7  
h-index

1473754

9  
g-index

12  
all docs

12  
docs citations

12  
times ranked

300  
citing authors

#	ARTICLE	IF	CITATIONS
1	Tapering Canakinumab Monotherapy in Patients With Systemic Juvenile Idiopathic Arthritis in Clinical Remission: Results From a Phase IIIb/IV Open-Label, Randomized Study. <i>Arthritis and Rheumatology</i> , 2021, 73, 336-346.	2.9	23
2	Therapeutic administration of etoposide coincides with reduced systemic HMGB1 levels in macrophage activation syndrome. <i>Molecular Medicine</i> , 2021, 27, 48.	1.9	7
3	Immunoprofiling of active and inactive systemic juvenile idiopathic arthritis reveals distinct biomarkers: a single-center study. <i>Pediatric Rheumatology</i> , 2021, 19, 173.	0.9	8
4	Expression of concern to: High systematic levels of the cytokine-inducing HMGB1 isoform secreted in severe macrophage activation syndrome. <i>Molecular Medicine</i> , 2020, 26, 17.	1.9	0
5	Juvenile idiopathic arthritis and risk of cancer before and after the introduction of biological therapies. <i>RMD Open</i> , 2019, 5, e001055.	1.8	9
6	Empowering Young People Living With Juvenile Idiopathic Arthritis to Better Communicate With Families and Care Teams: Content Analysis of Semistructured Interviews. <i>JMIR MHealth and UHealth</i> , 2019, 7, e10401.	1.8	20
7	High Systemic Levels of the Cytokine-Inducing HMGB1 Isoform Secreted in Severe Macrophage Activation Syndrome. <i>Molecular Medicine</i> , 2014, 20, 538-547.	1.9	45
8	HMGB1 mediates muscle fatigue via TLR4 - a possible mechanism for muscle fatigue in patients with inflammatory myopathies. <i>Annals of the Rheumatic Diseases</i> , 2012, 71, A42.2-A43.	0.5	0
9	Morphological characterization of intra-articular HMGB1 expression during the course of collagen-induced arthritis. <i>Arthritis Research and Therapy</i> , 2007, 9, R35.	1.6	36
10	Identification of Rat IL-1 $\beta$ , IL-2, IFN- $\gamma$ and TNF- $\alpha$ in Activated Splenocytes by Intracellular Immunostaining. <i>Biotechnic and Histochemistry</i> , 2000, 75, 101-109.	0.7	10