

Annemieke Kavelaars

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/77544/publications.pdf>

Version: 2024-02-01

77
papers

4,963
citations

71102

41
h-index

98798

67
g-index

80
all docs

80
docs citations

80
times ranked

5801
citing authors

#	ARTICLE	IF	CITATIONS
1	An HDAC6 inhibitor reverses chemotherapy-induced mechanical hypersensitivity via an IL-10 and macrophage dependent pathway. <i>Brain, Behavior, and Immunity</i> , 2022, 100, 287-296.	4.1	11
2	HDAC6 inhibition reverses long-term doxorubicin-induced cognitive dysfunction by restoring microglia homeostasis and synaptic integrity. <i>Theranostics</i> , 2022, 12, 603-619.	10.0	12
3	Targeting the A3 adenosine receptor to prevent and reverse chemotherapy-induced neurotoxicities in mice. <i>Acta Neuropathologica Communications</i> , 2022, 10, 11.	5.2	22
4	Targeting the Meningeal Compartment to Resolve Chemobrain and Neuropathy via Nasal Delivery of Functionalized Mitochondria. <i>Advanced Healthcare Materials</i> , 2022, 11, e2102153.	7.6	8
5	CD8+ T cell-derived IL-13 increases macrophage IL-10 to resolve neuropathic pain. <i>JCI Insight</i> , 2022, 7, .	5.0	31
6	Nasal administration of mesenchymal stem cells reverses chemotherapy-induced peripheral neuropathy in mice. <i>Brain, Behavior, and Immunity</i> , 2021, 93, 43-54.	4.1	23
7	Nasal administration of mitochondria reverses chemotherapy-induced cognitive deficits. <i>Theranostics</i> , 2021, 11, 3109-3130.	10.0	57
8	T Cells as Guardians of Pain Resolution. <i>Trends in Molecular Medicine</i> , 2021, 27, 302-313.	6.7	21
9	Inhibition of dual leucine zipper kinase prevents chemotherapy-induced peripheral neuropathy and cognitive impairments. <i>Pain</i> , 2021, 162, 2599-2612.	4.2	10
10	TTI-101: A competitive inhibitor of STAT3 that spares oxidative phosphorylation and reverses mechanical allodynia in mouse models of neuropathic pain. <i>Biochemical Pharmacology</i> , 2021, 192, 114688.	4.4	16
11	Immune regulation of pain: Friend and foe. <i>Science Translational Medicine</i> , 2021, 13, eabj7152.	12.4	24
12	Bexarotene normalizes chemotherapy-induced myelin decompaction and reverses cognitive and sensorimotor deficits in mice. <i>Acta Neuropathologica Communications</i> , 2020, 8, 193.	5.2	17
13	Interleukin-10 resolves pain hypersensitivity induced by cisplatin by reversing sensory neuron hyperexcitability. <i>Pain</i> , 2020, 161, 2344-2352.	4.2	55
14	Astrocytes rescue neuronal health after cisplatin treatment through mitochondrial transfer. <i>Acta Neuropathologica Communications</i> , 2020, 8, 36.	5.2	64
15	The fibroblast-derived protein PI16 controls neuropathic pain. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2020, 117, 5463-5471.	7.1	39
16	CD3+ T cells are critical for the resolution of comorbid inflammatory pain and depression-like behavior. <i>Neurobiology of Pain (Cambridge, Mass)</i> , 2020, 7, 100043.	2.5	24
17	GRK2 levels in myeloid cells modulate adipose-liver crosstalk in high fat diet-induced obesity. <i>Cellular and Molecular Life Sciences</i> , 2020, 77, 4957-4976.	5.4	5
18	Motivational changes that develop in a mouse model of inflammation-induced depression are independent of indoleamine 2,3 dioxygenase. <i>Neuropsychopharmacology</i> , 2019, 44, 364-371.	5.4	27

#	ARTICLE	IF	CITATIONS
19	Alleviation of paclitaxel-induced mechanical hypersensitivity and hyperalgesic priming with AMPK activators in male and female mice. <i>Neurobiology of Pain (Cambridge, Mass)</i> , 2019, 6, 100037.	2.5	30
20	T Cells as an Emerging Target for Chronic Pain Therapy. <i>Frontiers in Molecular Neuroscience</i> , 2019, 12, 216.	2.9	87
21	Chemotherapy accelerates age-related development of tauopathy and results in loss of synaptic integrity and cognitive impairment. <i>Brain, Behavior, and Immunity</i> , 2019, 79, 319-325.	4.1	31
22	Cisplatin educates CD8+ T cells to prevent and resolve chemotherapy-induced peripheral neuropathy in mice. <i>Pain</i> , 2019, 160, 1459-1468.	4.2	57
23	Cell-specific role of histone deacetylase 6 in chemotherapy-induced mechanical allodynia and loss of intraepidermal nerve fibers. <i>Pain</i> , 2019, 160, 2877-2890.	4.2	37
24	Beyond symptomatic relief for chemotherapy-induced peripheral neuropathy: Targeting the source. <i>Cancer</i> , 2018, 124, 2289-2298.	4.1	115
25	Orally active Epac inhibitor reverses mechanical allodynia and loss of intraepidermal nerve fibers in a mouse model of chemotherapy-induced peripheral neuropathy. <i>Pain</i> , 2018, 159, 884-893.	4.2	38
26	Cisplatin treatment induces attention deficits and impairs synaptic integrity in the prefrontal cortex in mice. <i>Scientific Reports</i> , 2018, 8, 17400.	3.3	28
27	Nasal administration of mesenchymal stem cells restores cisplatin-induced cognitive impairment and brain damage in mice. <i>Oncotarget</i> , 2018, 9, 35581-35597.	1.8	55
28	Mitochondrial transfer from mesenchymal stem cells to neural stem cells protects against the neurotoxic effects of cisplatin. <i>Acta Neuropathologica Communications</i> , 2018, 6, 139.	5.2	93
29	Pharmacological inhibition of HDAC6 reverses cognitive impairment and tau pathology as a result of cisplatin treatment. <i>Acta Neuropathologica Communications</i> , 2018, 6, 103.	5.2	44
30	Transition to chronic pain: opportunities for novel therapeutics. <i>Nature Reviews Neuroscience</i> , 2018, 19, 383-384.	10.2	113
31	Low GRK2 Underlies Hyperalgesic Priming by Glial Cell-Derived Neurotrophic Factor. <i>Frontiers in Pharmacology</i> , 2018, 9, 592.	3.5	14
32	Resolution of inflammation-induced depression requires T lymphocytes and endogenous brain interleukin-10 signaling. <i>Neuropsychopharmacology</i> , 2018, 43, 2597-2605.	5.4	83
33	Associations of inflammation with symptom burden in patients with acute myeloid leukemia. <i>Psychoneuroendocrinology</i> , 2018, 89, 203-208.	2.7	10
34	Identification of FAM173B as a protein methyltransferase promoting chronic pain. <i>PLoS Biology</i> , 2018, 16, e2003452.	5.6	22
35	Patient-reported fatigue prior to treatment is prognostic of survival in patients with acute myeloid leukemia. <i>Oncotarget</i> , 2018, 9, 31244-31252.	1.8	17
36	HDAC6 inhibition effectively reverses chemotherapy-induced peripheral neuropathy. <i>Pain</i> , 2017, 158, 1126-1137.	4.2	136

#	ARTICLE	IF	CITATIONS
37	Upregulation of neuronal kynurenine 3-monooxygenase mediates depression-like behavior in a mouse model of neuropathic pain. <i>Brain, Behavior, and Immunity</i> , 2017, 66, 94-102.	4.1	60
38	Pifithrin- α Prevents Cisplatin-Induced Chemobrain by Preserving Neuronal Mitochondrial Function. <i>Cancer Research</i> , 2017, 77, 742-752.	0.9	89
39	Inhibition of Mitochondrial p53 Accumulation by PFT- α Prevents Cisplatin-Induced Peripheral Neuropathy. <i>Frontiers in Molecular Neuroscience</i> , 2017, 10, 108.	2.9	68
40	Metformin Prevents Cisplatin-Induced Cognitive Impairment and Brain Damage in Mice. <i>PLoS ONE</i> , 2016, 11, e0151890.	2.5	108
41	Critical role for Epac1 in inflammatory pain controlled by GRK2-mediated phosphorylation of Epac1. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2016, 113, 3036-3041.	7.1	104
42	IL-10 Fusion Protein Is a Novel Drug to Treat Persistent Inflammatory Pain. <i>Journal of Neuroscience</i> , 2016, 36, 7353-7363.	3.6	67
43	Epac1 interacts with importin β 1 and controls neurite outgrowth independently of cAMP and Rap1. <i>Scientific Reports</i> , 2016, 6, 36370.	3.3	13
44	CD8 ⁺ T Cells and Endogenous IL-10 Are Required for Resolution of Chemotherapy-Induced Neuropathic Pain. <i>Journal of Neuroscience</i> , 2016, 36, 11074-11083.	3.6	164
45	Dorsal Root Ganglion Infiltration by Macrophages Contributes to Paclitaxel Chemotherapy-Induced Peripheral Neuropathy. <i>Journal of Pain</i> , 2016, 17, 775-786.	1.4	237
46	Critical Role of GRK2 in the Prevention of Chronic Pain. <i>Methods in Pharmacology and Toxicology</i> , 2016, , 187-213.	0.2	0
47	Prevention of chemotherapy-induced peripheral neuropathy by the small-molecule inhibitor pifithrin- α . <i>Pain</i> , 2015, 156, 2184-2192.	4.2	60
48	Cytokine production as a putative biological mechanism underlying stress sensitization in high combat exposed soldiers. <i>Psychoneuroendocrinology</i> , 2015, 51, 534-546.	2.7	31
49	Peripheral indoleamine 2,3-dioxygenase 1 is required for comorbid depression-like behavior but does not contribute to neuropathic pain in mice. <i>Brain, Behavior, and Immunity</i> , 2015, 46, 147-153.	4.1	40
50	Reversal of diet-induced obesity and insulin resistance by inducible genetic ablation of GRK2. <i>Science Signaling</i> , 2015, 8, ra73.	3.6	56
51	Pre-deployment differences in glucocorticoid sensitivity of leukocytes in soldiers developing symptoms of PTSD, depression or fatigue persist after return from military deployment. <i>Psychoneuroendocrinology</i> , 2015, 51, 513-524.	2.7	21
52	Mechanisms of chemotherapy-induced behavioral toxicities. <i>Frontiers in Neuroscience</i> , 2015, 9, 131.	2.8	133
53	Assessment of long-term safety and efficacy of intranasal mesenchymal stem cell treatment for neonatal brain injury in the mouse. <i>Pediatric Research</i> , 2015, 78, 520-526.	2.3	74
54	Neonatal glucocorticoid treatment: Long-term effects on the hypothalamus-pituitary-adrenal axis, immune system, and problem behavior in 14-17 year old adolescents. <i>Brain, Behavior, and Immunity</i> , 2015, 45, 128-138.	4.1	28

#	ARTICLE	IF	CITATIONS
55	The Anti-Diabetic Drug Metformin Protects against Chemotherapy-Induced Peripheral Neuropathy in a Mouse Model. PLoS ONE, 2014, 9, e100701.	2.5	132
56	Monocytes/Macrophages Control Resolution of Transient Inflammatory Pain. Journal of Pain, 2014, 15, 496-506.	1.4	98
57	The neuroimmune basis of fatigue. Trends in Neurosciences, 2014, 37, 39-46.	8.6	254
58	Intranasally administered mesenchymal stem cells promote a regenerative niche for repair of neonatal ischemic brain injury. Experimental Neurology, 2014, 261, 53-64.	4.1	132
59	Balancing GRK2 and EPAC1 levels prevents and relieves chronic pain. Journal of Clinical Investigation, 2013, 123, 5023-5034.	8.2	83
60	MicroRNA-124 as a novel treatment for persistent hyperalgesia. Journal of Neuroinflammation, 2012, 9, 143.	7.2	129
61	Microglial GRK2: A novel regulator of transition from acute to chronic pain. Brain, Behavior, and Immunity, 2011, 25, 1055-1060.	4.1	24
62	GRK2 in sensory neurons regulates epinephrine-induced signalling and duration of mechanical hyperalgesia. Pain, 2011, 152, 1649-1658.	4.2	43
63	Microglial/macrophage GRK2 determines duration of peripheral IL-1 β -induced hyperalgesia: Contribution of spinal cord CX3CR1, p38 and IL-1 signaling. Pain, 2010, 150, 550-560.	4.2	85
64	Low Nociceptor GRK2 Prolongs Prostaglandin E ₂ Hyperalgesia via Biased cAMP Signaling to Epac/Rap1, Protein Kinase C μ , and MEK/ERK. Journal of Neuroscience, 2010, 30, 12806-12815.	3.6	85
65	Nasal administration of stem cells: a promising novel route to treat neonatal ischemic brain damage. Pediatric Research, 2010, 68, 1.	2.3	96
66	Cell-specific roles of GRK2 in onset and severity of hypoxic-ischemic brain damage in neonatal mice. Brain, Behavior, and Immunity, 2010, 24, 420-426.	4.1	31
67	GRK2: A Novel Cell-Specific Regulator of Severity and Duration of Inflammatory Pain. Journal of Neuroscience, 2010, 30, 2138-2149.	3.6	103
68	A role for G protein-coupled receptor kinase β 2 in mechanical allodynia. European Journal of Neuroscience, 2007, 25, 1696-1704.	2.6	37
69	Stress, genetics, and immunity. Brain, Behavior, and Immunity, 2006, 20, 313-316.	4.1	6
70	Changes in innate and acquired immune responses in mice with targeted deletion of the dopamine transporter gene. Journal of Neuroimmunology, 2005, 161, 162-168.	2.3	51
71	G Protein-Coupled Receptor Kinase 2 in Multiple Sclerosis and Experimental Autoimmune Encephalomyelitis. Journal of Immunology, 2005, 174, 4400-4406.	0.8	105
72	Increased Acute Inflammation, Leukotriene B ₄ -Induced Chemotaxis, and Signaling in Mice Deficient for G Protein-Coupled Receptor Kinase 6. Journal of Immunology, 2003, 171, 6128-6134.	0.8	64

#	ARTICLE	IF	CITATIONS
73	Regulated expression of α -1 adrenergic receptors in the immune system. <i>Brain, Behavior, and Immunity</i> , 2002, 16, 799-807.	4.1	98
74	Decreased expression and activity of G-protein-coupled receptor kinases in peripheral blood mononuclear cells of patients with rheumatoid arthritis. <i>FASEB Journal</i> , 1999, 13, 715-725.	0.5	200
75	Role of endogenous pro-enkephalin A-derived peptides in human T cell proliferation and monocyte IL-6 production. <i>Journal of Neuroimmunology</i> , 1998, 84, 53-60.	2.3	53
76	beta-Endorphin: Cytokine and Neuropeptide. <i>Immunological Reviews</i> , 1991, 119, 41-63.	6.0	148
77	Induction of β -Endorphin Secretion by Lymphocytes after Subcutaneous Administration of Corticotropin-Releasing Factor. <i>Endocrinology</i> , 1990, 126, 759-764.	2.8	75