

# Josã© Tadeu Abreu de Oliveira

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/7758763/publications.pdf>

Version: 2024-02-01

87  
papers

2,249  
citations

218381

26  
h-index

253896

43  
g-index

88  
all docs

88  
docs citations

88  
times ranked

2383  
citing authors

#	ARTICLE	IF	CITATIONS
1	Antinutritional properties of plant lectins. <i>Toxicon</i> , 2004, 44, 385-403.	0.8	295
2	Proteomic analysis of the reproductive tract fluids from tropically-adapted Santa Ines rams. <i>Journal of Proteomics</i> , 2012, 75, 4436-4456.	1.2	83
3	Antifungal activity of plant and bacterial ureases. <i>Toxicon</i> , 2007, 50, 971-983.	0.8	72
4	Synthetic antimicrobial peptides: From choice of the best sequences to action mechanisms. <i>Biochimie</i> , 2020, 175, 132-145.	1.3	71
5	Composition and nutritional properties of seeds from <i>Pachira aquatica</i> Aubl, <i>Sterculia striata</i> St Hil et Naud and <i>Terminalia catappa</i> Linn. <i>Food Chemistry</i> , 2000, 70, 185-191.	4.2	70
6	An antifungal peptide from passion fruit ( <i>Passiflora edulis</i> ) seeds with similarities to 2S albumin proteins. <i>Biochimica Et Biophysica Acta - Proteins and Proteomics</i> , 2006, 1764, 1141-1146.	1.1	64
7	Synthetic antimicrobial peptides: Characteristics, design, and potential as alternative molecules to overcome microbial resistance. <i>Life Sciences</i> , 2021, 278, 119647.	2.0	64
8	Composition, toxic and antinutritional factors of newly developed cultivars of Brazilian soybean ( <i>Glycine max</i> ). <i>Journal of the Science of Food and Agriculture</i> , 1997, 75, 419-426.	1.7	63
9	Mo-CBP3-PepI, Mo-CBP3-PepII, and Mo-CBP3-PepIII are synthetic antimicrobial peptides active against human pathogens by stimulating ROS generation and increasing plasma membrane permeability. <i>Biochimie</i> , 2019, 157, 10-21.	1.3	57
10	A Protein Isolate from <i>Moringa oleifera</i> Leaves Has Hypoglycemic and Antioxidant Effects in Alloxan-Induced Diabetic Mice. <i>Molecules</i> , 2017, 22, 271.	1.7	50
11	DROUGHT-INDUCED EFFECTS AND RECOVERY OF NITRATE ASSIMILATION AND NODULE ACTIVITY IN COWPEA PLANTS INOCULATED WITH BRADYRHIZOBIUM SPP. UNDER MODERATE NITRATE LEVEL. <i>Brazilian Journal of Microbiology</i> , 2001, 32, 187-194.	0.8	49
12	A novel chitin-binding protein from <i>Moringa oleifera</i> seed with potential for plant disease control. <i>Biopolymers</i> , 2012, 98, 406-415.	1.2	48
13	Mo-CBP3, an Antifungal Chitin-Binding Protein from <i>Moringa oleifera</i> Seeds, Is a Member of the 2S Albumin Family. <i>PLoS ONE</i> , 2015, 10, e0119871.	1.1	47
14	Differential expression of antioxidant enzymes and PR-proteins in compatible and incompatible interactions of cowpea ( <i>Vigna unguiculata</i> ) and the root-knot nematode <i>Meloidogyne incognita</i> . <i>Plant Physiology and Biochemistry</i> , 2012, 51, 145-152.	2.8	45
15	Nutritional study of two Brazilian soybean ( <i>Glycine max</i> ) cultivars differing in the contents of antinutritional and toxic proteins. <i>Journal of Nutritional Biochemistry</i> , 2001, 12, 55-62.	1.9	44
16	New Insights into the Structure and Mode of Action of Mo-CBP3, an Antifungal Chitin-Binding Protein of <i>Moringa oleifera</i> Seeds. <i>PLoS ONE</i> , 2014, 9, e111427.	1.1	43
17	Induction of an anionic peroxidase in cowpea leaves by exogenous salicylic acid. <i>Journal of Plant Physiology</i> , 2006, 163, 1040-1048.	1.6	42
18	Effects of a Novel Pathogenesis-Related Class 10 (PR-10) Protein from <i>Crotalaria pallida</i> Roots with Papain Inhibitory Activity against Root-Knot Nematode <i>Meloidogyne incognita</i> . <i>Journal of Agricultural and Food Chemistry</i> , 2010, 58, 4145-4152.	2.4	42

#	ARTICLE	IF	CITATIONS
19	JcTI: a novel trypsin inhibitor from <i>Jatropha curcas</i> seed cake with potential for bacterial infection treatment. <i>Frontiers in Microbiology</i> , 2014, 5, 5.	1.5	42
20	PURIFICATION AND PARTIAL CHARACTERIZATION OF A LECTIN FROM THE SEEDS OF <i>DIOCLEA GUIANENSIS</i> . <i>Journal of Food Biochemistry</i> , 1991, 15, 137-154.	1.2	41
21	A Chitin-binding Protein Purified from <i>Moringa oleifera</i> Seeds Presents Anticandidal Activity by Increasing Cell Membrane Permeability and Reactive Oxygen Species Production. <i>Frontiers in Microbiology</i> , 2017, 8, 980.	1.5	41
22	RcAlb-PepII, a synthetic small peptide bioinspired in the 2S albumin from the seed cake of <i>Ricinus communis</i> , is a potent antimicrobial agent against <i>Klebsiella pneumoniae</i> and <i>Candida parapsilosis</i> . <i>Biochimica Et Biophysica Acta - Biomembranes</i> , 2020, 1862, 183092.	1.4	38
23	A molecular docking study revealed that synthetic peptides induced conformational changes in the structure of SARS-CoV-2 spike glycoprotein, disrupting the interaction with human ACE2 receptor. <i>International Journal of Biological Macromolecules</i> , 2020, 164, 66-76.	3.6	38
24	Purification and physicochemical characterization of a cotyledonary lectin from <i>Luetzelburgia auriculata</i> . <i>Phytochemistry</i> , 2002, 61, 301-310.	1.4	32
25	A trypsin inhibitor purified from <i>Cassia leiandra</i> seeds has insecticidal activity against <i>Aedes aegypti</i> . <i>Process Biochemistry</i> , 2017, 57, 228-238.	1.8	31
26	Production in <i>Pichia pastoris</i> , antifungal activity and crystal structure of a class I chitinase from cowpea ( <i>Vigna unguiculata</i> ): Insights into sugar binding mode and hydrolytic action. <i>Biochimie</i> , 2017, 135, 89-103.	1.3	28
27	Photosynthetic and biochemical mechanisms of an EMS-mutagenized cowpea associated with its resistance to cowpea severe mosaic virus. <i>Plant Cell Reports</i> , 2017, 36, 219-234.	2.8	28
28	<i>Canavalia brasiliensis</i> seeds. Protein quality and nutritional implications of dietary lectin. <i>Journal of the Science of Food and Agriculture</i> , 1994, 64, 417-424.	1.7	25
29	Proteome of Soybean Seed Exudates Contains Plant Defense-Related Proteins Active against the Root-Knot Nematode <i>Meloidogyne incognita</i> . <i>Journal of Agricultural and Food Chemistry</i> , 2015, 63, 5335-5343.	2.4	24
30	Castor bean cake contains a trypsin inhibitor that displays antifungal activity against <i>Colletotrichum gloeosporioides</i> and inhibits the midgut proteases of the dengue mosquito larvae. <i>Industrial Crops and Products</i> , 2015, 70, 48-55.	2.5	24
31	H <sub>2</sub> O <sub>2</sub> plays an important role in the lifestyle of <i>Colletotrichum gloeosporioides</i> during interaction with cowpea [ <i>Vigna unguiculata</i> (L.) Walp.]. <i>Fungal Biology</i> , 2015, 119, 747-757.	1.1	23
32	Bactericidal Activity Identified in 2S Albumin from Sesame Seeds and In silico Studies of Structure-Function Relations. <i>Protein Journal</i> , 2011, 30, 340-350.	0.7	22
33	Purification of a Chitin-Binding Protein from <i>Moringa oleifera</i> Seeds with Potential to Relieve Pain and Inflammation. <i>Protein and Peptide Letters</i> , 2011, 18, 1078-1085.	0.4	21
34	Synthetic antimicrobial peptides control <i>Penicillium digitatum</i> infection in orange fruits. <i>Food Research International</i> , 2021, 147, 110582.	2.9	21
35	Insulin-like plant proteins as potential innovative drugs to treat diabetes—The <i>Moringa oleifera</i> case study. <i>New Biotechnology</i> , 2017, 39, 99-109.	2.4	19
36	Mo-CBP4, a purified chitin-binding protein from <i>Moringa oleifera</i> seeds, is a potent antidermatophytic protein: In vitro mechanisms of action, in vivo effect against infection, and clinical application as a hydrogel for skin infection. <i>International Journal of Biological Macromolecules</i> , 2020, 149, 432-442.	3.6	19

#	ARTICLE	IF	CITATIONS
37	A ConA-like Lectin from <i>Dioclea guianensis</i> Benth. Has Antifungal Activity against <i>Colletotrichum gloeosporioides</i> , unlike Its Homologues, ConM and ConA. Journal of Agricultural and Food Chemistry, 2010, 58, 4090-4096.	2.4	18
38	Proteomic analysis of responsive stem proteins of resistant and susceptible cashew plants after <i>Lasiodiplodia theobromae</i> infection. Journal of Proteomics, 2015, 113, 90-109.	1.2	18
39	A 2S Albumin from the Seed Cake of <i>Ricinus communis</i> Inhibits Trypsin and Has Strong Antibacterial Activity against Human Pathogenic Bacteria. Journal of Natural Products, 2016, 79, 2423-2431.	1.5	18
40	Label-free Proteomic Reveals that Cowpea Severe Mosaic Virus Transiently Suppresses the Host Leaf Protein Accumulation During the Compatible Interaction with Cowpea ( <i>Vigna unguiculata</i> [L.] Tj ETQq0 0 0rgBT /Overlock 10 T		
41	CITI, a Kunitz trypsin inhibitor purified from <i>Cassia leiandra</i> Benth. seeds, exerts a candidicidal effect on <i>Candida albicans</i> by inducing oxidative stress and necrosis. Biochimica Et Biophysica Acta - Biomembranes, 2019, 1861, 183032.	1.4	18
42	A resistant cowpea ( <i>Vigna unguiculata</i> [L.] Walp.) genotype became susceptible to cowpea severe mosaic virus (CPSMV) after exposure to salt stress. Journal of Proteomics, 2019, 194, 200-217.	1.2	18
43	Gel-free/label-free proteomic, photosynthetic, and biochemical analysis of cowpea ( <i>Vigna unguiculata</i> ) Tj ETQq1 1 0.784314 rgBT /Overlock 10 T 76-91.	1.2	17
44	Anticandidal activity of synthetic peptides: Mechanism of action revealed by scanning electron and fluorescence microscopies and synergism effect with nystatin. Journal of Peptide Science, 2020, 26, e3249.	0.8	17
45	Soybean Toxin (SBTX), a Protein from Soybeans That Inhibits the Life Cycle of Plant and Human Pathogenic Fungi. Journal of Agricultural and Food Chemistry, 2010, 58, 10356-10363.	2.4	15
46	Cowpea-Meloidogyne incognita interaction: Root proteomic analysis during early stages of nematode infection. Proteomics, 2015, 15, 1746-1759.	1.3	15
47	A Bowman-Birk Inhibitor from the Seeds of <i>Luetzelburgia auriculata</i> Inhibits <i>Staphylococcus aureus</i> Growth by Promoting Severe Cell Membrane Damage. Journal of Natural Products, 2018, 81, 1497-1507.	1.5	15
48	Myracrodruon urundeuva seed exudates proteome and anthelmintic activity against <i>Haemonchus contortus</i> . PLoS ONE, 2018, 13, e0200848.	1.1	15
49	Scanning electron microscopy reveals deleterious effects of <i>Moringa oleifera</i> seed exuded proteins on root-knot nematode <i>Meloidogyne incognita</i> eggs. International Journal of Biological Macromolecules, 2020, 154, 1237-1244.	3.6	15
50	Gene expression and spatiotemporal localization of antifungal chitin-binding proteins during <i>Moringa oleifera</i> seed development and germination. Planta, 2019, 249, 1503-1519.	1.6	13
51	Role of membrane sterol and redox system in the anti-candida activity reported for Mo-CBP2, a protein from <i>Moringa oleifera</i> seeds. International Journal of Biological Macromolecules, 2020, 143, 814-824.	3.6	13
52	<i>Cratylia argentea</i> seed lectin, a possible defensive protein against plant-eating organisms: effects on rat metabolism and gut histology. Food and Chemical Toxicology, 2004, 42, 1737-1747.	1.8	12
53	A cotyledonary agglutinin from <i>Luetzelburgia auriculata</i> inhibits the fungal growth of <i>Colletotrichum lindemuthianum</i> , <i>Fusarium solani</i> and <i>Aspergillus niger</i> and impairs glucose-stimulated acidification of the incubation medium by <i>Saccharomyces cerevisiae</i> cells. Plant Science, 2005, 169, 629-639.	1.7	12
54	Enhanced Synthesis of Antioxidant Enzymes, Defense Proteins and Leghemoglobin in Rhizobium-Free Cowpea Roots after Challenging with <i>Meloidogyne incognita</i> . Proteomes, 2014, 2, 527-549.	1.7	12

#	ARTICLE	IF	CITATIONS
55	Proteomics changes during the incompatible interaction between cowpea and <i>Colletotrichum gloeosporioides</i> (Penz.) Penz and Sacc. <i>Plant Science</i> , 2014, 217-218, 158-175.	1.7	12
56	Drought increases cowpea ( <i>Vigna unguiculata</i> [L.] Walp.) susceptibility to cowpea severe mosaic virus (CPSMV) at early stage of infection. <i>Plant Physiology and Biochemistry</i> , 2016, 109, 91-102.	2.8	12
57	A cysteine protease from the latex of <i>Ficus benjamina</i> has in vitro anthelmintic activity against <i>Haemonchus contortus</i> . <i>Brazilian Journal of Veterinary Parasitology</i> , 2018, 27, 473-480.	0.2	12
58	<i>Machaerium acutifolium</i> lectin alters membrane structure and induces ROS production in <i>Candida parapsilosis</i> . <i>International Journal of Biological Macromolecules</i> , 2020, 163, 19-25.	3.6	11
59	ClCPI, a cysteine protease inhibitor purified from <i>Cassia leiandra</i> seeds has antifungal activity against <i>Candida tropicalis</i> by inducing disruption of the cell surface. <i>International Journal of Biological Macromolecules</i> , 2019, 133, 1115-1124.	3.6	10
60	Antidermatophytic activity of synthetic peptides: Action mechanisms and clinical application as adjuvants to enhance the activity and decrease the toxicity of Griseofulvin. <i>Mycoses</i> , 2020, 63, 979-992.	1.8	10
61	SBTX, a new toxic protein distinct from soyatoxin and other toxic soybean [ <i>Glycine max</i> ] proteins, and its inhibitory effect on <i>Cercospora sojina</i> growth. <i>Toxicon</i> , 2008, 51, 952-963.	0.8	9
62	ACE2-derived peptides interact with the RBD domain of SARS-CoV-2 spike glycoprotein, disrupting the interaction with the human ACE2 receptor. <i>Journal of Biomolecular Structure and Dynamics</i> , 2022, 40, 5493-5506.	2.0	9
63	Reference gene identification for real-time PCR analyses in soybean leaves under fungus ( <i>Cercospora</i> ) Tj ETQq1 1 0.784314 rgBT /Ove 2016, 45, 191-199.	0.5	8
64	Identification, characterization, and expression analysis of cowpea ( <i>Vigna unguiculata</i> [L.] Walp.) miRNAs in response to cowpea severe mosaic virus (CPSMV) challenge. <i>Plant Cell Reports</i> , 2020, 39, 1061-1078.	2.8	8
65	Synthetic peptides against <i>Trichophyton mentagrophytes</i> and <i>T. rubrum</i> : Mechanisms of action and efficiency compared to griseofulvin and itraconazole. <i>Life Sciences</i> , 2021, 265, 118803.	2.0	8
66	Quantum biochemistry, molecular docking, and dynamics simulation revealed synthetic peptides induced conformational changes affecting the topology of the catalytic site of SARS-CoV-2 main protease. <i>Journal of Biomolecular Structure and Dynamics</i> , 2022, 40, 8925-8937.	2.0	8
67	Towards a better understanding of <i>Ipomoea asarifolia</i> toxicity: Evidence of the involvement of a leaf lectin. <i>Toxicon</i> , 2011, 58, 502-508.	0.8	7
68	A protein extract and a cysteine protease inhibitor enriched fraction from <i>Jatropha curcas</i> seed cake have in vitro anti- <i>Toxoplasma gondii</i> activity. <i>Experimental Parasitology</i> , 2015, 153, 111-117.	0.5	7
69	The expression of the genes involved in redox metabolism and hydrogen peroxide balance is associated with the resistance of cowpea [ <i>Vigna unguiculata</i> (L.) Walp.] to the hemibiotrophic fungus <i>Colletotrichum gloeosporioides</i> . <i>Journal of Plant Physiology</i> , 2019, 233, 73-83.	1.6	7
70	Neutralizing Effect of Synthetic Peptides toward SARS-CoV-2. <i>ACS Omega</i> , 2022, 7, 16222-16234.	1.6	7
71	Soybean Toxin (SBTX) Impairs Fungal Growth by Interfering with Molecular Transport, Carbohydrate/Amino Acid Metabolism and Drug/Stress Responses. <i>PLoS ONE</i> , 2013, 8, e70425.	1.1	6
72	Mo-CBP3, a 2S albumin from <i>Moringa oleifera</i> , is a complex mixture of isoforms that arise from different post-translational modifications. <i>Plant Physiology and Biochemistry</i> , 2019, 140, 68-77.	2.8	6

#	ARTICLE	IF	CITATIONS
73	H <sub>2</sub> O <sub>2</sub> Accumulation, Host Cell Death and Differential Levels of Proteins Related to Photosynthesis, Redox Homeostasis, and Required for Viral Replication Explain the Resistance of EMS-mutagenized Cowpea to Cowpea Severe Mosaic Virus. <i>Journal of Plant Physiology</i> , 2020, 245, 153110.	1.6	6
74	Computational approach, scanning electron and fluorescence microscopies revealed insights into the action mechanisms of anticandidal peptide Mo-CBP3-PepIII. <i>Life Sciences</i> , 2021, 281, 119775.	2.0	6
75	Orally hypoglycemic activity of an insulin mimetic glycoprotein isolated from <i>Cnidocolus quercifolius</i> Pohl. (Euphorbiaceae) seeds, Cq-IMP. <i>International Journal of Biological Macromolecules</i> , 2020, 159, 886-895.	3.6	5
76	A peroxidase purified from cowpea roots possesses high thermal stability and displays antifungal activity against <i>Colletotrichum gloeosporioides</i> and <i>Fusarium oxysporum</i> . <i>Biocatalysis and Agricultural Biotechnology</i> , 2022, 42, 102322.	1.5	5
77	Synergistic Antifungal Activity of Synthetic Peptides and Antifungal Drugs against <i>Candida albicans</i> and <i>C. parapsilosis</i> Biofilms. <i>Antibiotics</i> , 2022, 11, 553.	1.5	5
78	In vitro and in vivo digestibility of the albumin and globulin fractions of eight Brazilian cowpea [ <i>Vigna unguiculata</i> (L) Walp] cultivars. <i>Journal of the Science of Food and Agriculture</i> , 2004, 84, 1823-1830.	1.7	4
79	Increased Levels of Antinutritional and/or Defense Proteins Reduced the Protein Quality of a Disease-Resistant Soybean Cultivar. <i>Nutrients</i> , 2015, 7, 6038-6054.	1.7	4
80	New Insights into Anthelmintic Mechanisms of Action of a Synthetic Peptide: An Ultrastructural and Nanomechanical Approach. <i>Polymers</i> , 2021, 13, 2370.	2.0	4
81	Biochemical, physicochemical and molecular characterization of a genuine 2-Cys-peroxiredoxin purified from cowpea [ <i>Vigna unguiculata</i> (L.) Walpers] leaves. <i>Biochimica Et Biophysica Acta - General Subjects</i> , 2012, 1820, 1128-1140.	1.1	2
82	Purification, characterization and antifungal activity of LPA a lectin from <i>Pachira aquatica</i> seeds: their possible role in plant defence. <i>BMC Proceedings</i> , 2014, 8, .	1.8	2
83	Gene expression during development and overexpression after <i>Cercospora kikuchii</i> and salicylic acid challenging indicate defensive roles of the soybean toxin. <i>Plant Cell Reports</i> , 2020, 39, 669-682.	2.8	1
84	In vitro toxicological characterisation of the antifungal compound soybean toxin (SBTX). <i>Toxicology in Vitro</i> , 2020, 65, 104824.	1.1	1
85	Inhibition of Protease and Egg Hatching of <i>Haemonchus contortus</i> by Soybean Seed Exudates. <i>Journal of Parasitology</i> , 2021, 107, 23-28.	0.3	1
86	JcTI-PepI, a synthetic peptide bioinspired in the trypsin inhibitor from <i>Jatropha curcas</i> , presents potent inhibitory activity against <i>C.Âkrusei</i> , a neglected pathogen. <i>Biochimie</i> , 2022, 200, 107-118.	1.3	1
87	A chemically sulfated derivative galactomannan from <i>Adenanthera pavonina</i> seeds elicits defense-related responses in cowpea and confers protection against <i>Colletotrichum gloeosporioides</i> . <i>Journal of Biological Control</i> , 2018, 33, 7-17.	0.2	0