

# Abbas Imani-Baran

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/7758800/publications.pdf>

Version: 2024-02-01

9  
papers

107  
citations

1478505

6  
h-index

1474206

9  
g-index

9  
all docs

9  
docs citations

9  
times ranked

175  
citing authors

#	ARTICLE	IF	CITATIONS
1	Vairimorpha ceranae was the only detected microsporidian species from Iranian honey bee colonies: a molecular and phylogenetic study. Parasitology Research, 2022, 121, 355-366.	1.6	1
2	Anthelmintic activity of crude powder and crude aqueous extract of Trachyspermum ammi on gastrointestinal nematodes in donkey (Equus asinus): An in vivo study. Journal of Ethnopharmacology, 2020, 248, 112249.	4.1	8
3	In vitro acaricidal activity of essential oil and alcoholic extract of Trachyspermum ammi against Dermanyssus gallinae. Veterinary Parasitology, 2020, 278, 109030.	1.8	18
4	Evaluation of oxidative stress and antioxidant status, serum trace mineral levels and cholinesterases activity in cattle infected with Anaplasma marginale. Microbial Pathogenesis, 2018, 123, 402-409.	2.9	25
5	Semi-nested polymerase chain reaction-based detection of Babesia spp. in small ruminants from Northwest of Iran. Veterinary World, 2018, 11, 268-273.	1.7	2
6	Molecular Determination of . Isolates from Domestic Ruminants Fecal Samples in the Northwest of Iran. Iranian Journal of Parasitology, 2017, 12, 243-250.	0.6	6
7	Morphological and molecular discrimination of fasciola species isolated from domestic ruminants of urmia city, iran. Iranian Journal of Parasitology, 2015, 10, 46-55.	0.6	13
8	Seasonal and Geographic Distribution of Cercarial Infection in Lymnaea gedrosiana (Pulmonata: Lymnaeidae) in Northwestern Iran. Iranian Journal of Parasitology, 2014, 9, 10-11.	0.6	11
9	Molecular study for detecting the prevalence of Fasciola gigantica in field-collected snails of Radix gedrosiana (Pulmonata: Lymnaeidae) in northwestern Iran. Veterinary Parasitology, 2012, 189, 374-377.	1.8	23