

# Adelson Marcal Rodrigues

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/7764994/publications.pdf>

Version: 2024-02-01

16  
papers

285  
citations

1039406

9  
h-index

1058022

14  
g-index

16  
all docs

16  
docs citations

16  
times ranked

437  
citing authors

#	ARTICLE	IF	CITATIONS
1	Relation between red blood cell distribution width and acute kidney injury in patients with sepsis. Einstein (Sao Paulo, Brazil), 2022, 20, eAO6828.	0.3	5
2	Kefir reduces nitrosative stress and upregulates Nrf2 in the kidney of diabetic rats. International Dairy Journal, 2021, 114, 104909.	1.5	6
3	P2X7 receptor-nitric oxide interaction mediates apoptosis in mouse immortalized mesangial cells exposed to high glucose. Jornal Brasileiro De Nefrologia: Orgao Oficial De Sociedades Brasileira E Latino-Americana De Nefrologia, 2021, , .	0.4	1
4	P2X7 siRNA targeted to the kidneys increases klotho and delays the progression of experimental diabetic nephropathy. Purinergic Signalling, 2020, 16, 175-185.	1.1	7
5	Inhibition of the P2X7 receptor improves renal function via renin-angiotensin system and nitric oxide on diabetic nephropathy in rats. Life Sciences, 2020, 251, 117640.	2.0	8
6	Esculin reduces P2X7 and reverses mitochondrial dysfunction in the renal cortex of diabetic rats. Life Sciences, 2020, 254, 117787.	2.0	7
7	Cupuaçu extract reduces nitrosative stress and modulates inflammatory mediators in the kidneys of experimental diabetes. Clinical Nutrition, 2019, 38, 364-371.	2.3	15
8	P2X7 receptor and klotho expressions in diabetic nephropathy progression. Purinergic Signalling, 2018, 14, 167-176.	1.1	13
9	N-acetylcysteine protects against diabetic nephropathy through control of oxidative and nitrosative stress by recovery of nitric oxide in rats. Nitric Oxide - Biology and Chemistry, 2018, 78, 22-31.	1.2	32
10	Moderate aerobic exercise on the recovery phase of gentamicin-induced acute kidney injury in rats. Life Sciences, 2017, 169, 37-42.	2.0	12
11	Evaluation of P2X 7 Receptor Expression in the Kidney at the Early Phase of Experimental Diabetes Mellitus. Free Radical Biology and Medicine, 2017, 112, 151-152.	1.3	0
12	Immunomodulation and nitric oxide restoration by a probiotic and its activity in gut and peritoneal macrophages in diabetic rats. Clinical Nutrition, 2016, 35, 1066-1072.	2.3	25
13	P2X7 Receptor in the Kidneys of Diabetic Rats Submitted to Aerobic Training or to N-Acetylcysteine Supplementation. PLoS ONE, 2014, 9, e97452.	1.1	32
14	Kefir administration reduced progression of renal injury in STZ-diabetic rats by lowering oxidative stress. Nitric Oxide - Biology and Chemistry, 2014, 37, 53-60.	1.2	53
15	Nitric oxide (NO) is associated with gentamicin (GENTA) nephrotoxicity and the renal function recovery after suspension of GENTA treatment in rats. Nitric Oxide - Biology and Chemistry, 2011, 24, 77-83.	1.2	40
16	Effects of training and nitric oxide on diabetic nephropathy progression in type I diabetic rats. Experimental Biology and Medicine, 2011, 236, 1180-1187.	1.1	29