## Jos de Keijser

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/7769356/publications.pdf Version: 2024-02-01



LOS DE KEUSED

#	Article	IF	CITATIONS
1	Prolonged Grief Disorder, depression, and posttraumatic stress disorder are distinguishable syndromes. Journal of Affective Disorders, 2010, 125, 374-378.	4.1	148
2	Trajectories of grief, depression, and posttraumatic stress in disasterâ€bereaved people. Depression and Anxiety, 2020, 37, 35-44.	4.1	83
3	Further validation of the Traumatic Grief Inventory-Self Report (TGI-SR): A measure of persistent complex bereavement disorder and prolonged grief disorder. Death Studies, 2019, 43, 351-364.	2.7	76
4	Cognitive–behavioral variables mediate the impact of violent loss on post-loss psychopathology Psychological Trauma: Theory, Research, Practice, and Policy, 2015, 7, 382-390.	2.1	67
5	Prolonged grief, depression, and posttraumatic stress in disaster-bereaved individuals: latent class analysis. HA¶gre Utbildning, 2017, 8, 1298311.	3.0	65
6	Valid measurement of DSM-5 persistent complex bereavement disorder and DSM-5-TR and ICD-11 prolonged grief disorder: The Traumatic Grief Inventory-Self Report Plus (TGI-SR+). Comprehensive Psychiatry, 2022, 112, 152281.	3.1	58
7	Psychopathology Among Homicidally Bereaved Individuals. Trauma, Violence, and Abuse, 2015, 16, 70-80.	6.2	31
8	Grief rumination mediates the association between self-compassion and psychopathology in relatives of missing persons. HA¶gre Utbildning, 2017, 8, 1378052.	3.0	30
9	Toward a Better Understanding of Psychological Symptoms in People Confronted With the Disappearance of a Loved One: A Systematic Review. Trauma, Violence, and Abuse, 2019, 20, 287-302.	6.2	29
10	Evaluation of benefit to patients of training mental health professionals in suicide guidelines: Cluster randomised trial. British Journal of Psychiatry, 2016, 208, 477-483.	2.8	28
11	Patterns, Predictors, and Prognostic Validity of Persistent Complex Bereavement Disorder Symptoms in Recently Bereaved Adults. Journal of Nervous and Mental Disease, 2019, 207, 913-920.	1.0	25
12	DSM-5-TR prolonged grief disorder and DSM-5 posttraumatic stress disorder are related, yet distinct: confirmatory factor analyses in traumatically bereaved people. Högre Utbildning, 2021, 12, 1-14.	3.0	22
13	Remotely Delivered Cognitive Behavior Therapy for Disturbed Grief During the COVID-19 Crisis: Challenges and Opportunities. Journal of Loss and Trauma, 2021, 26, 211-219.	1.5	21
14	Prolonged grief and post-traumatic stress among relatives of missing persons and homicidally bereaved individuals: A comparative study. Journal of Affective Disorders, 2017, 209, 1-2.	4.1	20
15	Treating complicated grief and posttraumatic stress in homicidally bereaved individuals: A randomized controlled trial. Clinical Psychology and Psychotherapy, 2018, 25, 497-508.	2.7	20
16	Reciprocal Associations Among Symptom Levels of Disturbed Grief, Posttraumatic Stress, and Depression Following Traumatic Loss: A Four-Wave Cross-Lagged Study. Clinical Psychological Science, 2019, 7, 1330-1339.	4.0	20
17	Treatment gap in bereavement care: (Online) bereavement support needs and use after traumatic loss. Clinical Psychology and Psychotherapy, 2021, 28, 907-916.	2.7	19
18	Revenge and psychological adjustment after homicidal loss. Aggressive Behavior, 2014, 40, 504-511.	2.4	18

Jos de Keijser

#	Article	IF	CITATIONS
19	Enhancing the effect of psychotherapy through systematic client feedback in outpatient mental healthcare: A cluster randomized trial Psychotherapy Research, 2022, 32, 710-722.	1.8	16
20	Cognitive-Behavioral Correlates of Psychological Symptoms Among Relatives of Missing Persons. International Journal of Cognitive Therapy, 2018, 11, 311-324.	2.2	11
21	Cognitive behavioural therapy and mindfulness for relatives of missing persons: a pilot study. Pilot and Feasibility Studies, 2019, 5, 93.	1.2	11
22	Online cognitive–behavioural therapy for traumatically bereaved people: study protocol for a randomised waitlist-controlled trial. BMJ Open, 2020, 10, e035050.	1.9	11
23	Online treatment of persistent complex bereavement disorder, posttraumatic stress disorder, and depression symptoms in people who lost loved ones during the COVID-19 pandemic: study protocol for a randomized controlled trial and a controlled trial. HĶgre Utbildning, 2021, 12, 1987687.	3.0	11
24	Cognitive behavioural therapy for psychopathology in relatives of missing persons: study protocol for a pilot randomised controlled trial. Pilot and Feasibility Studies, 2016, 2, 19.	1.2	9
25	Cognitive therapy and eye movement desensitization and reprocessing for reducing psychopathology among disaster-bereaved individuals: study protocol for a randomized controlled trial. HA¶gre Utbildning, 2017, 8, 1388710.	3.0	9
26	l've changed, but l'm not less happy: Interview study among nonclinical relatives of long-term missing persons. Death Studies, 2018, 42, 346-355.	2.7	9
27	Latent classes of posttraumatic stress symptoms in two samples of bereaved people International Journal of Stress Management, 2019, 26, 401-410.	1.2	7
28	Traumatic stress, depression, and non-bereavement grief following non-fatal traffic accidents: Symptom patterns and correlates. PLoS ONE, 2022, 17, e0264497.	2.5	6
29	Systematic client feedback to brief therapy in basic mental healthcare: study protocol for a four-centre clinical trial. BMJ Open, 2019, 9, e025701.	1.9	4
30	Associations of dimensions of anger with distress following traumatic bereavement Psychological Trauma: Theory, Research, Practice, and Policy, 2024, 16, 176-183.	2.1	4
31	Concurrent associations of dimensions of anger with posttraumatic stress, depression, and functional impairment following non-fatal traffic accidents. European Journal of Psychotraumatology, 2022, 13, 2068912.	2.5	0